

## The Comment

# Use of Pulse Dose Methylprednisolone in a Cohort of very Severe Covid-19 Patients in a resource-limited setting in Myanmar: a retrospective study

**Comment [A1]:** SEE THE FULL COMMENT AT THE OTHER FILES IN THIS ATTACHED

**Comment [A2]:** Writing the title is still too general, write a title more specific to your locus of research. See the full comment at the other files in this attached

## ABSTRACT

**Aims:** This study aims to explore the role of pulse dose methylprednisolone therapy in very severe COVID-19 patients in preventing the need for ICU care and death.

**Study design:** Retrospective record review study

**Place and Duration of Study:** Oak-ta-chat-thal-ta-pwint Covid-19 treatment center in Yangon, Myanmar between September 2021 to December 2021.

**Methodology:** This study included 13 confirmed Covid-19 patients with severe to critical illness, who were treated with pulse dose methylprednisolone therapy. We reviewed the patients' demographics, comorbidities and disease severity before starting pulse dose methylprednisolone therapy and changes in oxygen requirement, chest x ray scores, inflammatory markers, development of significant clinical events and 28 days mortality after therapy.

**Results:** Before pulse dose methylprednisolone therapy, all 13 patients had very severe disease (mean SPO<sub>2</sub>/FiO<sub>2</sub> = 173mmHg, mean SPO<sub>2</sub> = 88.54%, mean CRP = 115mg/L, mean ferritin = 1295.5ng/ml and mean Brixia Score = 6.54). They received 3-7 days (mean = 5.5) of pulse dose methylprednisolone. Ten patients (76%) survived in a setting with limited ICU care. High ferritin was a significant predictor of mortality. Improvement in oxygen requirement was noticeable after 1-11 days (mean = 5.6g). Hyperglycemia was common and confirmed bacterial infection was found in 3 patients, but all patients received empirical antibiotics therapy.

**Conclusion:** Pulse dose methylprednisolone therapy may be an effective salvage therapy in a carefully selected subset of very severe covid-19 patients. It might be a feasible alternative to other more expensive immunomodulating agents and organ support treatments in a resource-limited setting.

**Keywords:** Covid-19, Pulse dose corticosteroid, methylprednisolone, resource-limited settings

**Comment [A3]:** USE THE FOLLOWING STEPS IN DEVELOPING AN IDEAL ABSTRACT JOURNAL PERPUTATION (Glasman, 2010; EMERALD, 2020; Wibirama, 2019)

1. General Problem in Your Field
2. Specific Problem (Research Gap)
3. Your Proposed Solution/Proposed Method
4. Key Results/Key Findings
5. Brief Discussion and Implications

## 1. INTRODUCTION

Coronavirus disease 2019 (Covid-19) is a pandemic disease caused by a novel corona virus SARS-Cov-2. Recently, the world health organization declared an end to covid-19 as a disease of public health emergency. Yet, many mysteries remain about the disease. Since its emergence in 2019, more than six million lives have been lost to this disease [1]. Hypoxic respiratory failure due to ARDS and multisystem failure due to cytokine release syndromes are significant events in critical cases of covid-19. The immunopathology plays a major role in these critical stages of Covid-19 [2].

The use of low to moderate dose dexamethasone therapy in severe covid 19 has been well accepted [3]. In very severe cases, thrombo-inflammation in the lungs and alveolar damages progress despite low dose dexamethasone and antiviral agents, requiring ventilatory support and advanced ICU care. The prognoses in these patients are very poor despite every effort with the use of advanced ICU supports [4,5]. In a resource-limited setting, the advanced life supports are practically almost unavailable during a pandemic crisis and entering a critical stage in covid-19 is usually a one-way downward spiral to death. The pathogenesis in these later stages of disease is mainly due to aberrant and dysfunctional inflammatory response and suppressing the inflammatory response theoretically might benefit.

Corticosteroids are potent non-specific immunosuppressive and anti-inflammatory agents [6]. They are also widely available in every setting and their use has been familiar to most clinicians for their historical use in severe autoimmune conditions. Various doses of corticosteroids were reportedly used in severe covid-19 and the role of high dose corticosteroids is still uncertain. The studies on high dose corticosteroids in severe covid-19 were mainly retrospective observational studies and success rate has been varied due to the use of various doses of drug and choice of patients [7–19]. There are also concerns for potential harms like delay viral clearance and superimposed infections. Short term use of supra-physiological dose of corticosteroids is believed to act through nongenomic pathway to downregulate immune cells activation and proinflammatory cytokine production [20]. We hypothesized that the high dose corticosteroid would be beneficial if it is given in a critical time window at the start of severe unchecked inflammation.

This study aimed to explore the role of pulse dose methylprednisolone therapy in very severe COVID-19 patients in preventing the need for ICU care and death. The specific objectives were to describe the characteristics of patients treated with high-dose methylprednisolone therapy, ascertain the timing from disease onset and severity status at the initiation of pulse dose therapy, and describe the outcomes and complications of these patients.

## 2. METHODOLOGY

This is a retrospective medical record review study on a series of severe Covid-19 cases treated with pulse dose methylprednisolone therapy. The study was done in a temporary covid-19 treatment facility in Yangon, Myanmar. The study was conducted from September to December 2021, when the delta B.1.617.2 strain of SARS-Cov2 was actively transmitting in the country, coinciding with the later part of the catastrophic "third wave". All patients admitted to the facility during the study period were reviewed for eligibility. There was no sample size calculation and all patients who received the therapy of interest during the study period were included. All adult patients more than 18 years of age with confirmed Covid-19 disease and who received pulse dose methylprednisolone therapy (defined as MP  $\geq$  250mg or equivalent for at least 3 consecutive days by clinician discretion) were included in the review. The variables of interest were extracted from inpatient medical records of the electronic medical record systems. Outcomes data were extracted from the routine telephone follow-up record of the treatment facility until 28 days from date of admission. Mortality status (death or survive) was final outcome and receiving pulse dose methylprednisolone therapy (PDMPT) was exposure and changes in SPO<sub>2</sub>/FiO<sub>2</sub> status, Brixia Score before and after PDMPT and biochemical makers (CRP, ferritin and lymphocytes) were predictors; age and gender are potential confounders and days of covid 19 and underlying co-morbidity and use of other immunomodulators were effect modifiers.

The age, sex and underlying comorbid conditions were extracted from the initial clinical assessment record by the admitting medical officer. The dose and total duration of high-dose corticosteroid therapy was taken from the highest dose of corticosteroid which the

**Comment [A4]:** Preferred preliminary elements contain elements that clearly describe (in order): (1). A bit of general background is needed for this research; (2) state of the art (brief literature review) previous (similar) studies to justify the novelty of this article (there must be a reference to the journal within the last 5 years); (3) gap analysis or statement of gaps or novelty (novelty statement) or the unique difference of this research compared to previous research or based on state of the art (in terms of whether the research is important or the comparison of uniqueness with previous research); (4) problems and/or hypotheses (if any); (5) problem solving approach (if any); (6) the expected results or research objectives in this article.

Notes:

**The elements above have not been explained.**

patient received during admission and the duration of that dose, irrespective of the tapering doses or if-any previous smaller doses. The timing of therapy initiation was measured by days from onset of symptoms and severity of disease at the initiation was measured by oxygen requirement (SaO<sub>2</sub>/FiO<sub>2</sub> ratio estimated from dose of oxygen and pulse oxymetry measurement), chest Xray scores (Brixia score by Borghesi and Maroldi, 2020 [21]) and biochemical markers of inflammation. Any other treatments received were also recorded and they were grouped into antiviral agents, anti-thrombotic, other immunomodulators and antibiotics.

Any event after initiation of therapy until discharged was analyzed as potential complications of therapy. Infection was defined as any event of culture-confirmed or clinically suspected infection, requiring antibiotics therapy. Hyperglycemia was new onset or worsening hyperglycemia requiring any treatment or diet modifications. Clinical severity of disease was classified as asymptomatic, mild, moderate and severe or critical as defined by the Covid-19 treatment guideline by the National Institute of Health [22].

### 3. RESULTS AND DISCUSSION

Total of 13 patients received 3-7 days course of IV methylprednisolone therapy. Twelve patients received 1 G of methylprednisolone each day while only 1 patient was given 250mg of methylprednisolone therapy. Majority of them (n=11) were also given per oral baricitinib and 9 patients also received two doses of tocilizumab (according to availability). CRP level was the earliest to fall after high dose steroid therapy and improvement in hypoxia was noticeable only after 1-11 days with mean duration 5.6 days.

Table 1: Description of 13 Covid 19 cases under the study in accordance with their final outcomes

Variables	Outcomes		Total	Significant Test "p"	Remark
	Alive n(%)	Death n (%)			
<b>Age Group</b>				Fisher Exact Test $\chi^2 =$ 0.795 "p" = 1.00	
≤40 years	1(100)	0(0)	1(7.7)		
≥41-60 years	5(71.4)	2(28.6)	7 (53.8)		
>61 years	4 (80)	1(20)	5 (38.5)		
Minimum age – 27					
Maximum age -85					
Mean (±SD)- 56.63±14.86					
<b>Gender</b>				Fisher exact $\chi^2 = 5.424$	
Male	4 (57.1)	3 (42.9)	7 (53.8)		

Female	6(100)	0(0)	6 (46.2)	"P" = 0.192	
<b>Comorbidity *</b>				X <sup>2</sup> = 0.231	
Yes	8(80)	2 (20)	10(79.9*)	"P" = 1.00	
No	2 (66.7)	1 (33.3 )	3(23.1*)		
*Diabetes,Hypertension, IHD, Obesity,Peripheral Vaculopathy					
<b>Risk for Severe Covid</b>				X <sup>2</sup> = 0.231	
<b>Yes</b> (≥61 years± underlying d/s)	8 (80)	2 (20)	10(76.9*)	"P" = 1.00	
<b>No</b> (≤ 60 without underlying d/s)	2(66.6)	1 (33.3)	3 (23.1*)		
<b>Steroid duration (days)</b>				95% CI of the difference	
Minimum days	3	5		lower (-1.19)	
Maximum days	7	5		upper(2.93)	
Mean±SD	5.5(±1.84 )	5(±0.0)			
<b>Days from start of steroid to improve in hypoxia</b>				95% CI of the difference	
Minimum days	1	4		lower (-6.09)	
Maximum days	10	11		upper(4.22)	
Mean±SD	5.4(±3.41 )	6.33(±4.0 4)			
<b>Additional immunosuppressive</b>				X <sup>2</sup> = 1.501	
<b>Baricitinib</b>	3(100)	0 (0)	3 (23.1)	"P" = 0.27	
<b>Nil</b>	1(100)	0 (0)	1 (7.7)		
<b>Tocilizumab, Baricitinib</b>	6(66.6)	3 (33.4)	9 (69.2)		
<b>Symptoms days before admission</b>				X <sup>2</sup> = 0.258	
<b>&lt;7 days</b>	5 (83.3)	1(16.7)	6(46.2)	"P" = 1.000	
<b>≥7 days</b>	5 (71.4)	2(28.6)	7 (53.8)		

Table 2. Comparison of disease severity status before PDMP therapy of 13 Covid-19 cases under study in accordance with final outcomes

Disease severity parameters	Outcomes		Total	Significant Test "p"	Remark
	Alive	Death			
<b>SPO2/FiO2 Before Steroid Therapy</b>					
Minimum	96	100	96	95% CI of the difference lower (-130.06) upper(79.59) "P"= 0.61	
Maximum	303	240	303		
Mean (±SE)	168 (22.2)	193(±46.67)	173(±19.46)		
			6		
<b>CRP Before Steroid Therapy</b>					
Minimum	44.3	68.1	44.3	95% CI of the difference lower (-96.74) upper(112.13) "P"= 0.87	
Maximum	264.8	151.5	264.8		
Mean (±SE)	117(±24.4)	109.9(±46.67)	115(±10.16)		
<b>Lymphocytes Before Steroid Therapy</b>					
Minimum	0.37	0.63	0.37	95% CI of the difference lower (-0.76) upper(0.97)	
Maximum	2.47	1.05	2.47		
Mean (±SE)	0.96(±0.21)	0.86(±0.12)	0.93(±0.57)		
<b>Ferritin Before Steroid Therapy</b>					
Minimum	204	1392	204	95% CI of the difference lower (-2854.74) upper(455.07))	
Maximum	1902	3001	3001		
Mean (±SE)	944.76(±216.42)	2464.67(±536.3)	1295.5(±268.8)		
<b>Lowest SPO2 at Home</b>					
Minimum days	75	88	75	95% CI of the difference lower (-8.7) upper(5.77)	
Maximum days	93	92	93		
Mean±SD	88.2(±1.718)	89.67(±1.2)	88.54(±1.33)		

Table 3. Case Summary of 13 severe Covid-19 patients treated with pulse dose methylprednisolone therapy

Cas e No	A g e	Sex	Comorbid ities	SP O 2/ Fi O 2	CRP	Lymp hocyt e	Ferr itin	Bri xia sc or e	Ster oid dose	Ster oid dura tion	Additional immunos uppressiv es	Events after treatment	Final Outco me
1	60	F	Peripheral vascular disease	95.83	59.86	0.41	3267	7	1G	7	tocilizum ab, baricitini b	Hyperglyc emia	surviv e
2	43	F	Nil	101.04	73.23	1.01	204	10	1G	7	tocilizum ab, baricitini b	Hypergly cemia, Bacterial pneumo nia	surviv e
3	72	F	Hyperten sion, cardiova scular disease	194.00	264.76	1.09	411.3	2	250 mg	5	tocilizum ab, baricitini b	Hypergly cemia, Gl bleeding	surviv e
4	42	M	obesity, hyperten sion	98.96	108.9	0.37	1621	10	1G	6	tocilizum ab, baricitini b	Hypergly cemia, Staphylo coccus bacteriae mia,	surviv e



			disease														
10	27	F	Nil	194.00	79.07	1.33	1707.7	6	1G	7	baricitinib	Hyperglycemia, GI bleeding	survive				
11	65	F	diabetes	165.00	250.43	0.45	885.9	2	1G	3	nil	Hyperglycemia, Septicemic shock	survive				
12	46	F	hypertension	303.13	44.3	1.36	2537.7	7	1G	3	baricitinib	Hyperglycemia	survive				
13	67	M	Coronary artery disease, benign prostate hypertrophy, renal stones disease	190.00	121.87	0.71	487.4	3	1G	3	baricitinib	Hyperglycemia, shock	survive				




Figure 1. Box plots showing serial changes in disease severity parameters after PDMPT. Day 1 represents the first day of PDMPT. The horizontal line within each box represents the median and the cross represents the mean, the lower and upper border of each box represent the lower and upper quartiles and the T bar represents 1.5 x IQR and the dots are outliers. For measurement of hypoxia, SpO<sub>2</sub>/FiO<sub>2</sub> was used as an estimate of PaO<sub>2</sub>/FiO<sub>2</sub>, the value of 200-300 represents mild degree, 100-200 represents moderate degree and <100 represents severe degree of ARDS.

## Discussion

In this study, we reported the outcome of 13 very severe covid 19 cases, treated with pulse dose methylprednisolone therapy. These patients were treated in a time of pandemic crisis in a very resource-limited setting where advanced ICU care was practically unavailable. Pulse dose methylprednisolone therapy was used as a last resort salvage therapy and we could save 76% of them.

Current treatment guidelines on Covid-19 recommend 6mg of dexamethasone in severe cases requiring oxygen therapy [3]. This recommendation was based on results of clinical trials randomizing all severe patients and lack of statistically significant results in comparison with slightly larger dose of dexamethasone. However, we experienced certain subgroup of patients whose covid-19 related alveolar damage and markers of inflammatory responses were not adequately controlled and who were at imminent risk of progressing into critical stage. In those patients, we tried pulse dose methylprednisolone therapy with the hope that dose-dependent non-specific immunosuppressive and anti-inflammatory effect of corticosteroids might rescue those hopeless patients. There were also reports of successful uses of similarly high doses of corticosteroids. These studies were small retrospective

observational studies. There were also mixed results with different studies using different doses and different criteria for starting high dose corticosteroids.

In those studies, benefits of high dose corticosteroids were seen in terms of mortality [7,13]), oxygen requirement [16], ICU admission [13] and inflammatory markers [17]. These benefits were found especially in age less than 70 years of age (7) and if given not too early or too late [16]. Tromp et al (2021) also reported that high dose corticosteroid therapy with methylprednisolone 1000mg/day for 3 days did not increase viral replication [17]. However, recent systematic review and meta-analysis of 12 studies involving 2759 patients reported that high dose corticosteroid was not better than conventional dose corticosteroid [15]. The trials in this meta-analysis used various doses, duration and regimen of high dose corticosteroid, resulting in high heterogeneity. A retrospective cohort study by Yaqoob et al (2022) also found that pulse dose steroid (MP 1G/day) was not better than conventional dose, with higher rate of acute kidney injury [18]. However, this study was done in patients admitted already to ICU, which is different from our patient population who are mostly inaccessible to ICU care. Another cohort study of 573 patients also reported higher mortality with corticosteroid doses higher than 250mg/day [12]. In this study, remdesivir use was one of exclusion criteria and it may be one reason why more intensive immunosuppression resulted in higher mortality due to unchecked viral replication.

In our cohort of patients, we used higher doses of corticosteroids (1 gram of methylprednisolone for 3 to 7 days) and we started this regime as soon as the standard treatment was not apparently working. The criteria for initiation were high oxygen requirement more than 10 litre/minute, more than 50% of parenchymal involvement in chest x rays, and high inflammatory markers in the absence of confirmed or suspected co-infection. Rapid progression of disease severity in terms of oxygen requirement and serial chest x rays was also an important factor in considering the more aggressive treatment. All of our patients were also treated with intravenous remdesivir for 5 days. Intravenous tocilizumab was also given if the patient afforded the cost of drug. However, this therapy was hard to start immediately since it was not readily available at the time. Nine out of 13 patients received tocilizumab therapy while the other 4 did not. The outcome was not significantly associated with the addition of additional immunomodulator therapy like tocilizumab. All 4 patients who could not afford this therapy survive but some treated with tocilizumab were lost.

After initiation of therapy, CRP levels invariably fall after a day or two, but rise again in the 3rd week in some cases (Figure 1.). Superimposed infection was identified in some of those cases and successfully treated. However, improvement in oxygen level was noticeable only after a week. Ferritin level and Chest x ray scores showed no noticeable changes after PDMPT until 3 weeks of follow-up. This can be explained by the facts that serum ferritin was not frequently measured in our cohorts and follow up period was short to detect the usually slow radiological improvements. In two cases of mortality, there was no improvement in hypoxia, CRP level rise again after initial falling. This indicates that inflammatory response was not adequately controlled or smoldering co-infection might have been missed. These cases had very high ferritin and CRP before starting PDMPT. In the third case of mortality, oxygenation and inflammatory markers improved but patient was lost due to a CNS complication, a thrombotic event or infection, which we could not confirm. High ferritin was only significantly associated with mortality in our cohort.

Generally, the cases we treated with PDMPT were very severe from the outset and failed to respond to standard treatments. Remdesivir and PDMPT (in combination with tocilizumab in some cases) successfully rescued most of them, even in a setting without advanced ICU care. Hyperglycemia was very common after PDMPT as expected and empirical antibiotics therapies were used in all cases but definitive infective infection was proved only in three cases.

We recommend further randomized controlled trials of pulse dose methylprednisolone therapy in very severe covid-19 patients with high oxygen requirement,

diffuse parenchymal involvement in chest x ray or rapidly changing x rays and high markers of inflammation. Since contextual factors like availability of high ends supportive care are varied in different health care settings, defining health care resources settings in future trials might be useful. We observed several limitations of this study. First, we reported the story of a small number of cases and these cases were not randomly selected, limiting the generalizability of the study results. The cases were followed up for only 28 days and we did not study the long-term complications.

#### 4. CONCLUSION

Intravenous Pulse dose methylprednisolone 1 gram per day for 3 to 7 days might be an effective salvage therapy for very severe covid 19 patients. This is especially hopeful for patients in resource-limited setting. Future studies are warranted in carefully selected population of very severe disease.

#### CONSENT

Written informed consent was obtained from the patient or the next of kin for publication of this case series. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

#### ETHICAL APPROVAL

All authors hereby declare that this study was approved by the local ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

#### REFERENCES

1. WHO Coronavirus (COVID-19) Dashboard | WHO Coronavirus (COVID-19) Dashboard With Vaccination Data [Internet]. [cited 2023 May 25]. Available from: <https://covid19.who.int/>
2. Lamers MM, Haagmans BL. SARS-CoV-2 pathogenesis. Nature Reviews Microbiology 2022 20:5 [Internet]. 2022 Mar 30 [cited 2023 May 25];20(5):270–84. Available from: <https://www.nature.com/articles/s41579-022-00713-0>
3. Corticosteroids | COVID-19 Treatment Guidelines [Internet]. [cited 2023 May 28]. Available from: <https://www.covid19treatmentguidelines.nih.gov/therapies/immunomodulators/corticosteroids/>
4. Anesi GL, Jablonski J, Harhay MO, Atkins JH, Bajaj J, Baston C, et al. Characteristics, Outcomes, and Trends of Patients With COVID-19-Related Critical Illness at a Learning Health System in the United States. Ann Intern Med [Internet]. 2021 May 1 [cited 2023 May 28];174(5):613–21. Available from: <https://pubmed.ncbi.nlm.nih.gov/33460330/>
5. Biccard BM, Gopalan PD, Miller M, Michell WL, Thomson D, Ademuyiwa A, et al. Patient care and clinical outcomes for patients with COVID-19 infection admitted to African high-care or intensive care units (ACCCOS): a multicentre, prospective, observational cohort study. Lancet [Internet]. 2021 May 22 [cited 2023 May 28];397(10288):1885–94. Available from: <https://pubmed.ncbi.nlm.nih.gov/34022988/>
6. Suran K, Pushpa S. Pulse therapy – A newer approach. Indian Journal of Multidisciplinary Dentistry. 2017;7(1):41.

7. Climente-Martí M, Ruiz-Millo O, López-Cruz I, Atienza-García Á, Martínez-Moragón E, Garijo-Gómez E, et al. Impact of intermediate to high doses of methylprednisolone on mortality rate in patients with COVID-19 pneumonia-induced severe systemic inflammation. *Int J Clin Pract*. 2021 Sep 1;75(9).
8. Fan Q, Deng K, Huang H, He R, Deng X, Lan Y, et al. Kinetics of Severity Biomarkers and Immunological Features of Methylprednisolone Therapy for Severe COVID-19 Patients. *Front Immunol*. 2022 Mar 8;13.
9. Go RC, Nyirenda T, Bojarian M, Hosseini DK, Kim K, Rahim M, et al. Racial/ethnic disparities on inflammation and response to methylprednisolone in severe COVID-19 pneumonia. *BMC Infect Dis*. 2022 Dec 1;22(1).
10. Kumar V, Kashyap A, Kaur S, John M, Sibia R, Chopra V, et al. Comparison of Tocilizumab and High-dose Methylprednisolone Pulse on Outcomes in Severe Corona Virus Disease-2019: TAME-COVID, a Retrospective Multicentric Study. *Int J Appl Basic Med Res [Internet]*. 2021 [cited 2022 Mar 14];11(4):263. Available from: <https://pubmed.ncbi.nlm.nih.gov/34912692/>
11. Lim PC, Wong KL, Rajah R, Chong MF, Chow TS, Subramaniam S, et al. Comparing the efficacy of tocilizumab with corticosteroid therapy in treating COVID-19 patients: a systematic review and meta-analysis. Vol. 30, DARU, *Journal of Pharmaceutical Sciences*. Springer Science and Business Media Deutschland GmbH; 2022. p. 211–28.
12. Monreal E, Sainz de la Maza S, Natera-Villalba E, Beltrán-Corbellini Á, Rodríguez-Jorge F, Fernández-Velasco JI, et al. High versus standard doses of corticosteroids in severe COVID-19: a retrospective cohort study. *European Journal of Clinical Microbiology and Infectious Diseases*. 2021 Apr 1;40(4):761–9.
13. Pinzón MA, Ortiz S, Holguín H, Betancur JF, Arango DC, Laniado H, et al. Dexamethasone vs methylprednisolone high dose for Covid-19 pneumonia. *PLoS One [Internet]*. 2021 May 1 [cited 2022 Dec 27];16(5). Available from: <https://pubmed.ncbi.nlm.nih.gov/34033648/>
14. Ro S, Nishimura N, Imai R, Tomishima Y, So C, Murakami M, et al. Identification of patients with COVID-19 who are optimal for methylprednisolone pulse therapy.
15. Tan RSJ, Ng KT, Xin CE, Atan R, Yunus NM, Hasan MS. High-Dose versus Low-Dose Corticosteroids in COVID-19 Patients: a Systematic Review and Meta-analysis. *J Cardiothorac Vasc Anesth*. 2022 Sep 1;36(9):3576–86.
16. Tokano M, Tarumoto N, Imai K, Sakai J, Ishibashi N, Yokota K, et al. Timing of Initiation of Methylprednisolone Pulse Therapy in Patients with COVID-19. *COVID*. 2022 Apr 21;2(5):532–9.
17. Tromp K, van der Zee P, Rokx C, van Kampen J, Gommers D, Endeman H. Effect of Methylprednisolone on Inflammation and Coagulation in Patients with Severe COVID-19: A Retrospective Cohort Study. Vol. 16, *Biomarker Insights*. SAGE Publications Ltd; 2021.
18. Yaqoob H, Greenberg D, Hwang F, Lee C, Vernik D, Manglani R, et al. Comparison of pulse-dose and high-dose corticosteroids with no corticosteroid treatment for COVID-19 pneumonia in the intensive care unit. *J Med Virol*. 2022 Jan 1;94(1):349–56.
19. Zúñiga MÁL, Moreno-Moral A, Ocaña-Granados A, Padilla-Moreno FA, Castillo-Fernández AM, Guillamón-Fernández D, et al. High-dose corticosteroid pulse therapy increases the survival rate in COVID-19 patients at risk of hyper-inflammatory response. *PLoS One [Internet]*. 2021 Jan 1 [cited 2023 May 17];16(1):e0243964. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0243964>
20. Kumar A, Singh S, Sherawat V. Pulse therapy – A newer approach. *Indian Journal of Multidisciplinary Dentistry*. 2017 Oct 26;8(11):1739.

21. Borghesi A, Maroldi R. COVID-19 outbreak in Italy: experimental chest X-ray scoring system for quantifying and monitoring disease progression. *Radiologia Medica* [Internet]. 2020 May 1 [cited 2023 May 17];125(5):509–13. Available from: <https://link.springer.com/article/10.1007/s11547-020-01200-3>
22. Clinical Spectrum | COVID-19 Treatment Guidelines [Internet]. [cited 2023 May 17]. Available from: <https://www.covid19treatmentguidelines.nih.gov/overview/clinical-spectrum/>

#### **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

**ARDS:** acute respiratory distress syndrome

**Covid-19:** Coronavirus disease 2019

**CRP:** C-reactive protein level in serum

**ICU:** intensive care unit

**PDMP:** pulse dose methylprednisolone therapy

**SPO<sub>2</sub>:** peripheral capillary oxygen saturation

**SPO<sub>2</sub>/FI<sub>O2</sub>:** ratio of peripheral capillary oxygen saturation to inspired fraction of oxygen

UNDER PEER REVIEW