

## Effect of Plant growth regulators and Micro nutrient on quality of Ber (*Zizyphus mauritiana* Lamk.) cv. Gola.

### ABSTRACT

The present experiment was carried out on twenty eight year old ber orchard planted under sodic soil condition which is located at the Main Experiment Station Horticulture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya. The objective of experiment was to assess the effect of PGRs and micro nutrient on quality of ber. The application of different growth regulators affected different characters of ber significantly i.e., Fruit weight, Fruit size, TSS, acidity, ascorbic acid, reducing sugar, non-reducing sugar and Total sugar. The maximum fruit size, length (3.87 and 3.92cm) and breadth (3.38 and 3.41 cm) was recorded under the treatment of T<sub>8</sub>- 0.15 % Promalin+0.5% Borax, which was statistically at par with T<sub>10</sub>-15 ppm GA<sub>4+7</sub>+0.5 % Borax (3.79 and 3.84cm) and (3.33 and 3.36cm). The maximum fruit weight was recorded (27.69g and 28.89g) under the treatment of T<sub>8</sub>- 0.15 % Promalin + 0.5% Borax, which was statistically at par with T<sub>10</sub>-15 ppm GA<sub>4+7</sub> + 0.5 % Borax (26.30g). The maximum TSS (19.42 and 19.49) °Brix was recorded under the treatment of T<sub>8</sub>- 0.15 % Promalin + 0.5 % Borax, which was statistically at par with T<sub>10</sub>-15 ppm GA<sub>4+7</sub> + 0.5 % Borax (19.02 and 19.09) °Brix. The minimum Acidity (0.24% and 0.23 %) was recorded under the treatment of T<sub>8</sub>- 0.15 % Promalin + 0.5 % Borax, which was statistically at par with T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax. The maximum Ascorbic acid (85.33 and 86.02) was recorded under the treatment of T<sub>8</sub>- 0.15 % Promalin + 0.5 % Borax which was statistically at par with T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax (84.71 and 84.62). The maximum Reducing sugar (6.23 and 6.25) was recorded under the treatment of T<sub>7</sub>- 0.13 % Promalin + 0.5 % Borax. The maximum non-reducing sugar (7.92 and 7.95) was recorded under the treatment of T<sub>7</sub>- 0.13% Promalin +0.5 % Borax. The maximum Total sugar (14.15 and 14.20) was recorded under the treatment of T<sub>7</sub>- 0.13 % Promalin +0.5 % Borax.

**Keywords:** Acidity, Ber, Boron, GA<sub>4+7</sub>, Promalin, Sugars

Not mentioned the study years and strictly avoid the repetition, also give the proper keywords.

### INTRODUCTION

The Indian Ber (*Zizyphus mauritiana* Lamk.) belong to family Rhamnaceae and genus *Zizyphus* which includes about fifty species, and among those, 18-20 species are native to India (Pareek,1983). It is tetraploid in nature with a chromosome number  $2n=4x=48$ . Origin place of fruit ber is believed to be India to South - Western Asia. Ber is a very famous ancient fruit crop of India and China. It is also called as Chinese date or Chinese fig or plum. It is an important minor fruit of India which is reported to be growing in other countries also such as Iran, Syria, Australia, USA, Russia, Myanmar etc. Ber is cultivated in various part of country particularly in arid and semi-arid regions comprising of 53,000 ha area, producing 5.70 lakh MT of fruits (National Horticulture Board, 2021-22). The major ber growing regions are Punjab, Haryana, Uttar Pradesh, Rajasthan, Maharashtra, Andhra Pradesh, Bihar, and Madhya Pradesh. Ber tree bears its inflorescence in the axils of leaves on current season's growth. The flowering period lasts for about two and a half months from September to November. The fruit setting starts in second week of October and continues up to first fortnight of November.

Comment [AS1]: Comma after this

Comment [AS2]: First full form the short in bracket

Comment [AS3]: Small

Comment [AS4]: Small

Comment [AS5]: Small

Comment [AS6]: during 2021-22 and 2022-23, respectively

Comment [AS7]: highest

Comment [AS8]: noted

Comment [AS9]: significantly similar

Comment [AS10]: Second year figure

Comment [AS11]: Higher

Comment [AS12]: observed

Comment [AS13]: that

Comment [AS14]: Small

Comment [AS15]: Significantly same

Comment [AS16]: Small

Comment [AS17]: Comma after this

Comment [AS18]: Small

Comment [AS19]: Comma after this

Comment [AS20]: Comma after this

The ripe ber fruit have high nutritive values and conventionally it is considered a “Poor man’s fruit”. The ber fruit is richer than apple in protein, phosphorus, calcium and Vitamin ‘C’. and one hundred gram of edible ber contains moisture (85.9%), protein(0.8g), fat (0.1g), carbohydrates (12.88%), calcium (0.03g), phosphorus (0.03g), iron (0.8g), beta carotene (70 IU), vitamin C (50-100mg/100g pulp) etc. Ripe ber fruits are eaten fresh. Fruits is also dried and used as dessert. It gives an excellent product when candy. Other processed products made are ber butter, ber juice or squash, RTS etc.

Comment [AS21]: ?

Plant bio-regulators and micronutrient such as Promalin, GA<sub>4+7</sub> and borax play an important role in increase fruits size, fruits set, fruit yield, self-life and quality improvement. Promalin plays an important role for increase the size and shape of the fruit. Promalin containing gibberellins and cytokine promotes both cell division and cell enlargement.

Comment [AS22]: Avoid repetition

Comment [AS23]: Avoid repetition

GA<sub>4+7</sub> plays an important role for induce fruit elongation and to increase the size and quality, fruit set, changing fruit shape and size, increasing weight of fruit,

Comment [AS24]: Avoid repetition

Boron is for most important in the micro nutrients. Boron also plays important role for nucleic acid and lignin synthesis, pollen tube growth, it is associated with carbohydrate and fat metabolism and translocation and transformation of sugars.

Not given the findings of the previous studies related to the treatments whether need to study in ber

## MATERIALS AND METHODS

The present study was conducted on twenty eight years old uniform tree of ber cv. Gola, planted at a distance of 8 x 8m in Main Experimental Station of Horticulture, Acharya Narendra Deva University of Agriculture & Technology, Narendra Nagar, Kumarganj, Ayodhya. (26<sup>o</sup>.47<sup>o</sup>N latitude, 82.12<sup>o</sup>E longitude of 113 meters altitude away from mean sea stratum) during 2021-22 and 2022-23. Plants growth regulators and micro-nutrient, viz. promalin (0.13% and 0.15%), GA<sub>4+7</sub> (10 ppm and 15 ppm) and borax (0.5%). Experiment was laid out in randomized block design (RBD) with three replication per treatment and considering one tree per replication. Data on fruit drop was recorded periodically from October to March. Ber fruits from sprayed plants were harvested month of March and analyzed for quality attributes.

Comment [AS25]: 8m

Comment [AS26]: Capital

Comment [AS27]: Give properly treatments

Comment [AS28]: replications

Comment [AS29]: in month

Ten randomly selected fruits from each replication were used to assess quality attributes of ber. Physical parameters like fruit size in term of length and breadth (cm) were measured with help of vernier’s calliper and expressed in centimetres. Fruit weight were recorded with help of digital weighing machine and expressed in gram. Among bio-chemical parameters, TSS content was determined with hand refractometer and expressed in per cent (Ranganna, 1986). Acidity was expressed as per cent of citric acid as method given by Kaur, A. and Kalia, M.(2017). Ascorbic acid content was expressed in mg/100g of pulp and was estimated using 2,6-dichlorophenol indophenol dye by visual titration method(AOAC 1975).Total sugar reducing, non-reducing expressed in per cent was estimated by method suggested by AOAC (1975).The experiment data were analyzed following

Comment [AS30]: the

Comment [AS31]: estimated

Comment [AS32]: presented

Comment [AS33]: noted

Comment [AS34]: Kaur and Kalia

Comment [AS35]: indophenols

Comment [AS36]: Comma after this

RBD to test the significance of using the Statistix-10 software. The differences in quantified concentrations were evaluated using F test at  $P < 0.05$ . Statistical analysis was done as described by Gomez and Gomez (1984).

## RESULTS AND DISCUSSION

### Fruit Size (cm)

#### Fruit Length (cm)

A perusal of data on fruit length presented in Table (1.) indicated that all treatments have a significant effect in improving the fruit length over control .The maximum fruit length (3.87cm and 3.92 cm) was recorded with  $T_8$ -0.15% Promalin+0.5% Borax, which was significantly superior over the rest of the treatments and at par with  $T_{10}$  -15 ppm  $GA_{4+7}$  +0.5% Borax (3.79cm & 3.84cm) followed by  $T_9$ -10 ppm  $GA_{4+7}$ + 0.5% Borax (3.64cm and 3.69 cm ),  $T_5$  -15 ppm  $GA_{4+7}$  (3.54cm and 3.59cm) and  $T_7$ - 0.13% Promalin+0.5% Borax (3.39cm and 3.43cm),  $T_3$ - 0.15 % Promalin (3.28cm and 3.32cm),  $T_4$ - 10 ppm  $GA_{4+7}$  (3.22cm and 3.25cm),  $T_2$ - 0.13 % Promalin (3.13 and 3.17),  $T_6$  -0.5 % Borax(3.10&3.14cm) respectively. While minimum fruit length (3.08cm and 3.11cm) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS37]: No bracket

Comment [AS38]: had

Comment [AS39]: and

Comment [AS40]: and

Comment [AS41]: Before this comma, why is respectively

#### Fruit breadth (cm)

A perusal of data on fruit length presented in Table 1. indicated that all treatments have a significant effect in improving the fruit breadth over control .The maximum fruit breadth (3.38 cm and 3.41 cm ) was recorded with  $T_8$  - 0.15% Promalin+0.5% Borax, which was significantly superior over the rest of the treatments and at par with  $T_{10}$  15 ppm  $GA_{4+7}$ +0.5% Borax( 3.33cm and 3.36cm) and  $T_9$ -10 ppm  $GA_{4+7}$ + 0.5% Borax (3.32 cm and 3.35 cm ), followed by  $T_5$  -15 ppm  $GA_{4+7}$  (3.30 cm and 3.33cm ),  $T_7$ - 0.13% Promalin+0.5% Borax (3.26 cm and 3.29cm )  $T_3$ - 15ppm Promalin (3.22 and 3.25 cm ),  $T_4$ -10 ppm  $GA_{4+7}$  ( 3.18cm and 3.21cm) and  $T_2$ -0.13 % Promalin(3.10cm and 3.13cm),  $T_6$ - 0.5% Borax (3.00cm and 3.03cm ), respectively. While minimum fruit breadth (2.60cm and 2.63cm) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS42]: given

Comment [AS43]: Repetition it should be avoided

Comment [AS44]: highest

Comment [AS45]: noted

Comment [AS46]: statistically

Comment [AS47]: . It was followed

Comment [AS48]: In 2021-22 and 2022-23, respectively

Comment [AS49]: Repetition so should be avoided

Comment [AS50]: Comma after this

Fruit size was reported to be a genetic characteristic of the cultivars. The increase in size of fruit (length and width) over control as a result of foliar application of borax , Promalin(6BA) and  $GA_{4+7}$  in present investigation might be due to their involvement in cell division, cell elongation, increased volume of intercellular spaces in the mesocarpic cells (Brahmachari *et al.*, 1997), Promalin ,  $GA_{4+7}$  is also reported to promote growth by increasing plasticity of the cell wall followed by the hydrolysis of starch into sugar which reduces the cell water potential, resulting in the entry of water into the cell and causing elongation (Kassem *et al.*, 2011).

Increased in fruit size with plant growth regulators and micro-nutrient application has also been reported by Stembridge and Morell (1972) reported that maximum fruit size application of

Comment [AS51]: , the

Comment [AS52]: Size was obtained with the

promalin @ 50 ppm. Unarth (1974) foliar application of 25 and 50ppm promalin sprayed in apple, Cibulsky (1977); Looney (1979); Comai *et al.* (1981) in apple, Curracy and Williams (1982); Eccher and Maffi (1986) in apple, Pal (1997) in apple.

### Fruit weight (g) / fruit

A perusal of data on fruit weight shown in Table1. indicated that all treatments have a significant effect in improving the fruit weight over control. The maximum fruit weight (27.69 g and 28.89g) was recorded with T<sub>8</sub> - 0.15% Promalin+0.5%Borax, which was significantly superior over the rest of the treatments and at par with T<sub>10</sub> -15 ppm GA<sub>4+7</sub> +0.5% Borax ( 26.30g in 2021-22), followed by T<sub>9</sub> -10 ppm GA<sub>4+7</sub>+ 0.5% Borax (25.87g,) T<sub>5</sub> -15 ppm GA<sub>4+7</sub> (23.96 g and 25.16 g ),T<sub>7</sub> - 0.13% Promalin+0.5% Borax ( 23.40g and 24.60g) , T<sub>3</sub>- 15ppm Promalin (23.24g and 24.44g),T<sub>4</sub> -10 ppm GA<sub>4+7</sub> ( 22.83g and 24.03g), T<sub>2</sub>-0.13 % Promalin (21.97g and 23.17g) respectively. While minimum fruit weight (19.89g and 21.09) was recorded with control during 2021-22 and 2022-23, respectively.

**Table 1.: Effect of foliar feeding of plant growth regulators and micro-nutrient on Fruit size, and Fruit weight of ber fruit**

Treatment	Parameters					
	Length (cm)		Breadth (cm)		Fruit weight (g)/fruit	
	2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T <sub>1</sub> - Control	3.08 <sup>i</sup>	3.11 <sup>e</sup>	2.60 <sup>j</sup>	2.63 <sup>f</sup>	19.89 <sup>f</sup>	21.09 <sup>f</sup>
T <sub>2</sub> - 0.13 % Promalin	3.13 <sup>ef</sup>	3.17 <sup>de</sup>	3.10 <sup>d</sup>	3.13 <sup>de</sup>	21.97 <sup>de</sup>	23.17 <sup>de</sup>
T <sub>3</sub> - 0.15 % Promalin	3.28 <sup>cd</sup>	3.32 <sup>cd</sup>	3.22 <sup>c</sup>	3.25 <sup>bcd</sup>	23.24 <sup>cd</sup>	24.44 <sup>c</sup>
T <sub>4</sub> - 10 ppm GA <sub>4+7</sub>	3.22 <sup>de</sup>	3.25 <sup>de</sup>	3.18 <sup>c</sup>	3.21 <sup>cd</sup>	22.83 <sup>cde</sup>	24.03 <sup>cd</sup>
T <sub>5</sub> - 15 ppm GA <sub>4+7</sub>	3.54 <sup>b</sup>	3.59 <sup>b</sup>	3.30 <sup>b</sup>	3.33 <sup>abc</sup>	23.96 <sup>c</sup>	25.16 <sup>c</sup>
T <sub>6</sub> - 0.5 % Borax	3.10 <sup>f</sup>	3.14 <sup>e</sup>	3.00 <sup>e</sup>	3.03 <sup>e</sup>	21.36 <sup>ef</sup>	22.56 <sup>e</sup>
T <sub>7</sub> - 0.13 % Promalin + 0.5 % Borax	3.39 <sup>c</sup>	3.43 <sup>c</sup>	3.26 <sup>bc</sup>	3.29 <sup>abc</sup>	23.40 <sup>cd</sup>	24.60 <sup>c</sup>
T <sub>8</sub> - 0.15 % Promalin + 0.5 % Borax	3.87 <sup>a</sup>	3.92 <sup>a</sup>	3.38 <sup>a</sup>	3.41 <sup>a</sup>	27.69 <sup>a</sup>	28.89 <sup>a</sup>
T <sub>9</sub> - 10 ppm GA <sub>4+7</sub> + 0.5 % Borax	3.64 <sup>b</sup>	3.69 <sup>b</sup>	3.32 <sup>ab</sup>	3.35 <sup>abc</sup>	25.87 <sup>b</sup>	27.07 <sup>b</sup>
T <sub>10</sub> -15 ppm GA <sub>4+7</sub> + 0.5 % Borax	3.79 <sup>a</sup>	3.84 <sup>a</sup>	3.33 <sup>ab</sup>	3.36 <sup>ab</sup>	26.30 <sup>ab</sup>	27.50 <sup>b</sup>
<b>LSD (p≤0.05)</b>	<b>0.113</b>	<b>0.155</b>	<b>0.079</b>	<b>0.135</b>	<b>1.699</b>	<b>1.199</b>

Increase in fruit weight with the spray of PGRs and micro-nutrient might be due to faster loading and mobilization into fruits (Brahmachari *et al.* 1997) and involvement in hormonal metabolism, increased cell division and expansion of cell. This may also be attributed to greater photosynthetic activity, resulting the increased production and accumulation of carbohydrates and favourable effect on vegetative growth and retention of fruits, which might have increased size and weight.

The present findings regarding the increase in fruit size and fruit weight with application of promalin and GA<sub>4+7</sub> may be due to cell division and cell elongation. These results in close conformity with results of Looney *et al.* (1992), McArtney (2010), Chen *et al.* (2012), Ito *et al.* (2015), Sekhar (2012) and Al-madhagi *et al.* (2012) in strawberry. Kumar *et al.* (2003) and Yildirim *et al.* (2014) also

**Comment [AS53]:** Not clear simply given foliar application in apple so no results given, all the references very old

**Comment [AS54]:** The

**Comment [AS55]:** Just repetition so change it accordingly

**Comment [AS56]:** that

**Comment [AS57]:** Repetition

**Comment [AS58]:** Repetition

**Comment [AS59]:** Small

**Comment [AS60]:** Delete

**Comment [AS61]:** Comma after this

found significant increase in fruit weight and size with application of promalin and GA<sub>4+7</sub> in fruit crops.

### TSS (Brix<sup>0</sup>)

A perusal of data depicted in Chart 1 that all treatments have a significant in improving the TSS. The maximum TSS (19.42 & 19.49) was recorded under the treatment of T<sub>8</sub>- 0.15 % Promalin+0.5 % Borax, which was statistically *at par* with T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax (19.02 and 19.09) and T<sub>9</sub>-10ppm GA<sub>4+7</sub> + 0.5 % Borax (18.76 and 18.83), followed by T<sub>7</sub>- 0.13 % Promalin+0.5% Borax (18.41 and 18.48), T<sub>6</sub>- Borax 0.5% (18.03 and 18.10), T<sub>5</sub>- 15ppm GA<sub>4+7</sub> (17.58 and 17.65), T<sub>3</sub>- 0.15 % Promalin (17.20 and 17.27), T<sub>4</sub>- 10 ppm GA<sub>4+7</sub> (16.56 and 16.63) and T<sub>2</sub>- 0.13 % Promalin (15.83 and 15.90), while minimum TSS (14.80 and 14.87) was recorded with control during 2021-22 and 2022-23, respectively.

Increased TSS due to metabolic modification of starch and pectin, hydrolysis of complex polysaccharides, metabolite synthesis, and fast sugar translocation from leaves to developing fruits, foliar application of PGRs and micronutrients enhances total soluble solids (TSS) content in fruits. Total soluble solids (TSS) are also increased by promalin and boron, with a maximum TSS of 0.5% Borax and 0.15% Promalin applied topically. These results are also in close conformity with those of Jindal *et al.* (2004) in apple, Pandey and Kumar (2022) and Pal *et al.* (2022) in ber cv. Gola, Montalti *et al.* (1984) in apple, Canli *et al.* (2009) in pear, Wismer *et al.* (1995) in apples, Cheolku *et al.* (2000) in pear, and Sharma (2001) in apple cv. Starking Delicious.

Comment [AS62]: Fig.

Comment [AS63]: and

Comment [AS64]: Repetition

Comment [AS65]: Repetition

Comment [AS66]: Similar with the findings




Chart 1: Effect of foliar feeding of plant growth regulators and micro-nutrient on TSS, Acidity and Ascorbic acid of ber fruit.

Comment [AS67]: Fig.

Comment [AS68]: Small

Comment [AS69]: Small

### Acidity (%)

A perusal of data Chart 1 revealed that all treatments have a significant in improving the Acidity per cent. The minimum Acidity per cent (0.24 and 0.23) was recorded under the treatment of T<sub>8</sub>-0.15% Promalin+0.5% Borax, which was statistically *at par* with T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> +0.5 % Borax (0.25 and 0.24) , followed by T<sub>9</sub>-10 ppm GA<sub>4+7</sub> +0.5% Borax (0.26 and 0.25), T<sub>7</sub>-0.13 % Promalin+ 0.5% Borax (0.27 and 0.26), T<sub>6</sub>- Borax 0.5% (0.28 and 0.27),T<sub>5</sub>- 15 ppm GA<sub>4+7</sub>(0.29 and 0.28), T<sub>3</sub>- 0.15 % Promalin (0.29 and 0.28), T<sub>4</sub>- 10 ppm GA<sub>4+7</sub>(0.30 and 0.29), T<sub>2</sub>- 0.13% Promalin, while maximum Acidity (0.32 and 0.31) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS70]: Fig.

Comment [AS71]: Repetition

Comment [AS72]: Small

Comment [AS73]: small

Comment [AS74]: Small

Comment [AS75]: Same pattern so change it

The decrease in the acidity might be due to higher accumulation of sugars in fruits, better translocation of sugars into fruit tissues and conversion of organic acids into sugars by foliar application of PGRs and micro-nutrient. The reduction in the acidity under boric acid treatment might be owing to increased TSS of the fruits. These results also elucidate the finding of Jindal *et al.* (2004) in apple cv. Starking Delicious, Canli *et al.* (2009) in “Deveci” pear, Pandey and Kumar (2022), Pal *et al.* (2022) in ber cv. Gola, Khalid *et al.* (2012) in Kinnow.

### Ascorbic acid (mg)

A perusal of data Chart 1 revealed that all treatments have a significant in improving the Acidity per cent .The maximum Ascorbic acid (85.33 and 86.02) was recorded under the treatment of T<sub>8</sub>-0.15 % Promalin+0.5% Borax which was statistically *at par* with T<sub>10</sub>-15 ppm GA<sub>4+7</sub> + 0.5 % Borax(84.71 and 84.62)followed by T<sub>9</sub>-10 ppm GA<sub>4+7</sub> + 0.5 % Borax(83.69 and 84.14), T<sub>7</sub>-0.13% Promalin + 0.5 % Borax (83.43 and 84.12), T<sub>6</sub> -Borax 0.5% (82.52 and 83.87), T<sub>5</sub>- 15ppm GA<sub>4+7</sub>(80.37 and 81.39), T<sub>3</sub>-0.15% Promalin (79.83 and 80.85), T<sub>4</sub>- 10 ppm GA<sub>4+7</sub> (78.20 and 79.55) and T<sub>2</sub>-0.13 % Promalin (77.74 and 78.43), while minimum ascorbic acid (76.82 and 77.84) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS76]: Fig.

Comment [AS77]: Small

Comment [AS78]: Change the presentation pattern

The increase in ascorbic acid content might be speculated due to increased activity of enzyme responsible for the synthesis of the ascorbic acid precursor i.e., glucose-6-phosphate or inhibition of its conversion into dehydro ascorbic acid by enzyme ascorbic acid oxidase or both and also the reduction in the rate of respiration by these chemicals (Brahmachari and Rani 2001). Similar result have been reported by Pal *et al.*,(2022) and Pandey and Kumar (2022) in ber cv. Gola, Khalid *et al.*, (2012) in Kinnow and Cheolku *et al.* (2000) in pear.

Comment [AS79]: Comma after this

### Reducing sugar (%)

A perusal of data Chart 2 revealed that all treatments have a significant in improving the Acidity percent. The maximum reducing sugar (6.23 and 6.25) was recorded under the treatment of T<sub>7</sub> -0.13 % Promalin + 0.5 % Borax followed by T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax (5.96 and 5.98), T<sub>9</sub>-10 ppm GA<sub>4+7</sub> +0.5 % Borax(5.84 and 5.87), T<sub>8</sub>- 0.15 % Promalin+0.5 % Borax (5.73 and 5.75), T<sub>6</sub> - Borax 0.5% (5.59 and 5.62), T<sub>5</sub>- 15 ppm GA<sub>4+7</sub>(5.43 and 5.45),T<sub>3</sub>- 0.15 % Promalin(5.30 and 5.32),

Comment [AS80]: Fig.

T<sub>4</sub>- 10 ppm GA<sub>4+7</sub>(5.07 and 5.14) and T<sub>2</sub>-0.13 % Promalin (4.82 and 4.84), while minimum reducing sugar (4.47 and 4.49) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS81]: Change it

Foliar spray of PGRs and micro-nutrients is helpful to increase the reducing sugar level, which could be due to accumulation of more carbohydrates to the fruit which results the better accessibility of nutrition for developing fruits and increases the reducing sugar level of fruits in present investigation. The result in conformity with those of Pandey and Kumar (2022) and Pal *et al.*(2022) in ber cv. Gola, Jindal *et al.* (2004) in apple, Canli *et al.* (2009) in pear and Khalid *et al.*, (2012) in Kinnow and Cheolku *et al.* (2000) in pear.

### Non-reducing sugar (%)

A perusal of data Chart 2 revealed that all treatments have a significant in improving the Acidity per cent. The maximum Non-reducing sugar (7.92 and 7.95) was recorded under the treatment of T<sub>7</sub>- 0.13 % Promalin +0.5 % Borax followed by T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax (7.87 and 7.90), T<sub>9</sub>- 10 ppm GA<sub>4+7</sub> + 0.5 % Borax (7.72 and 7.74), T<sub>8</sub>- 0.15 % Promalin + 0.5 % Borax (7.56 and 7.59), T<sub>6</sub> - Borax 0.5% (7.39 and 7.41), T<sub>5</sub>- 15ppm GA<sub>4+7</sub> (7.17 and 7.20), T<sub>3</sub>- 0.15 % Promalin (6.99 and 7.02), T<sub>4</sub>-10 ppm GA<sub>4+7</sub> (6.69 and 6.72) and T<sub>2</sub>- 0.13 % Promalin (6.37 and 6.40), while minimum non-reducing sugar (5.89 and 5.92) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS82]: Fig.

Comment [AS83]: Change it




Chart 2: Effect of foliar feeding of plant growth regulators and micro-nutrient on Reducing sugar, Non-reducing sugar and Total sugar of ber fruit.

Comment [AS84]: Fig.

Comment [AS85]: Small

Comment [AS86]: Small

Comment [AS87]: small

Comment [AS88]: fruits was

The possible reason for increase in non-reducing sugar content of fruits due to hydrolysis of polysaccharides to simpler form i.e. mono and disaccharides and better transportation of nutrients to plant from leaves to their place of utilization. These results corroborate the earlier records of Pandey and Kumar (2022) and Pal *et al.*(2022) in ber cv. Gola, Khalid *et al.*, (2012) in Kinnow, Jindal *et al.* (2004) in apple and Cheolku *et al.* (2000) in pear.

## Total sugar (%)

A perusal of data Chart 2 revealed that all treatments have a significant in improving the Acidity per cent. The maximum Total sugar (14.15 and 14.20) was recorded under the treatment of T<sub>7</sub>- 0.13 % Promalin+0.5 % Borax followed by T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax (13.83 and 13.88), T<sub>9</sub>-10 ppm GA<sub>4+7</sub> + 0.5 % Borax (13.56 and 13.61), T<sub>8</sub>-0.15 % Promalin + 0.5 % Borax (13.29 and 13.34), T<sub>6</sub> -Borax 0.5% (12.98 and 13.03), T<sub>5</sub>- 15 ppm GA<sub>4+7</sub> (12.60 and 12.65), T<sub>3</sub>- 0.15 % Promalin (12.29 and 12.34), T<sub>4</sub>- 10 ppm GA<sub>4+7</sub> (11.76 and 11.81) and T<sub>2</sub>- 0.13% Promalin (11.19 and 11.24), while minimum Total sugar (10.36 and 10.41) was recorded with control during 2021-22 and 2022-23, respectively.

Comment [AS89]: Fig.

The possible reason for increase in total sugar content may be due to hydrolysis of starch yielding mono and disaccharide, which owned a simplest form of sugar and boron, promalin and GA<sub>4+7</sub> probably augmented the conversion of starch. Gauch and Dugger (1953) reported that the improvement in total sugar content may be due to more translocation of sugars from leaves to developing fruits. Our results are also in line with the finding of Pandey and Kumar (2022) and Pal *et al.*, (2022) in ber cv. Gola, Chaudhary *et al.* (2018) in aonla, Cheolku *et al.* (2000) in pear, Wismer *et al.* (1995) in apples and Sharma (2001) in apple cv. Starking Delicious.

Comment [AS90]: Change pattern

## CONCLUSION

It may be concluded that the application of different growth regulators resulted that improve the characters of ber *i.e.* Fruit weight, fruit size, TSS, acidity, ascorbic acid, reducing sugar, non-reducing sugar and total sugar. The sugar and ascorbic acid, parameters were significantly affected by the application of 0.15% Promalin+0.5% Borax (T<sub>8</sub>) and T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax. Based on cost benefit ratio it may be concluded that T<sub>10</sub>- 15 ppm GA<sub>4+7</sub> + 0.5 % Borax could be recommended to farmers for commercially increasing yield and quality of ber. Normally people don't prefer Gola variety ber over Apple ber due to it's small size but with the application of 15 ppm GA<sub>4+7</sub> + 0.5 % Borax we can increase fruit size in a very economical way resultant consumer's preference will also increase.

Comment [AS91]: delete

Comment [AS92]: Small

## REFERENCES:

1. A.O.A.C., 1975. Official method of analysis. Assoc. of Official Analytical Chemists, Benjamin Franklin Station, Washington D.C., 12th edition, 575.
2. Al-madhagi, IAH.; Hasan, SMZ.; Ahmad, A.; Zain, AM. and Yusoff, A. 2012. Influence of exogenous hormone on the flowering of strawberry (*Fragaria x ananassa* Duch). *Journal of Biology Agriculture and Healthcare*. 2(4):46-52.
3. Anonymous, 2021-22. Horticulture Data Base, National Horticulture Board, Gurgaon, Ministry of Agriculture, India.
4. Brahmachari, V.S and Rani, R. 2001. Effect of growth substance on productivity, cracking, ripening and quality of fruits in litchi. *Orissa journal of Horticulture*, 29(1):41-45.

Comment [AS93]: Delete

Comment [AS94]: Al-madhagi, I.A.H.; Hasan, S.M.Z.; Ahmad, A.; Zain, A.M.

5. Canli, A.F.; Pektas, M.; and Kelen, M. 2009 . Effects of Pre-harvest Plant Growth Regulator Sprays on Fruit Quality of ‘Deveci’ Pear (*Pyrus communis* L.). *Journal of Applied Biological Sciences*. 3 (1): 81-84.
6. Chaudhary, M.; Singh, D. P.; Singh, J. P. and Tomar, S. 2018. Effect of foliar application of PGRs and mineral nutrients on physico-chemical characteristics of Aonla fruit cv. NA-7. *Journal of Pharmacognosy and Phytochemistry* 7(4), 2719-2723.
7. Chen, X.; Bao, J. Chen, Y.; Chen, T.; Zhang, C.; and Huang, X. 2012. Effect of hormone treatments on deformed fruit development in pear. *African journal of biotechnology*, 11(44), 10207-10209.
8. Cheolku, Y.; Seonkyukim, S.L. and Kwansoon, C. 2000. Effect of application time of promalin on tree growth and fruit quality of pears. *Journal of Korean Society for Horticultural Science* 41(3): 289-291.
9. Cibulsky, R.J. 1977. Effect of bloom application of promalin on apple fruit. *Proceedings of the Plant Growth Regulator Society of America* 4: 223-230.
10. Comai, M.; Widemann, L. and Gianotti, G. 1981. The effect of gibberellins and promalin on fruit shape and quality of apple. *Esperienze Ricerche Stazione Sperimentale Agraria Forestale Adige*. 11: 47-61.
11. Curracy, E A. and Williams, M W. 1982. Promalin or GA<sub>3</sub> increase pedicel and fruit length and leaf size of 'Delicious' apple treated with paclobutrazol. *Horticulture Science*. 18(2): 214-215.
12. Eccher, T. and Maffi, A. 1986. Treatments for prevention of “Golden Delicious” russetting. *Acta Horticulturae*. 179:767-70.
13. Gauch, H.G. and Dugger, W.W.J 1953. The physiological action of boron in higher plants review and interpretation. Technical Bulletin, University Maryland Agriculture Experiment Station, USA, 80p.
14. Gomez, K.A. and Gomez, A.A. 1984. *Statistical Procedures of Agricultural Research (2nd Edn.)*, John Willey and Sons Inc., New York.
15. Ito, A.; Sakamoto, D.; Itai, A.; Nishijima, T.; Oyama-Okubo, N.; Nakamura, Y.; Moriguchi, T. and Nakajima, I. 2015. Effects of GA<sub>3+4</sub> and GA<sub>4+7</sub> application either alone or combined with prohexadione-Ca on fruit development of Japanese pear ‘Kosui’. *The Japanese Society for Horticulture Science*, 85(3), 201-208.
16. Jindal, K.K.; Pal, S.; Chauhan, P.S. and Mankotia, M.S. 2004. Effect of promalin and mixtatal on fruit growth, yield efficiency and quality of “Starking Delicious” apple. *Acta Horticulturae*. 636: 533-536.
17. Khalid S.; Malik, A.U.; Khan, A.S. and Jamil, A. 2012. Influence of exogenous applications of plant growth regulators on fruit quality of Young 'Kinnow' Mandarin (*Citrus nobilis* × *C. deliciosa*) Trees. *Int. J. Agric. Biol.*; 14(2):229-234.
18. Kumar, J.; Rana, S.S.; Verma, H.S. and Jindal, K.K. 2003. Effect of promalin on fruit shape, set, yield and quality of apple cv. Starking Delicious. *Indian Journal of Horticulture*. 60(2): 143-146.
19. Looney, N E. 1979. Some effects of GA<sub>4+7</sub> and BA on fruit weight, shape, quality and calcium content of 'Spartan' apple. *Journal of the American Society for Horticultural Science*. 104(3): 389-391.
20. McCartney, S. J. 2010. Exogenous gibberellin affects biennial bearing and the fruit shape of ‘Braeburn’ apple. *New Zealand Journal of Crop and Horticultural Science*, Vol. 22: 343-346
21. Montalti, P.; Cristofferi, B. and Sansavini, S. 1984. Effect of GA<sub>4+7</sub> and shoot pinching on hormonal levels in conference pear. *Acta Horticulturae*. 161: 163-170.

Comment [AS95]: Use comma

Comment [AS96]: Delete

Comment [AS97]: Use comma at all places

Comment [AS98]: Use comma at all places

22. Pal, R.; Kumar, A.; and Kumar, A. 2021. Effect of foliar feeding on chemical attributes of Ber (*Zizyphus mauritiana* L.) under Sodic soil. *International Journal of Chemical Studies*. 9 (1), 1558-1560.
23. Pal, S.J. 1997. Effect of plant bio regulators on fruit growth, fruit yield and quality of Stalking Delicious apples. M.Sc. Thesis Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, (H.P.).
24. Pandey, A.; and Kumar, A. 2022. Effect of ZnSO<sub>4</sub>, Borax and GA<sub>3</sub> on quality and yield of ber (*Zizyphus mauritiana* Lamk.) Fruits Cv. Gola. *The Pharma Innovation Journal*. 11(8): 1788-1790.
25. Pareek, B.S. 1983. The Ber- 1<sup>st</sup> edition *Indian Council of Agricultural Research*, New Delhi. pp 31-36.
26. Ranganna, S. (1986). Manual of Analysis of fruits and vegetable products, Tata Mc Graw-Hill Publishing Company Ltd; New Delhi, pp.1-30
27. Sharma, G. 2001. Effect of pre bloom foliar applications of nutrients and plant bio regulators on fruit set and yield in apple. M.Sc. Thesis Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, (H.P.).
28. Stembridge, G.E. and Morrell, G. 1972. Effect of gibberellins and 6-benzyl-adenine on the shape and fruit set of "Delicious" apples. *Journal of the American Society for Horticultural Science*. 97:464-67.
29. Unrath, C.R. 1974. The commercial implications of gibberellin A<sub>4</sub> plus benzyladenine for Improving shape and yield of 'Delicious' apples. *Journal of the American Society for Horticultural Science*. 99(4): 381-384.
30. Wismer, P T.; Proctor, J T A. and Elfving, D.C. 1995. Benzyladenine affects cell division and cell size during apple fruit thinning. *Journal of the American Society for Horticultural Science*. 120: 802-807.
31. Yildirim, F.A.; Kepenek, G.; San, B.; Yildirim, A.N. and Kacal, E. 2014. Effects of Ba + GA<sub>4+7</sub> treatments on fruit quality in "Fuji" apple variety. *Turkish Journal of Agricultural and Natural Sciences*. 2: 1387-1390.

Comment [AS99]: G.E

Comment [AS100]: C.R.

Comment [AS101]: Wismer, P. T.; Proctor, J. T. A., and Elfving, D.C.

Comment [AS102]: F.A.,

Comment [AS103]: A.N.

Need to give the latest more references to support the results