

Effect of Plant growth regulators and Micro nutrient on quality of Ber (*Zizyphus mauritiana* Lamk.) cv. Gola.

Formatted: Font: 14 pt

ABSTRACT

The present experiment was carried out on twenty eight year old ber orchard planted under sodic soil condition which is located at the Main Experiment Station Horticulture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya. The objective of experiment was to assess the effect of PGRs and micro nutrient on quality of ber. The application of different growth regulators affected different characters of ber significantly *i.e.*, **Fruit-fruit** weight, **Fruit-fruit** size, TSS, acidity, ascorbic acid, reducing sugar, non-reducing sugar and Total sugar. The maximum fruit size, length (3.87 and 3.92cm) and breadth (3.38 and 3.41 cm) was recorded under the treatment of **T₈**- 0.15 % Promalin + 0.5% Borax, which was statistically *at par* with **T₁₀**-15 ppm GA₄₊₇+0.5 % Borax (3.79 and 3.84cm) and (3.33 and 3.36 cm). The maximum fruit weight was recorded (27.69g and 28.89 g) under the treatment of **T₈**- 0.15 % Promalin + 0.5% Borax, which was statistically *at par* with **T₁₀**-15 ppm GA₄₊₇ + 0.5 % Borax (26.30g). The maximum TSS (19.42 and 19.49)⁰Brix was recorded under the treatment of **T₈**- 0.15 % Promalin + 0.5 % Borax, which was statistically *at par* with **T₁₀**-15 ppm GA₄₊₇ + 0.5 % Borax (19.02 and 19.09)⁰Brix. The minimum Acidity (0.24% and 0.23 %) was recorded under the treatment of **T₈**- 0.15 % Promalin + 0.5 % Borax, which was statistically *at par* with **T₁₀**- 15 ppm GA₄₊₇ + 0.5 % Borax. The maximum Ascorbic acid (85.33 and 86.02) was recorded under the treatment of **T₈**- 0.15 % Promalin + 0.5 % Borax which was statistically *at par* with **T₁₀**- 15 ppm GA₄₊₇ + 0.5 % Borax (84.71 and 84.62). The maximum Reducing sugar (6.23 and 6.25) was recorded under the treatment of **T₇**- 0.13 % Promalin + 0.5 % Borax. The maximum non-reducing sugar (7.92 and 7.95) was recorded under the treatment of **T₇**- 0.13% Promalin +0.5 % Borax. The maximum Total sugar (14.15 and 14.20) was recorded under the treatment of **T₇**- 0.13 % Promalin +0.5 % Borax.

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Keywords: Acidity, Ber, Boron, GA₄₊₇, Promalin, Sugars.

INTRODUCTION

The Indian Ber (*Zizyphus mauritiana* Lamk.) belong to family Rhamnaceae and genus *Zizyphus* which includes about fifty species, and among those, 18-20 species are native to India (Pareek,1983). It is tetraploid in nature with a chromosome number $2n=4x=48$. Origin place of fruit ber is believed to be India to South - Western Asia. Ber is a very famous ancient fruit crop of India and China. It is also called as Chinese date or Chinese fig or plum. It is an important minor fruit of India which is reported to be growing in other countries also such as Iran, Syria, Australia, USA, Russia, Myanmar *etc.* Ber is cultivated in various part of country particularly in arid and semi-arid regions comprising of 53,000 ha area, producing 5.70 lakh MT of fruits (National Horticulture Board, 2021-22). The major ber growing regions are Punjab, Haryana, Uttar Pradesh, Rajasthan, Maharashtra, Andhra Pradesh, Bihar, and Madhya Pradesh. Ber tree bears its inflorescence in the axils of leaves on current season's growth. The

Formatted: Font: Italic

flowering period lasts for about two and a half months from September to November. The fruit setting starts in second week of October and continues up to first fortnight of November.

The ripe ber fruit have high nutritive values and conventionally it is considered a “Poor man’s fruit”. The ber fruit is richer than apple in protein, phosphorus, calcium and Vitamin ‘C’. and one hundred gram of edible ber contains moisture (85.9%), protein (0.8g), fat (0.1g), carbohydrates (12.88%), calcium (0.03g), phosphorus (0.03g), iron (0.8g), beta carotene (70 IU), vitamin C (50-100mg/100g pulp) etc. Ripe ber fruits are eaten fresh. Fruits is also dried and used as dessert. It gives an excellent product when candy. Other processed products made are ber butter, ber juice or squash, RTS *etc.*

Formatted: Font: Italic

Plant bio-regulators and micronutrient such as Promalin, GA₄₊₇ and borax play an important role in increase fruits size, fruits set, fruit yield, self-life and quality improvement. Promalin plays an important role for increase the size and shape of the fruit. Promalin containing gibberellins and cytokine promotes both cell division and cell enlargement.

GA₄₊₇ plays an important role for induce fruit elongation and to increase the size and quality, fruit set, changing fruit shape and size, increasing weight of fruit,

Boron is for most important in the micro nutrients. Boron also plays important role for nucleic acid and lignin synthesis, pollen tube growth, it is associated with carbohydrate and fat metabolism and translocation and transformation of sugars.

MATERIALS AND METHODS

The present study was conducted on twenty eight years old uniform tree of ber cv. Gola, planted at a distance of 8 m x 8 m in Main Experimental Station of Horticulture, Acharya Narendra Deva University of agriculture & Technology, Narendra Nagar, Kumarganj, Ayodhya. (26^o.47^oN latitude, 82.12^oE longitude of 113 meters altitude away from mean sea stratum) during 2021-22 and 2022-23. Plants growth regulators and micro-nutrient, viz. ~~promalin~~Promalin (0.13% and 0.15%), GA₄₊₇ (10 ppm and 15 ppm) and borax (0.5%). Experiment was laid out in randomized block design (RBD) with three replication per treatment and considering one tree per replication. Data on fruit drop was recorded periodically from October to March. Ber fruits from sprayed plants were harvested month of March and analyzed for quality attributes.

Ten randomly selected fruits from each replication were used to assess quality attributes of ber. ~~Physical-physical~~ parameters like fruit size in term of length and breadth (cm) were measured with help of vernier’s calliper and expressed in centimetres. Fruit weight were recorded with help of digital weighing machine and expressed in gram. Among bio-

chemical parameters, TSS content was determined with hand refractometer and expressed in per cent (Ranganna, 1986). Acidity was expressed as per cent of citric acid as method given by Kaur, A. and Kalia, M. (2017). Ascorbic acid content was expressed in mg/100g of pulp and was estimated using 2,6-dichlorophenol indophenol dye by visual titration method (AOAC 1975). Total sugar reducing, non-reducing expressed in per cent was estimated by method suggested by AOAC (1975). The experiment data were analyzed following RBD to test the significance of using the Statistix-10 software. The differences in quantified concentrations were evaluated using F test at $P < 0.05$. Statistical analysis was done as described by Gomez and Gomez (1984).

RESULTS AND DISCUSSION

Fruit Size (cm)

Fruit Length (cm)

A perusal of The data on fruit length presented in (Table 1.) indicated that all treatments have a significant effect in improving the fruit length over control. The maximum fruit length (3.87cm and 3.92 cm) was recorded with T_8 -0.15% Promalin + 0.5% Borax, which was significantly superior over the rest of the treatments and at par with T_{10} -15 ppm GA_{4+7} +0.5% Borax (3.79 cm & 3.84 cm) followed by T_9 -10 ppm GA_{4+7} + 0.5% Borax (3.64cm and 3.69 cm), T_5 -15 ppm GA_{4+7} (3.54 cm and 3.59 cm) and T_7 -0.13% Promalin+0.5% Borax (3.39cm and 3.43cm), T_3 - 0.15 % Promalin (3.28cm and 3.32cm), T_4 - 10 ppm GA_{4+7} (3.22cm and 3.25cm), T_2 - 0.13 % Promalin (3.13 and 3.17), T_6 - 0.5 % Borax (3.10 & 3.14 cm) respectively. While minimum fruit length (3.08 cm and 3.11cm) was recorded with control during 2021-22 and 2022-23, respectively.

Formatted: Font: Bold

Fruit breadth (cm)

A perusal of The data on fruit length presented in (Table 1.) indicated that all treatments have a significant effect in improving the fruit breadth over control. The maximum fruit breadth (3.38 cm and 3.41 cm) was recorded with T_8 - 0.15% Promalin+0.5% Borax, which was significantly superior over the rest of the treatments and at par with T_{10} 15 ppm GA_{4+7} +0.5% Borax (3.33 cm and 3.36 cm) and T_9 -10 ppm GA_{4+7} + 0.5% Borax (3.32 cm and 3.35 cm), followed by T_5 -15 ppm GA_{4+7} (3.30 cm and 3.33 cm), T_7 - 0.13% Promalin+0.5% Borax (3.26 cm and 3.29 cm), T_3 - 15ppm Promalin (3.22 and 3.25 cm), T_4 -10 ppm GA_{4+7} (3.18cm and 3.21cm) and T_2 -0.13 % Promalin (3.10 cm and 3.13cm), T_6 - 0.5%

Formatted: Font: Bold

Borax (3.00cm and 3.03cm), respectively. While minimum fruit breadth (2.60 cm and 2.63 cm) was recorded with control during 2021-22 and 2022-23, respectively.

Fruit size was reported to be a genetic characteristic of the cultivars. The increase in size of fruit (length and width) over control as a result of foliar application of borax , Promalin_(6BA_) and GA₄₊₇ in present investigation might be due to their involvement in cell division, cell elongation, increased volume of intercellular spaces in the mesocarpic cells (Brahmachari *et al.* 1997), Promalin , GA₄₊₇ is also reported to promote growth by increasing plasticity of the cell wall followed by the hydrolysis of starch into sugar which reduces the cell water potential, resulting in the entry of water into the cell and causing elongation (Kassem *et al.*, 2011).

Increased in fruit size with plant growth regulators and micro-nutrient application has also been reported by Stenbridge and Morell (1972) reported that maximum fruit size application of promalin @ 50 ppm. Unarth (1974) foliar application of 25 and 50ppm promalin sprayed in apple, Cibulsky (1977); Looney (1979); Comai *et al.* (1981) in apple, Curraey and Williams (1982); Eccher and Maffi (1986) in apple, Pal (1997) in apple.

Fruit weight (g) / fruit

A perusal of data on fruit weight presented in (Table1) indicated that all treatments have a significant effect in improving the fruit weight over control. The maximum fruit weight (27.69 g and 28.89 g) was recorded with T₈ - 0.15% Promalin+0.5% Borax, which was significantly superior over the rest of the treatments and at par with T₁₀ -15 ppm GA₄₊₇ +0.5% Borax (26.30 g in 2021-22), followed by T₉ -10 ppm GA₄₊₇ + 0.5% Borax (25.87g,) T₅ -15 ppm GA₄₊₇ (23.96 g and 25.16 g),T₇ - 0.13% Promalin+0.5% Borax (23.40g and 24.60g) , T₃- 15ppm Promalin (23.24g and 24.44g),T₄ -10 ppm GA₄₊₇ (22.83 g and 24.03 g), T₂-0.13 % Promalin (21.97g_ and 23.17g) respectively. While minimum fruit weight (19.89g_8 g and 21.09) was recorded with control during 2021-22 and 2022-23, respectively.

Table 1.1. Effect of foliar feeding of plant growth regulators and micro-nutrient on Fruit size, and Fruit weight of ber fruit

Treatment	Parameters					
	Length (cm)		Breadth (cm)		Fruit weight (g)/fruit	
	2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T ₁ - Control	3.08 ^f	3.11 ^e	2.60 ^f	2.63 ^f	19.89 ^f	21.09 ^f
T ₂ - 0.13 % Promalin	3.13 ^{ef}	3.17 ^{de}	3.10 ^d	3.13 ^{de}	21.97 ^{de}	23.17 ^{de}

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold, English (United

Formatted: Font: Bold

Formatted: Font: Bold, English (United States)

T₃ - 0.15 % Promalin	3.28 ^{cd}	3.32 ^{cd}	3.22 ^c	3.25 ^{bcd}	23.24 ^{cd}	24.44 ^c
T₄ - 10 ppm GA₄₊₇	3.22 ^{de}	3.25 ^{de}	3.18 ^c	3.21 ^{cd}	22.83 ^{cde}	24.03 ^{cd}
T₅ - 15 ppm GA₄₊₇	3.54 ^b	3.59 ^b	3.30 ^b	3.33 ^{abc}	23.96 ^c	25.16 ^c
T₆ - 0.5 % Borax	3.10 ^f	3.14 ^e	3.00 ^e	3.03 ^e	21.36 ^{ef}	22.56 ^e
T₇ - 0.13 % Promalin + 0.5 % Borax	3.39 ^c	3.43 ^c	3.26 ^{bc}	3.29 ^{abc}	23.40 ^{cd}	24.60 ^c
T₈ - 0.15 % Promalin + 0.5 % Borax	3.87 ^a	3.92 ^a	3.38 ^a	3.41 ^a	27.69 ^a	28.89 ^a
T₉ - 10 ppm GA₄₊₇ + 0.5 % Borax	3.64 ^b	3.69 ^b	3.32 ^{ab}	3.35 ^{abc}	25.87 ^b	27.07 ^b
T₁₀ -15 ppm GA₄₊₇ + 0.5 % Borax	3.79 ^a	3.84 ^a	3.33 ^{ab}	3.36 ^{ab}	26.30 ^{ab}	27.50 ^b
LSD (p≤0.05)	0.113	0.155	0.079	0.135	1.699	1.199

Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)
Formatted: Font: Bold
Formatted: Font: Bold, English (United)

Increase in fruit weight with the spray of PGRs and micro-nutrient might be due to faster loading and mobilization into fruits (Brahmachari *et al.* 1997) and involvement in hormonal metabolism, increased cell division and expansion of cell. This may also be attributed to greater photosynthetic activity, resulting the increased production and accumulation of carbohydrates and favourable effect on vegetative growth and retention of fruits, which might have increased size and weight.

The present findings regarding the increase in fruit size and fruit weight with application of ~~promalin~~Promalin and GA₄₊₇ may be due to cell division and cell elongation. These results in close conformity with results of Looney *et al.* (1992), McArtney (2010), Chen *et al.* (2012), Ito *et al.* (2015), Sekhar (2012) and Al-madhagi *et al.* (2012) in strawberry. Kumar *et al.* (2003) and Yildirim *et al.* (2014) also found significant increase in fruit weight and size with application of ~~promalin~~Promalin and GA₄₊₇ ~~in~~ in fruit crops.

TSS (Brix⁰)

~~A perusal of~~The data ~~depicted~~ indepicted in Chart 1 that all treatments have a significant in improving the TSS. The maximum TSS (19.42_& 19.49) was recorded under the treatment of **T₈**- 0.15 % Promalin+0.5 % Borax, which was statistically *at par* with **T₁₀**- 15 ppm GA₄₊₇ + 0.5 % Borax_(19.02 and 19.09) and **T₉**-10ppm GA₄₊₇ + 0.5 % Borax_(18.76 and 18.83), followed by **T₇**- 0.13 % Promalin+0.5% Borax (18.41 and 18.48), **T₆**- Borax 0.5% (18.03 and 18.10), **T₅**- 15ppm GA₄₊₇ (17.58 and 17.65), **T₃**- 0.15 % Promalin (17.20

and 17.27), T₄- 10 ppm GA₄₊₇(16.56 and 16.63) and T₂- 0.13 % Promalin (15.83 and 15.90), while minimum TSS (14.80 and 14.87) was recorded with control during 2021-22 and 2022-23, respectively.

Increased TSS due to metabolic modification of starch and pectin, hydrolysis of complex polysaccharides, metabolite synthesis, and fast sugar translocation from leaves to developing fruits, foliar application of PGRs and micronutrients enhances total soluble solids (TSS) content in fruits. Total soluble solids (TSS) are also increased by ~~promalin~~Promalin and boron, with a maximum TSS of 0.5% Borax and 0.15% Promalin applied topically. These results are also in close conformity with those of Jindal *et al.* (2004) in apple, Pandey and Kumar (2022) and Pal *et al.* (2022) in ber cv. Gola, ~~Mentalti~~Montalti *et al.* (1984) in apple, Canli *et al.* (2009) in pear, Wismer *et al.* (1995) in apples, Cheolku *et al.* (2000) in pear, and Sharma (2001) in apple cv. Starking Delicious.

Formatted: Font: Italic




Chart 1: Effect of foliar feeding of plant growth regulators and micro-nutrient on TSS, Acidity and Ascorbic acid of ber fruit.

Acidity (%)

A perusal of data Chart 1 revealed that all treatments have a significant in improving the ~~Acidity per~~Acidity per cent. The minimum Acidity per cent (0.24 and 0.23) was recorded under the treatment of T₈- 0.15% Promalin + 0.5% Borax, which was statistically *at par* with T₁₀- 15 ppm GA₄₊₇ + 0.5 % Borax (0.25 and 0.24) , followed by T₉-10 ppm GA₄₊₇ + 0.5% Borax (0.26 and 0.25), T₇-0.13 % Promalin+ 0.5% Borax (0.27 and 0.26), T₆- Borax 0.5% (0.28 and 0.27), T₅- 15 ppm GA₄₊₇(0.29 and 0.28), T₃- 0.15 % Promalin (0.29 and 0.28), T₄-

10 ppm GA₄₊₇(0.30 and 0.29), T₂- 0.13% Promalin, while maximum Acidity (0.32 and 0.31) was recorded with control during 2021-22 and 2022-23, respectively.

The decrease in the acidity might be due to higher accumulation of sugars in fruits, better translocation of sugars into fruit tissues and conversion of organic acids into sugars by foliar application of PGRs and micro-nutrient. The reduction in the acidity under boric acid treatment might be owing to increased TSS of the fruits. These results also elucidate the finding of Jindal *et al.* (2004) in apple cv. Starking Delicious, Canli *et al.* (2009) in “Deveci” pear, Pandey and Kumar (2022), Pal *et al.* (2022) in ber cv. Gola, Khalid *et al.* (2012) in Kinnow.

Ascorbic acid (mg)

A perusal of data **Chart 1** revealed that all treatments have a significant in improving the Acidity per cent. The maximum Ascorbic acid (85.33 and 86.02) was recorded under the treatment of T₈-0.15 % Promalin+0.5% Borax which was statistically *at par* with T₁₀-15 ppm GA₄₊₇ + 0.5 % Borax (84.71 and 84.62) followed by T₉-10 ppm GA₄₊₇ + 0.5 % Borax (83.69 and 84.14), T₇-0.13% Promalin + 0.5 % Borax (83.43 and 84.12), T₆ -Borax 0.5% (82.52 and 83.87), T₅- 15ppm GA₄₊₇(80.37 and 81.39), T₃-0.15% Promalin (79.83 and 80.85), T₄- 10 ppm GA₄₊₇ (78.20 and 79.55) and T₂-0.13 % Promalin (77.74 and 78.43), while minimum ascorbic acid (76.82 and 77.84) was recorded with control during 2021-22 and 2022-23, respectively.

Formatted: Font: Bold

The increase in ascorbic acid content might be speculated due to increased activity of enzyme responsible for the synthesis of the ascorbic acid precursor i.e., glucose-6-phosphate or inhibition of its conversion into ~~dehydro~~ascorbic acid by enzyme ascorbic acid oxidase or both and also the reduction in the rate of respiration by these chemicals (Brahmachari and Rani 2001). Similar result have been reported by Pal *et al.*,(2022) and Pandey and Kumar (2022) in ber cv. Gola, Khalid *et al.*, (2012) in Kinnow and Cheolku *et al.* (2000) in pear.

Reducing sugar (%)

A perusal of data **Chart 2** revealed that all treatments have a significant in improving the Acidity percent. The maximum reducing sugar (6.23 and 6.25) was recorded under the treatment of T₇ -0.13 % Promalin + 0.5 % Borax followed by T₁₀- 15 ppm GA₄₊₇ + 0.5 % Borax (5.96 and 5.98), T₉-10 ppm GA₄₊₇ +0.5 % Borax (5.84 and 5.87), T₈- 0.15 % Promalin +0.5 % Borax (5.73 and 5.75), T₆ -Borax 0.5% (5.59 and 5.62), T₅- 15 ppm GA₄₊

Formatted: Font: Bold

T₇(5.43 and 5.45), T₃- 0.15 % Promalin(5.30 and 5.32), T₄- 10 ppm GA₄₊₇(5.07 and 5.14) and T₂-0.13 % Promalin (4.82 and 4.84), while minimum reducing sugar (4.47 and 4.49) was recorded with control during 2021-22 and 2022-23, respectively.

Foliar spray of PGRs and micro-nutrients is helpful to increase the reducing sugar level, which could be due to accumulation of more carbohydrates to the fruit which results the better accessibility of nutrition for developing fruits and increases the reducing sugar level of fruits in present investigation. The result in conformity with those of Pandey and Kumar (2022) and Pal *et al.*(2022) in ber cv. Gola, Jindal *et al.* (2004) in apple, Canli *et al.* (2009) in pear and Khalid *et al.*, (2012) in Kinnow and Cheolku *et al.* (2000) in pear.

Non-reducing sugar (%)

A perusal of data **Chart 2** revealed that all treatments have a significant in improving the Acidity per cent. The maximum Non-reducing sugar (7.92 and 7.95) was recorded under the treatment of T₇- 0.13 % Promalin +0.5 % Borax followed by T₁₀- 15 ppm GA₄₊₇ + 0.5 % Borax (7.87 and 7.90), T₉- 10 ppm GA₄₊₇ + 0.5 % Borax (7.72 and 7.74), T₈- 0.15 % Promalin + 0.5 % Borax (7.56 and 7.59), T₆ -Borax 0.5% (7.39 and 7.41), T₅- 15ppm GA₄₊₇ (7.17 and 7.20), T₃- 0.15 % Promalin (6.99 and 7.02), T₄-10 ppm GA₄₊₇ (9.69 and 6.72) and T₂- 0.13 % Promalin (6.37 and 6.40), while minimum non-reducing sugar (5.89 and 5.92) was recorded with control during 2021-22 and 2022-23, respectively

Formatted: Font: Bold




Chart 2: Effect of foliar feeding of plant growth regulators and micro-nutrient on Reducing sugar, Non-reducing sugar and Total sugar of ber fruit.

The possible reason for increase in non-reducing sugar content of fruits due to hydrolysis of polysaccharides to simpler form i.e. mono and disaccharides and better transportation of

nutrients to plant from leaves to their place of utilization. These results corroborate the earlier records of Pandey and Kumar (2022) and Pal *et al.*(2022) in ber cv. Gola, Khalid *et al.*, (2012) in Kinnow, Jindal *et al.* (2004) in apple and Cheolku *et al.* (2000) in pear.

Total sugar (%)

A perusal of the data **Chart 2** revealed that all treatments have a significant in improving the Acidity per cent. The maximum Total sugar (14.15 and 14.20) was recorded under the treatment of **T₇**- 0.13 % Promalin + 0.5 % Borax followed by **T₁₀**- 15 ppm GA₄₊₇ + 0.5 % Borax (13.83 and 13.88), **T₉**-10 ppm GA₄₊₇ + 0.5 % Borax (13.56 and 13.61), **T₈**-0.15 % Promalin + 0.5 % Borax (13.29 and 13.34), **T₆** -Borax 0.5% (12.98 and 13.03), **T₅**- 15 ppm GA₄₊₇ (12.60 and 12.65), **T₃**- 0.15 % Promalin (12.29 and 12.34), **T₄**- 10 ppm GA₄₊₇ (11.76 and 11.81) and **T₂**- 0.13% Promalin (11.19 and 11.24), while minimum Total sugar (10.36 and 10.41) was recorded with control during 2021-22 and 2022-23, respectively.

The possible reason for increase in total sugar content may be due to hydrolysis of starch yielding mono and disaccharide, which owned a simplest form of sugar and boron, **Promalin** and GA₄₊₇ probably augmented the conversion of starch. Gauch and Dugger (1953) reported that the improvement in total sugar content may be due to more translocation of sugars from leaves to developing fruits. Our results are also in line with the finding of Pandey and Kumar (2022) and Pal *et al.*,(2022) in ber cv. Gola, Chaudhary *et al.*(2018) in **Anola**, Cheolku *et al.* (2000) in **Pear**, Wismer *et al.* (1995) in **Apples** and Sharma (2001) in **Apple** cv. Starking Delicious.

CONCLUSION

It may be concluded that the application of different growth regulators resulted that improve the characters of ber *i.e.* Fruit weight, fruit size, TSS, acidity, ascorbic acid, reducing sugar, non-reducing sugar and total sugar. The sugar and ascorbic acid, parameters were significantly affected by the application of 0.15% Promalin + 0.5% Borax (**T₈**) and **T₁₀**- 15 ppm GA₄₊₇ + 0.5 % Borax. Based on cost benefit ratio it may be concluded that **T₁₀**- 15 ppm GA₄₊₇ + 0.5 % Borax could be recommended to farmers for commercially increasing yield and quality of ber. Normally people don't prefer Gola variety ber over Apple ber due to

Formatted: Font: Bold

small size but with the application of 15 ppm GA₄+7 + 0.5 % Borax we can increase fruit size in a very economical way resultant consumer's preference will also increase.

Formatted: Indent: First line: 0.5", Space Before: 0 pt

REFERENCES:

1. A.O.A.C., **1975**. Official method of analysis. Assoc. of Official Analytical Chemists, Benjamin Franklin Station, Washington D.C., 12th edition, 575.
2. Al-madhagi, IAH.; Hasan, SMZ.; Ahmad, A.; Zain, AM. and Yusoff, A. **2012**. Influence of exogenous hormone on the flowering of strawberry (*Fragaria x ananassa* Duch). *Journal of Biology Agriculture and Healthcare*. 2(4):46-52.
3. Anonymous, **2021-22**. Horticulture Data Base, National Horticulture Board, Gurgaon, Ministry of Agriculture, India.
4. Brahmachari, V.S and Rani, R. **2001**. Effect of growth substance on productivity, cracking, ripening and quality of fruits in litchi. *Orissa journal of Horticulture*,29(1):41-45.
5. Canli,AF.; Pektas, M.; and Kelen, M. **2009** . Effects of Pre-harvest Plant Growth Regulator Sprays on Fruit Quality of 'Deveci' Pear (*Pyrus communis* L.). *Journal of Applied Biological Sciences*. 3 (1): 81-84.
6. Chaudhary, M.; Singh, D. P.; Singh, J. P. and Tomar, S. **2018**. Effect of foliar application of PGRs and mineral nutrients on physico-chemical characteristics of Aonla fruit cv. NA-7. *Journal of Pharmacognosy and Phytochemistry*7(4), 2719-2723.
7. Chen, X.; Bao, J. Chen, Y.; Chen, T.; Zhang, C.; and Huang, X. **2012**. Effect of hormone treatments on deformed fruit development in pear. *African journal of biotechnology*, 11(44), 10207-10209.
8. Cheolku, Y.; Seonkyukim, S.L. and Kwansoon,C. **2000**. Effect of application time of promalin on tree growth and fruit quality of pears. *Journal of Korean Society for Horticultural Science* 41(3): 289-291.
9. Cibulsky, RJ. **1977**. Effect of bloom application of promalin on apple fruit. *Proceedings of the Plant Growth Regulator Society of America* 4: 223-230.
10. Comai, M.; Widemann,L. and Gianotti,G. **1981**. The effect of gibberellins and promalin on fruit shape and quality of apple. *Esperienze Ricerche Stazione Sperimentale Agraria Forestale Adige*. 11: 47-61.
11. Curracy, E A. and Williams, M W. **1982**. Promalin or GA₃ increase pedicel and fruit length and leaf size of 'Delicious' apple treated with paclobutrazol. *Horticulture Science*. 18(2): 214-215.
12. Eccher, T. and Maffi, A. **1986**. Treatments for prevention of "Golden Delicious" russetting. *Acta Horticulturae*. 179:767-70.
13. Gauch, H.G. and Dugger,W.W.J **1953**. The physiological action of boron in higher plants review and interpretation. Technical Bulletin, University Maryland Agriculture Experiment Station, USA, 80p.

Formatted: Font: Bold

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Black, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

Formatted: Font: Bold, English (United States)

14. Gomez, K.A. and Gomez, A.A. **1984**. *Statistical Procedures of Agricultural Research (2nd Edn.)*, John Willey and Sons Inc., New York. **Formatted: Font: Bold, English (United)**
15. Ito, A.; Sakamoto, D.; Itai, A.; Nishijima, T.; Oyama-Okubo, N.; Nakamura, Y.; Moriguchi, T. and Nakajima, I. **2015**. Effects of GA₃₊₄ and GA₄₊₇ application either alone or combined with prohexadione-Ca on fruit development of Japanese pear 'Kosui'. *The Japanese Society for Horticulture Science*, 85(3), 201-208. **Formatted: Font: Bold, English (United)**
16. Jindal, K.K.; Pal, S.; Chauhan, P.S. and Mankotia, M.S. **2004**. Effect of promalin and mixtadol on fruit growth, yield efficiency and quality of "Starking Delicious" apple. *Acta Horticulturae*. 636: 533-536. **Formatted: Font: Bold, English (United)**
17. Khalid S.; Malik, A.U.; Khan, A.S. and Jamil, A. **2012**. Influence of exogenous applications of plant growth regulators on fruit quality of Young 'Kinnow' Mandarin (*Citrus nobilis* × *C. deliciosa*) Trees. *Int. J. Agric. Biol.*; 14(2):229-234. **Formatted: Font: Bold, English (United)**
18. Kumar, J.; Rana, SS.; Verma, HS. and Jindal, KK. **2003**. Effect of promalin on fruit shape, set, yield and quality of apple cv. Starking Delicious. *Indian Journal of Horticulture*. 60(2): 143-146. **Formatted: Font: Bold, English (United)**
19. Looney, N E. **1979**. Some effects of GA₄₊₇ and BA on fruit weight, shape, quality and calcium content of 'Spartan' apple. *Journal of the American Society for Horticultural Science*. 104(3): 389-391. **Formatted: Font: Bold, English (United)**
20. McArtney, S. J. **2010**. Exogenous gibberellin affects biennial bearing and the fruit shape of 'Braeburn' apple. *New Zealand Journal of Crop and Horticultural Science*, Vol. 22: 343-346. **Formatted: Font: Bold, English (United)**
21. Montalti, P.; Cristofferi, B. and Sansavini, S. **1984**. Effect of GA₄₊₇ and shoot pinching on hormonal levels in conference pear. *Acta Horticulturae*. 161: 163-170. **Formatted: Font: Bold, English (United)**
22. Pal, R.; Kumar, A.; and Kumar, A. **2021**. Effect of foliar feeding on chemical attributes of Ber (*Zizyphus mauritiana* L.) under Sodic soil. *International Journal of Chemical Studies*. 9 (1), 1558-1560. **Formatted: Font: Bold, English (United)**
23. Pal, S.J. **1997**. Effect of plant bio regulators on fruit growth, fruit yield and quality of Stalking Delicious apples. M.Sc. Thesis Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, (H.P.). **Formatted: Font: Bold, English (United)**
24. Pandey, A.; and Kumar, A. **2022**. Effect of ZnSO₄, Borax and GA₃ on quality and yield of ber (*Zizyphus mauritiana* Lamk.) Fruits Cv. Gola. *The Pharma Innovation Journal*. 11(8): 1788-1790. **Formatted: Font: Bold, English (United)**
25. Pareek, B.S. **1983**. The Ber- 1st edition *Indian Council of Agricultural Research*, New Delhi. pp 31-36. **Formatted: Font: Bold, English (United)**
26. Ranganna, S. **(1986)**. *Manual of Analysis of fruits and vegetable products*, Tata Mc Graw-Hill Publishing Company Ltd; New Delhi, pp.1-30. **Formatted: Font: Bold**
27. Sharma, G. **2001**. Effect of pre bloom foliar applications of nutrients and plant bio regulators on fruit set and yield in apple. M.Sc. Thesis Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, (H.P.). **Formatted: Font: Bold, English (United)**
28. Stembridge, GE. and Morrell, G. **1972**. Effect of gibberellins and 6-benzyl-adenine on the shape and fruit set of "Delicious" apples. *Journal of the American Society for Horticultural Science*. 97:464-67. **Formatted: Font: Bold, English (United)**

29. Unrath, CR. **1974**. The commercial implications of gibberellin A₄ plus benzyladenine for Improving shape and yield of 'Delicious' apples. *Journal of the American Society for Horticultural Science* .99(4): 381-384.
30. Wismer, P T.; Proctor, J T A. and Elfving, DC. **1995**. Benzyladenine affects cell division and cell size during apple fruit thinning. *Journal of the American Society for Horticultural Science*. 120: 802-807.
31. Yildirim, FA.; Kepenek, G.; San, B.; Yildirim, AN. and Kacal, E. **2014**. Effects of Ba + GA₄₊₇ treatments on fruit quality in “Fuji” apple variety. *Turkish Journal of Agricultural and Natural Sciences*.2: 1387-1390.

Formatted: Font: Bold, English (United)

Formatted: Font: Bold, English (United)

Formatted: Font: Bold, English (United)

Formatted: Font: Bold, English (United)

UNDER PEER REVIEW