

Efficacy of selected insecticides with Botanicals and Bio-agents against Diamondback moth, *Plutella xylostella* (L.) in cabbage, *Brassica oleracea var capitata* (L.)

ABSTRACT

Field trial was conducted during *rabi* season 2022-2023 at Central Research Farm (CRF) , SHUATS. The experiment was laid out in Randomised block design with eight treatments each replicated thrice using a variety Green Soccer (546). The treatments are T₁ Chlorantraniliprole 18.5 % SC, T₂ Indoxacarb 14.5% SC, T₃ Flubendiamide 39.3 SC, T₄ *Bacillus thuringiensis* (1×10^8 CFU/ml), T₅ *Metarhizium anisopilae* (1×10^8 CFU/ml), T₆ NSKE 5%, T₇ Spinosad 45% SC and along with an untreated T₈ control against *Plutella xylostella* in Cabbage. Mean reduction in the larval population of Diamond back moth per plant revealed that all treatments significantly superior over control (4.200). Among all the treatments T₁ Chlorantraniliprole 18.5 % SC recorded highest larval reduction with mean (1.000) of *Plutella Xylostella* followed by T₂ Indoxacarb 14.5% SC (1.178), T₇ Spinosad 45% SC (1.311) ,T₃ Flubendiamide 39.3 SC (1.322) , T₅ *Metarhizium anisopilae* (1×10^8 CFU/ml) (1.456), T₄ *Bacillus thuringiensis* (1×10^8 CFU/ml) (1.522) and T₆ NSKE 5% (1.734) and least recorded in T₈ control (4.200). Highest yield 336 q/ha as well as B:C ratio (1:5.35) was obtained from the treatment Chlorantraniliprole 18.5% SC followed by T₂ Indoxacarb 14.5% SC (1:5.28), T₇ Spinosad 45% SC (1:4.85) T₃ Flubendiamide 39.3 SC (1:4.83) , T₅ *Metarhizium anisopilae* (1×10^8 CFU/ml) (1:4.18), T₄ *Bacillus thuringiensis* (1×10^8 CFU/ml) (1:4.71) and T₆ NSKE 5% (1:3.52) superior over the control (1:2.87).

Key words: *Bacillus thuringiensis*, Chlorantraniliprole, Cost benefit ratio, *Metarhizium anisopilae*, NSKE, *Plutella xylostella*.

INTRODUCTION

“Cabbage, (*Brassica oleracea* var. *capitata*) an herbaceous plant of Family Brassicaceae. It is one of the most popular cole vegetables grown in India. It is originated in Europe. It is commonly used fresh as boiled vegetables cooked in curries, process, salad and used in herbal medicine. Cabbage juice can reduce constipation and has also been used as a laxative, as an antidote to mushroom poisoning, or a treatment for hangovers and headaches. In fact, cabbage has historically been used to stop sunstroke or to relieve fevers”. [25]

“Regular consumption of dark green leafy vegetables is highly recommended because of their potential in reducing chronic disease and glucosinolates in cabbage reduced risk of cancer induction and development. It is known to possess medicinal properties and its enlarged terminal buds is a rich source of Ca, P, Na, K, S, vitamins A, Vitamin C and dietary fiber. It is said to be good for person suffering from diabetes”. (Khan and Tayde, 2022).

“Leaves are low in calories (27 per cent), fat (0.1 per cent) and carbohydrates (4.6 per cent). It is a good source of protein (1.3 per cent) which contains all essential amino acids, particularly Sulphur containing amino acids. Cabbage is an exceptional source of minerals such as calcium (39 mg), iron (0.8 mg), magnesium (10 mg), sodium (14.1 mg), potassium (114 mg) and phosphorus (44 mg). It has considerable amounts of β carotene provitamin A, ascorbic acid, riboflavin, niacin, and thiamine. Ascorbic acid content varies from 30-65 mg per 100 g fresh weight. Flavor in cabbage leaves is due to the glycoside sinigrin. It contains goitrogens which cause enlargement of thyroid glands. The Lumbee tribe of North Carolina has traditionally used the leaves of *Brassica oleracea* in medicine that they believed to have cleansing qualities, as well as a mild laxative, an anti-inflammatory, and treatment for glaucoma and pneumonia”. (Jacquet et al., 2017).

“In 2020, world production of cabbages was 71 million tones, led by China with 48% of the world total (table). Other substantial producers were India, Russia, and South Korea. (UN Food and Agriculture Organization, Corporate Statistical Database)” (FAOSTAT, 2022). “India is the largest producer of cabbage after China. India accounts for 8755000 tons of productivity in an area of 388000/ha. In India, Uttar Pradesh accounts for production of 5.7 million tones in an area of 0.72 million ha”. (Kommoji and Tayde, 2022).

“The major insect pests, which cause maximum yield losses in cabbage are Diamond back moth (*Plutella xylostella* L), cabbage butterfly, (*Pieris brassicae* L), cabbage aphid (*Brevicoryne brassicae* L.), cabbage semilooper (*Trichoplusia ni*), leaf webber (*Crocidolomia binotalis*), cabbage head borer, (*Hellula undalis* Fab.). Diamondback moth is the most destructive pest in cabbage growing areas and the yield loss were reported up to 52% in India” (Krishnamoorthy, 2004).

“However, the set back to optimum cabbage production is the attack of insect pests, the most important of which is the diamondback moth (DBM), *Plutella xylostella* which has become a single limiting factor in the production of quality heads. It is one of the most destructive insect pests of cruciferous vegetables, currently accounting for US\$2.7 billion worth of annual worldwide crop losses. Management of this pest depends largely on imposing heavy quantities of synthetic chemical pesticides all over the world”. (Maity *et al.*, 2015). The DBM has developed resistance to all major classes of insecticides. More management of this pest poses serious concern due to development of insecticide resistance to carbamate (Sun *et al.*, 2007) and synthetic pyrethroids (Liu *et al.*, 2000).

MATERIALS AND METHODS

Field experiment was conducted at Central Research Farm (CRF), Department of Entomology, SHUATS, Naini, Prayagraj, Uttar Pradesh, India in a Randomized Block Design with eight treatments replicated three times using the variety Green Soccer in a plot size of (2mx1m) at a spacing of (60x45cm) with a recommended package of practices excluding plant protection. The soil of the experimental site was well drained and medium high.

All of the insecticides used in the study were sprayed as foliar application. The eight different treatments were used with dosage consisting of T₁ Chlorantraniliprole 18.5% SC 0.5ml/L, T₂ Indoxacarb 14.5% SC 0.6ml/L, T₃ Flubendiamide 39.3 SC 0.2ml/L, T₄ *Bacillus thuringiensis* (1x 10⁸ CFU/ml) 2ml/L, T₅ *Metarhizium anisopilae* (1x10⁸CFU/ml) 4gm /L, T₆ NSKE 5gm/L, T₇ Spinosad 45% SC 0.5ml/L, T₈ Control. Two sprays were carried out at intervals of 15 days during the experiment to assess the effectiveness of pesticides. On five randomly chosen and tagged plants in each plot, pre and post -treatment observations on the percent damage of Diamondback moth infestation was made shortly before 24 hours and 3rd, 7th and 14th days, respectively.

Application of treatments was started when the threshold action level of one diamond back moth per plant. Spray was made using manually operated knapsack hand.

Preparation of insecticidal spray solutions:

The Insecticidal spray solution of desired concentration as per treatment was freshly prepared every time at the site of experimentation just before the start of spraying operations. The spray solution of a desired concentration was preparing by adopting the following formula,

$$V = \frac{(CXA)}{\%a.i}$$

Where,

V = Volume/ weight of formulated insecticide required.

C = Concentration required.

A = Volume of solution to be prepared.

a.i. = given percentage of active ingredient.

OBSERVATION:

The population of caterpillar was recorded on 5 plants randomly selected and tagged from each plot. After that mean of three replications was calculated for each treatment and the same was done with the untreated plot. The population of *Plutella xylostella* was recorded 1 day before spraying and on 3rd day, 7th day and 14th day after insecticidal application. The average population of larvae of diamondback moth per plant was worked out for statistical analysis. While recording yield data, only marketable heads was considered. Yield obtained from net plot was converted into per hectare.

Cost benefit ratio of treatments :

Gross returns was calculated by multiplying total yield with market price of the produce. Cost of cultivation and cost of treatments was deducted from the gross returns, to find out returns and cost benefit of ratio by following formula,

$$\text{BCR} = \frac{\text{Gross Returns}}{\text{Cost of Cultivation}}$$

Where,

Benefit Cost Ratio

Table:1 Efficacy of selected Insecticides with Botanicals and Bio-agents against Diamondback moth, *Plutella xylostella* (L.) in cabbage , *Brassica oleracea var capitata* (L.)

S.No	Treatments	Larval Population of diamondback moth								Overall mean	Yield (q/ha)	C:B Ratio
		First Spray				Second spray						
		1 DBS	3 DAS	7 DAS	14 DAS	1 DAS	3 DAS	7 DAS	14 DAS			
T1	Chlorantraniliprol 18.5%SC	3.333	2.000 ^c	0.600 ^c	1.067 ^d	1.067 ^d	0.800 ^d	0.600 ^c	0.733 ^c	1.000 ^b	336	1:5.35
T2	Indoxacarb 14.5% SC	3.000	2.067 ^c	0.733 ^{de}	1.400 ^{cd}	1.400 ^{cd}	0.867 ^d	0.733 ^{de}	1.000 ^{bc}	1.178 ^b	325	1:5.28
T3	Flubendiamide 39.3 SC	2.800	2.200 ^{bc}	1.000 ^{bcd}	1.467 ^c	1.467 ^c	1.138 ^{bcd}	1.000 ^{bcd}	1.200 ^{bc}	1.322 ^b	265	1:4.83
T4	<i>Bacillus thuringiensis</i> (1x10 ⁸ CFU/ml)	2.867	2.267 ^{bc}	1.133 ^{bc}	1.600 ^{bc}	1.600 ^{bc}	1.467 ^{bc}	1.133 ^{bc}	1.400 ^b	1.522 ^b	241	1:4.71
T5	<i>Metarhizium anisopilae</i> (1x10 ⁸ CFU/ml)	2.867	2.333 ^{bc}	1.067 ^{bcd}	1.533 ^c	1.533 ^c	1.400 ^{bc}	1.067 ^{bcd}	1.267 ^b	1.456 ^b	217	1:4.18
T6	NSKE	3.333	2.600 ^b	1.267 ^b	1.933 ^b	1.933 ^b	1.533 ^b	1.267 ^b	1.400 ^b	1.734 ^b	192	1:3.52
T7	Spinosad 45% SC	3.000	2.133 ^{bc}	0.867 ^{cde}	1.400 ^{cd}	1.400 ^{cd}	1.067 ^{cd}	0.867 ^{cde}	1.333 ^b	1.311 ^b	318	1:4.85
T8	Control	2.733	3.533 ^a	4.600 ^a	3.867 ^a	3.867 ^a	4.467 ^a	4.600 ^a	5.000 ^a	4.200 ^a	140	1:2.87
	F-test	NS	S	S	S	S	S	S	S	S	--	
	C.D. (P=0.05)		0.480	0.338	0.389	0.389	0.446	0.383	0.495	0.894	--	
	C.V	8.978	11.451	13.291	12.449	12.499	15.988	15.533	16.945	22.048	--	

DBS**-Day Before Spray*, DAS***- Day After Spray **, NS= Non-Significant***, S-Significant ***

RESULTS AND DISCUSSION

The present study entitled, “Efficacy of selected insecticides with Botanical and Bio-agents against diamondback moth, *Plutella xylostella* (L.) in Cabbage, *Brassica oleracea var capitata* (L.)” The data so obtained through observation on various aspects were subjected to statistical analysis wherever necessary and the data was compiled. Results thus obtained are presented aspect wise here under, the data on the larval population of Diamondback moth *Plutella xylostella* in cabbage 3rd,7th and 14th after first spray revealed that all the chemical treatments, were significantly superior over control (3.711). Among all the treatments lowest larval population was recorded in T₁ Chlorantraniliprole 18.5 % SC (1.289), T₂ Indoxacarb 14.5% SC (1.489), T₇ Spinosad 45% SC (1.533) T₃ Flubendiamide 39.3 SC (1.533) followed by T₅ *Metarhizium anisopilae* (1x10⁸ CFU/ml) (1.666), T₄ *Bacillus thuringiensis* (1x10⁸ CFU/ml) (1.711) and T₆ NSKE 5% (2.067).

The data on the larval population of Diamondback moth *Plutella xylostella* in cabbage 3rd,7th and 14th day after Second spray revealed that all the chemical treatments were significantly superior over control. Among all the treatments lowest larval population was recorded in T₁ Chlorantraniliprole 18.5 % SC (0.711), T₂ Indoxacarb 14.5% SC (0.867), T₇ Spinosad 45% SC (1.089) T₃ Flubendiamide 39.3 SC (1.111) followed by T₅ *Metarhizium anisopilae* (1x10⁸ CFU/ml) (1.245), T₄ *Bacillus thuringiensis* (1x10⁸ CFU/ml) (1.333) and T₆ NSKE 5% (1.400) and control plot (4.689).

The data revealed on population of *Plutella xylostella* over control on Overall mean revealed that all the treatments were significantly superior over control (4.200). Among all the treatments minimum larval population was recorded in T₁ Chlorantraniliprole 18.5 % SC (1.000) similar findings made by **Kommoji et al.(2022)**, T₂ Indoxacarb 14.5% SC (1.178),similar findings made by **Pavithra et al.,(2016)**, T₇ Spinosad 45% SC (1.311) similar findings made by **Devi and Tayde, (2017)**, T₃ Flubendiamide 39.3 SC (1.322) similar findings made by **Lal et al., (2021)** followed by T₅ *Metarhizium anisopilae* (1x10⁸ CFU/ml) (1.456),similar findings made by **Devi and Tayde, (2017)**, T₄ *Bacillus thuringiensis* (1x10⁸ CFU/ml) (1.522) similar findings made by **Purushotam et al.,(2022)** and T₆ NSKE 5% (1.734) similar findings made by **Khan and Tayde, (2022)**.

All the insecticides were found very effective and significantly superior over control. The minimum larval population and the highest yield was recorded in T₁ Chlorantraniliprole 18.5 % SC (336q/ha), followed by T₂ Indoxacarb 14.5% SC (325q/ha), T₇ Spinosad 45% SC (318q/ha) T₃ Flubendiamide 39.3 SC (265q/ha) followed by T₅ *Metarhizium anisopilae* (1x10⁸ CFU/ml) (217q/ha), T₄ *Bacillus thuringeninsis* (1x10⁸ CFU/ml) (241q/ha) and T₆ NSKE 5% (192q/ha) and the lowest yield is recorded in T₀ Control (140q/ha) similar findings made by **Kumar and Kumar, (2020), Khan and Tayde, (2022), Kommoji et al.(2022).**

Cost benefit ratio (CBR)

When cost benefit ratio worked out, interesting result was achieved, among the treatment studied, the best and most economical treatment is in T₁ Chlorantraniliprole 18.5 % SC (1:5.35) similar findings made by **Kommoji et al.(2022)**, T₂ Indoxacarb 14.5% SC (1:5.28), similar findings made by **Pavithra et al.,(2016)**, T₇ Spinosad 45% SC (1:4.85) similar findings made by **Devi and Tayde, (2017)**, T₃ Flubendiamide 39.3 SC (1:4.83) similar findings made by **Lal et al., (2021)** followed by T₅ *Metarhizium anisopilae* (1x10⁸ CFU/ml) (1:4.18), similar findings made by **Devi and Tayde, (2017)**, T₄ *Bacillus thuringeninsis* (1x10⁸ CFU/ml) (1:4.71) similar findings made by **Purushotam et al.,(2022)** and T₆ NSKE 5% (1:3.52) similar findings made by **Khan and Tayde, (2022)** and control T₀ (1:2.87) similar findings made by **Kumar and Kumar, (2020), Khan and Tayde, (2022), Kommoji et al.(2022).**

CONCLUSION

From the present study finding, it was revealed that among all treatment T₁ Chlorantraniliprole 18.5%SC recorded best and proved best effective for Diamondback moth, (*Plutella xylostella*) followed by T₂ Indoxacarb 14.5% SC, T₇ Spinosad 45% SC and T₃ Flubendiamide 39.3 SC in controlling (*Plutella xylostella*) in cabbage (*Brassica oleracea*) on mean larval population. Therefore, the botanicals *i.e.*, T₅ *Metarhizium anisopilae* (1×10^8 CFU/ml), T₄ *Bacillus thuringiensis* (1×10^8 CFU/ml) and T₆ NSKE 5% may be useful in devising proper integrated pest management strategy against Diamond Back Moth (*Plutella xylostella*).

References

1. **Chandraker, M. K., Rana, D. K. and Saran, B. (2022).** Efficacy of botanicals and microbial agents for the management of diamondback moth (*Plutella xylostella*) in cabbage. *The Pharma Innovation journal*, **11**(6): 2348-2351.
2. **Devi, H. D. and Tayde, A. R. (2017).** Comparative efficacy of bio-agents and botanicals on the management of diamondback moth (*Plutella xylostella* Linn.) on cabbage under Allahabad agroclimatic conditions. *International Journal of Current Microbiology and Applied Sciences*, **6**(7): 711-716.
3. **Dotasara, S. K., Agrawal, N., & Swami, D. (2017).** Efficacy of some new insecticides against diamond back moth (*Plutella xylostella* L.) on cauliflower. *THE INDIAN SOCIETY OF AGRICULTURAL SCIENCE*, 227.
4. **Fletcher, T.B. (1914)** Some south Indian pests. *Superintendent Government Press, Madras*.
 - a. 5:65-66.
5. **Gaddam, N. R., Srivastava, V. K., Tayde, A. R. and Tripathi, A. (2021).** Comparative efficacy of microbials and botanicals against diamondback moth (*Plutella xylostella* Linn) on cabbage. *Journal of Entomology and Zoology Studies*. **9**(1): 497-499.
6. **Gomez, K.A. and A.A. Gomez, (1976).** Statistical procedures for agricultural research.
 - a. *Library of Congress Cataloging in Publication*, **630**(72):83-14556.
7. **Hamid, d. S. and kalaf , J. M. (2021)** Effect of some insecticides in controlling the diamond back moth larvae, *Plutella xylostella* (lepidoptera: plutellidae) of cabbage *Biochemical and Cellular Archives*, **21**(1): 611-615.
8. **Harika, G., Dhurua, S., Suresh, M. and Sreesandhya, N. (2019).** Evaluation of Certain Insecticides against Diamondback Moth (DBM) *Plutella xylostella* on Cauliflower. *International Journal of Bio-resource and stress Management*, **10**(1), 070076.
9. **Jakhar, M., Singh, S. K. and Choudhary, A. L. (2019).** Evaluate of bio-efficacy on different insecticides against diamondback moth on cauliflower. *Journal of Entomology and Zoology Studies*, **7**(1): 1077-1081.

10. **Kommoji, T. and Tayde, A. R. (2022).** Efficacy of Newer Selected Insecticides, *Beaveria bassiana* and Neem Oil against Diamond Back Moth *Plutella xylostella*(L.) in Cabbage (*Brassica oleraceavar capitata*). *International Journal of Plant and Soil Science*, **34**(20): 485491.
11. **Krishnamoorthy, A. (2004)** Biological control of diamondback moth *Plutella xylostella* (L.), an Indianscenario with reference to past and future strategies. *In Proceedingsof the International Symposium*, **2**(7):204-11.
12. **Kumar, A. N. and Kumar, A. (2020).** Comparative efficacy of certain chemicals with biopesticides against diamondback moth, *Plutella xylostella* (L.) in cabbage, *Brassica oleracea* (L.). *Journal of Entomology and Zoology Studies*, **8**(6): 1350- 1353.
13. **Lal, J., Swaminathan, R., Meena, A. K. and Nagar, R. (2021).** Field efficacy of biorational insecticides against major pestiferous insects of cabbage. *International Journal of Pest Management*, 1-7.
14. **Mahmoudvand, M., Garjan, A. S. and Abbasipour, H. (2011).** Ovicidal effect of some insecticides on the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: yponomeutidae). *Chilean Journal of Agricultural Research*, **71**(2): 226-230.
15. **Maity, L., Padhi, G., & Samanta, A. (2018).** Population dynamics and management of diamond backmoth *Plutella xylostella* (L.) in cabbage ecosystem of West Bengal. *International Journal of ChemicalStudies*, **6**(1): 381-385.
16. **Nayak, R., Naveen, J. B. and Pradeep, S. (2019).** Evaluation of insecticides against diamondback moth *Plutella xylostella* on cabbage during 2016. *International Journal of Chemical Studies*, **7**(6): 50-51.
17. **Patel, Y., Garg, V. K. and Jamliya, G. (2020).** Efficacy of insecticides against diamond back moth *Plutella xylostella* (L.) in cabbage (*Brassica oleracea* var. *capitata*). *Journal of Entomology and Zoology Studies*, **8**(3): 1896-1902.

18. Patil, R. H., Shaila, H. M. and Awaknavar, J. S. (2017). Management of diamondback moth, *Plutella xylostella* L. through biorationals in cabbage. *Mysore Journal of Agricultural Sciences*, **51**(1): 148-152.
19. Patra, B., Das, B. C., Alam, S. F., Dhote, V., Patra, S., Chatterjee, M. L. and Samanta, A. (2015). Evaluation of new insecticides against diamond back moth, *Plutellaxylostella* (L.) on red cabbage. *International Journal of Bio-resource and StressManagement*, **6**(2): 280-284.
20. Reddy, M. S., Sathua, S. K., Sulagitti, A. and Singh, N. N. (2018). Bio-efficacy of different novel insecticides and their interaction between numbers of sprays against diamondback moth (*Plutella xylostella* L.) infesting cabbage. *Journal of Entomological Research*, **42**(1): 51-56.
21. Sawant, C. G. and Patil, C. S. (2018). Bio-efficacy of newer insecticides against diamondback moth (*Plutella xylostella* Linn.) in cabbage at farmer's field. *International Journal of Current Microbiology and Applied Sciences*, **7**(7): 2986- 2998.
22. Sharma, P., Sharma, A., Jat, S. M., Kumawat, K. C. and prasad, B. (2022). Bioefficacy of different insecticides against diamond back moth. *The Pharma Innovation Journal*, **11**(1): 1380- 1382.
23. Venugopal, U., Kumar, A. S. H. W. A. N. I., Kota, S. A. T. H. I. S. H. and Ramya, V. (2017). Efficacy of certain insecticides against diamondback moth (*Plutella*
 - a. L.) on cabbage (*Brassica oleracea* var. capitata L.). *Agriculture Update, xylostella*
 - b. **12**(6): 1612-1616.
24. Yadav, S., Raju, S., Yadav, M., Srivastava, A. and Dwivedi, P. (2017). Bio-efficacy of new insecticidal molecules against the Diamondback moth (*Plutella xylostella* L.) on cauliflower. *Journal of Experimental Zoology*, **20**(1): 465-469.
25. Tayde AR, Khan SA. Comparative Efficacy and Economics of Chemicals Insecticides and Bioagents against Diamondback Moth (*Plutella xylostella*) on Cabbage (*Brassica oleracea* var. Capitata L.). *International Journal of Plant & Soil Science*. 2022 Sep 8:1485-9.

