

Evaluate the efficacy of different Biopesticides with Imidacloprid against maize spotted stem borer (*Chilo partellus swinhoe*) on maize under field conditions

ABSTRACT

An experiment of Maize stem borer *Chilo partellus* using different Bio-pesticides in field condition was carried out during Kharif 2022-2023 at central Research field SHUATS Prayagraj, UP., India. The management of Maize stem borer was done using 8 different treatments and benefit cost ratios of all the treatments were calculated. Total two sprays were applied to protect the crop from *Chilo partellus* using randomized block design with three replications. The observations of *Chilo partellus* 24 hours before (Pre-treatment) and 3rd, 7th and 14th day after spraying (post-treatment) were recorded for computing the percent of larval population reduction. The data were subjected to statistical analysis after appropriate transformation for interpretation. The treatment with recommended insecticide cost benefit ratio and yield is in Imidacloprid 200 SL (T₇) (1:5.01) (40.49q/ha) followed by *Beauveria bassiana* (2CFU×10⁸ ml) (T₁) (1:4.53) (38.45q/ha), *Bacillus thuringiensis* (10⁸ CFU/ml) (T₄) (1:4.41) (37.7q/ha), *Metarhizium anisopliae* (T₂) (1:4.20) (36.79q/ha), NPV (T₃) (1:4.03) (35.98q/ha), NSKE @ 5% (T₅) (1:3.97) (34.6q/ha), Azadirachtin 0.03% (T₆) (1:3.85) (33.57q/ha) Least monetary return and yield was obtained with control (T₀) (1:3.52) (28.4q/ha).

KeyWords: Biopesticides, *Chilopartellus*, Management , Maize stem borer , Imidacloprid 200SL

INTRODUCTION

Maize (*Zea mays* L.) has become a staple food in many parts of the world, with total production surpassing that of wheat or rice. It is an important staple food crop in Asia and Africa. Maize occupies a pride place among cereal crops in India and due to its high yield potential is called the queen of cereals (**Kumar and Alam 2017**).

Maize grain has elevated nutritive value as it contains about 72% starch, 10% protein, 4.8% oil, 5.8% fibre and 3% sugar. At present, out of the total maize produced, 55% is used for food purpose, about 14% for livestock, 18 % for poultry feed, 12 % for starch and one percent as seed (**Krishna and Kumar, 2018**). It is cultivated in nearly 201 m/ha with a production of 1162 m tonnes and productivity of 5754.7 kg/ha all over the world, having wider diversity of soil, climate, biodiversity and management practices (**FAOSTAT 2020**). India produced 31.51 million tonnes in an area of 9.9 million hectares in 2020-21, whereas in *kharif* 2021-22, maize production was 21.24 million tonnes (1st advance estimates) in an area of 8.15 million hectares.

Out of 66 insects reported in maize field, there are 14 major pests such as Maize stem borer *Chilo partellus* Swinhoe), Fall Armyworm (*Spodoptera frugiperda* Smith), White grub etc. among which Maize stem borer *Chilo partellus* is more complex now a days. (**Pokharel et. al.2021**).

Chilo partellus was first mentioned by **Charles Swinhoe in 1885 (CABI 2019)**. *Chilo partellus* is cosmopolitan in nature having its origin in Asia and its severity was also reported in the African region. (**Pokharel et. al.2021**). Stem borer can cause severe damage at different stages in the development of cereal crops from seedling to maturity. When infestation is severe, there is a physiological disruption of plant growth hence tassel emergence and grain formation are severely affected (**Addo-Bediako and Thanguane,2012**). *Chilo partellus*, popularly known as stalk borer that occurs during monsoon season is a major pest throughout the country. *Chilo partellus* lays eggs 10-25 days after germination on the lower side of the leaves. The larva of the *Chilo partellus* enters the whorl and causes damage to the leaves. The losses ranging 26.7 to 80.4 per cent have been reported due to *Chilo partellus* Swinhoe.

Out of them, *Chilo Partellus* (Swinhoe) is a serious pest of maize throughout India during *kharif* season causing grain yield loss of 24.3 to 36.3 percent. Almost 75% damage of the crop occurs due to attack of maize stem borer (**Dinesh et al., 2018**).

The applications of various insecticides with different modes of action strengthen insecticide resistance management strategy. Thus, to demonstrate these promising tools of pest management at farmers' fields and economic comparison of different insecticidal treatments is

necessary (Anonymous 2016).

MATERIALS AND METHODS

The present investigation was conducted at the Central Research Farm, SHUATS, Prayagraj during *kharif season 2022-23*. The experiment was conducted during the *kharif season 2022* at SHUATS, Central research farm, Prayagraj, is situated at 25.27°C north latitude 80.50°C East longitude. The research trails was laid out during the *Kharif season of 2022* in Randomized Block Design (RBD) with three replications, seven treatments and untreated control.

In the experiment maize variety, Aarhoe was sown, The site selected was uniform, cultivable with typical sandy loam soil having good drainage. Each block were sub-divided into 2m×1m with maintaining 30cm borders as bund and the treatment were assigned randomly. The field trail was conducted with seven insecticides + biopesticides treatments in which include (T₁) *Beauveria bassiana* 10⁸ CFU/ml@1g/lit, (T₂)*Metarhizium anisopliae*@2.5ml/lit, (T₃)Nuclear polyhedrosis virus @400-500ppm, (T₄) *Bacillus thuringiensis* 10⁸CFU/m@750ml/ha,(T₅)Neem seed kernel extract 5% @5 ml/lit, (T₆) Azadirachtin 0.03% @5.0ml/l,(T₇) Imidachloropid 200SL 0.5ml/lit, (T₀) Control.

Application of treatments for the management of the *Chilo partellus* was initiated as soon as 5% ETL of Larval population observed in experimental field. Subsequent application was under taken at an interval of 3, 7, 14 days were made during experimental period. The observation was recorded on weekly intervals throughout the cropping season. To assess the incidence of stem borer at weekly intervals the total number of plants and number of plants Larval population observed was counted from each plot.

The Larval population observed of the maize stem borer was calculated according to the following equation:

$$\text{Larval population} = \frac{\text{No. of larvae}}{\text{Total no. of plants}}$$

Results and Discussion

The present study entitled “Efficacy of certain chemicals and biopesticides against maize stem borer, *Chilo partellus* (Swinhoe)” was undertaken the data so obtained through observation on various aspects were subjected to statistical analysis wherever necessary and the compiled mean data are present in the following pages.

The data on the effect of various treatments on the reduction of larval population of Maize spotted Stem borer *Chilo partellus* (Swinhoe) after 1st and 2nd spray overall mean revealed that All the insecticides were significantly superior over control in reducing the larval population of stem borer which were recorded after insecticidal application Imidacloprid 200SL was found significantly superior (3.37) Over other treatments (**Haq et al., 2018**), followed by *Beauveria bassiana* (3.75), *Bacillus thuringiensis* (4.22) (**Hegde et al., 2017**), *Metarhizium anisopliae* (4.71) (**Adhikari et al. (2021)**), *Nuclear polyhedrosis virus* (5.17) (**Tayde A.R and Ramesh,M 2022**). Neem seed kernel extract (5.60), Azadirachtin (6.06) (**Rani et al., 2018**) as compared to control (8.96).

The data showed that the highest grain yield Imidacloprid (40.49q/ha) these findings are similar with **Neupane et al.(2016)**with the result of (3.38t/ha), followed by the *Beauveria bassiana* (38.45q/ha) and these findings are similar with **Adhikari (et al.,2021)** with the result of (42.50q/ha), The next best treatment is *Bacillus thuringiensis* (37.70q/ha) these findings are supported by **Dhaliwal et al.,(2018)**with the result of (34.86q/ha), the next best treatment is *Metarhizium anisopliae* (36.79q/ha) are similar with **Hegde et al.,(2017)** with the result of (35.46q/ha), the next best treatment is *Nuclear polyhedrosis virus* (35.98 q/ha). Neem seed kernel extract (34.60 q/ha) and these findings are very close to **Dinesh et al.,(2018)**with result of (32.70q/ha). The next best treatment is Azadirachtin (33.75 q/ha) these findings are similar to **Dinesh et al. (2018)** with the result of (33.70 q/ha) as compared to control (28.40 q/ha).

When the cost benefit ratio is worked out, interesting result was achieved. Among the treatments studied, the best and economical treatment was Imidacloprid (1:5.01) and was found significantly superior over rest of the treatments and these findings are similar to (**Reddy and kumar,2021**) with the result of (1:4.89). Which was followed by *Beauveria bassiana* with (1:4.53) these findings are similar to **Babu and kumar(2022)**with the result of (1:3.31), the next best treatment is *Bacillus thuringiensis* (1:4.41) these findings are similar to **Hegde et al 2017**) with the result (1:3:5), *Metarhizium anisopliae* (1:4.20) these finding similar to **Divya and mariappan(2019)** with the result (1:3.8), *Nuclear polyhedrosis virus* with (1:4.03), these findings similar to **Ramesh et al. (2022)** with the result (1:3.60) the next best treatment is Neem seed kernel extract (1:3.97) these findings are close with **Babu and Kumar (2022)** with the

result of (1:2.39) and Azadirachtin 0.03% with (1:3.85) these findings are supported by **Dinesh *et al.* (2018)** with the result of (1:1.33) and untreated control (1:3.52).

Table 1: Efficacy of different biopesticides with imidacloprid against maize spotted stem borer (*Chilo partellus swinhoe*) (1st and 2nd spray):

S. No.	Treatments	Dosage	Larval Population of <i>chilo partellus</i>									Yield(q/ha)	C:B Ratio	
			First spray					Second spray						Overall mean
			1DBS	3DAS	7DAS	14DAS	Mean	3DAS	7DAS	14DAS	Mean			
T ₁	<i>Beauveria bassiana</i> (10 ⁸ CFU/ml)	1g/lit	7.200	5.000 ^e	4.000 ^e	4.600 ^{ef}	4.533 ^{fg}	3.533 ^e	3.133 ^e	2.267 ^g	2.978 ^{fg}	3.756 ^{de}	38.45	1:4.53
T ₂	<i>Metarhizium anisopliae</i>	2.5ml/lit	6.133	5.667 ^d	5.267 ^c	5.467 ^{cd}	5.467 ^{de}	4.533 ^d	4.133 ^d	3.200 ^e	3.955 ^{de}	4.711 ^{bcd} _e	36.79	1:4.20
T ₃	<i>Nuclear polyhedrosis virus</i>	400-500ppm	6.667	6.133 ^c	5.600 ^c	5.867 ^c	5.867 ^{cd}	5.133 ^c	4.600 ^d	3.733 ^d	4.489 ^{cd}	5.178 ^{bed}	35.98	1:4.03
T ₄	<i>Bacillus thuringiensis</i> (10 ⁸ CFU/m)	750 ml/ha	6.400	5.400 ^d	4.533 ^d	5.000 ^d	4.978 ^{ef}	4.133 ^d	3.600 ^e	2.667 ^f	3.467 ^{ef}	4.223 ^{cde}	37.70	1:4.41
T ₅	Neem seed kernel extract @5%	5 ml/lit	6.867	6.533 ^b	6.133 ^b	6.400 ^b	6.355 ^{bc}	5.533 ^c	4.867 ^c	4.133 ^c	4.844 ^c	5.600 ^{bc}	34.60	1:3.97
T ₆	Azadirachtin (0.03%)	5 ml/lit	7.467	6.800 ^b	6.400 ^b	6.533 ^b	6.578 ^b	6.400 ^b	5.467 ^b	4.800 ^b	5.556 ^b	6.067 ^b	33.57	1:3.85
T ₇	Imidachloropid 200SL	0.5 ml/lit	6.200	4.667 ^e	3.800 ^e	4.400 ^f	4.289 ^g	3.133 ^e	2.467 ^f	1.800 ^h	2.467 ^g	3.378 ^e	40.49	1:5.01
T ₈	Control	-	7.667	7.800 ^a	8.767 ^a	8.900 ^a	8.489 ^a	9.200 ^a	9.400 ^a	9.733 ^a	9.444 ^a	8.967 ^a	28.40	1:3.52
F-Test			NS	S	S	S	S	S	S	S	S	S		
S.Ed.(±)			NS	0.16	0.23	0.22	0.26	0.23	0.24	0.17	0.29	0.91		
CD (0.05)			NS	0.33	0.48	0.48	0.583	0.49	0.504	0.360	0.623	1.87		-

CONCLUSION

From the present study, the results showed that Imidacloprid 200 SL (T7), is the most effective treatment against maize stem borer and produces maximum yield, and recorded the highest Cost-Benefit ratio compared to other treatments. From the critical analysis of the present findings it can be concluded that maize stem borer increases with maximum temperature and decreased with decline in minimum temperature. Insecticides like *Beauveria bassiana* 1×10^8 CFU/ml (T1) and *Bacillus thuringiensis* (T4), was found significantly superior than other treatments. Followed next effective treatment found was *Metarhizium anisopilae* (T2), *Npv* (T3). However, NSKE @5% (T5) and *Azadirachtin* 0.03% (T6) found to be least effective in managing maize stem borer (*Chilo patellus*) as an effective tool under chemical control. Hence, it is suggested that effective insecticides may be alternated along with biopesticides with the existing Integrated pest management programs to avoid the problems associated with insecticidal resistance, pest resurgence etc.

Acknowledgements

The authors are grateful to Prof. (Dr.) Rajendra B. Lal, Hon'ble Vice Chancellor, SHUATS, Prof. (Dr.) Shailesh Marker, Director of Research, Prof. (Dr.) Biswaroop Mehra, Dean, Naini Agriculture Institute and Dr. Ashwani Kumar, Associate Prof. and Head, Department of Entomology, SHUATS for taking their keen interest and encouragement to carry out this research work.

REFERENCES

- Anonymous. 2016.** <http://www.indiastat.org/agriculture/2/foodgrains/17180/maize>
- Adhikari, B., Sapkota, R., Thapa, R. B., Bhandari, G. and Dahal, K. C. (2021)** Biorational Management of Maize Stem Borer *Chilo partellus* (Swinhoe) *Azarian journal of agriculture*.**9(8):** 38-44.
- AAddo-Bediako, and NThanguane (2012)** Stem borer distribution in different sorghum cultivars as influenced by soil fertility *Agricultural Science Research Journal* **2(4)**189-194.
- Babu, J. S. and Kumar, A. (2022).** Efficacy of certain chemicals and biopesticides against maize stem borer, *Chilo partellus* (Swinhoe). *The Pharma Innovation Journal*. **11(7):** 4156-4159.
- Dinesh, M. V., Simon, S. and Nagar, S. (2018).** Biorational management of stem borer, *Chilo partellus* in maize. *Journal of Pharmacognosy and Phytochemistry*.**7(4):** 1142-1145.
- Divya, S., and Mariappan, V. N. (2019).** Integrated approach for the management of major insect pest in maize agro-ecosystem in Perambalur district.
- Dhaliwal, A. K., Brar, D. S. and Jindal, J. (2018).** Evaluation of new insecticides against maize stem borer, *Chilo partellus* (Swinhoe). *Indian Journal of Entomology*, **80(3):** 975-978.
- Hegde, K., Manjunatha, M., Sharanabasappa, K., Kalleshwaraswamy, C.M. and Adaha, S.K. (2017)** Effect of application of biopesticides and insecticides on stem borers and yield of maize (*Zea mays* L.) *International journal of pure applied biosciences*. **5(1):**42-47.
- Haq, I., Sattar, S., Ahmed, B., Zeb, Q. and Usman, A. (2018).** Compatibility of chemical and biological control for the management of maize stem borer, *Chilo Partellus*, (Swinhoe) (Lepidoptera; Pyralidae). *Sarhad Journal of Agriculture*. **34(4):** 896-903.
- Krishna, D.V. and Kumar, A (2018)** Comparative efficacy of certain chemicals against Maize stem borer (*Chilo partellus*) in Allahabad region. *International journal of scientific and Engineering research* .**9(8):**696-700.
- Kumar, A and Kumar, A (2017)** Field efficacy of seven insecticides against *Chilo partellus* (Swinhoe) on maize (*Zea mays* L.) in Allahabad. *Journal of Pharmacognosy and Phytochemistry*. **6(4):** 1591-1593.
- Prakash, V., Singh, D.V., Singh, R., Singh, G. and Kumar, S. (2017)** Efficacy of some novel insecticide against maize stem borer, *Chilo partellus* (Swinhoe) in maize. *Journal of Pharmacognosy and Phytochemistry* .SP1: 481-484
- Rani, D.S., SatyaSri, C, N. Kumar, K.A and Venkatesh, M.N (2018)** Economic evaluation and efficacy of various insecticides against maize stem borers. *Journal of Pharmacognosy and*

Phytochemistry. 7(3): 15-20.

Ramesh, K., Kumar, A. and Kalita, H. (2012). Full paper| Indian Journal of Plant Protection Vol. 40. No. 3, 2012 (208-213) Biorational Management of Stem borer, *Chilo partellus* in Maize. *Indian Journal of Plant Protection*, 40(3), 208-213.

Reddy, G. V and Kumar, A. (2021). Efficacy of various insecticides against maize stem borer *Chilo partellus* (Swinhoe) and their cost-benefit analysis at Prayagraj. *Journal of Entomology and Zoology Studies* 2021. 9(6): 130-134.

Renuka, S., Ramanujam, B. and Poornesha, B. (2015). Screening of *Beauveria bassiana* (Balsamo) Vuillemin isolates against maize Stem Borer, *Chilo partellus* (Lepidoptera: Pyralidae) and the effect of solid substrates on conidial production and virulence. *Journal of Pure Applied Microbiology*. 9(4):2979-2986.

Malav, S.K., Lyall, H., Sharma, A.K., Sharma, S., Mehta, V. and Malav, H. (2018) Efficacy of certain chemical insecticides and neem oil against stem borer (*Chilo partellus* Swin) on maize (*Zea mays* L.) *International Journal of Advanced Scientific Research and Management, Special Issue (I)*.

Neupane, S., Bhandari, G., Sharma, S. D., Yadav, S. and Subedi, S. (2016). Management of stem borer (*Chilo partellus* Swinhoe) in maize using conventional pesticides in Chitwan, Nepal. *Journal of Maize Research and Development*, 2(1): 13-19.

Parajuli, D., Ojha, B., Panthee, A. A., Pokharel, A. and Gautam, B. (2021). Management practices of maize stem borer (*Chilo partellus* Swinhoe) in Nepal. *Acta Entomology and Zoology*. 2(2): 13-22.

Saranya, V.S.L and Samiayyan, K. (2017). Efficacy of different botanicals as ovicidal molecules and oviposition deterrents against maize stem borer, *Chilo partellus* (Swinhoe). *Annals of Plant Protection Sciences*, 25(1), 12-15.

Tayde A.R and Ramesh, M (2022). Comparative efficacy of selected chemicals and biopesticides against maize stem borer (*Chilo partellus*) on Maize. *International Journal Of Plant And Soil Science*. 15(6):466-467.

