

# Effect of fodder cutting management on quality of different varieties of oat (*Avena sativa* L.)

## ABSTRACT

A field experiment was conducted to effect of cutting and splitting of nitrogen management on quality of different varieties of fodder oat during *Rabi* 2019-2020. The experiment was carried out with three treatments consisting of three varieties (RO-19, JHO-851, UPO-212) and four fodder cutting and splitting of nitrogen *viz.* (T<sub>1</sub>) Two cuttings (55 DAS & 50% flowering) + 60% N as basal + 40% N at I<sup>st</sup> cut; (T<sub>2</sub>) Two cuttings (55 DAS & 50% flowering) + 50% N as basal + 50% N at I<sup>st</sup> cut; (T<sub>3</sub>) Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% N as basal + 25% N at I<sup>st</sup> cut + 25% N at II<sup>nd</sup> cut and (T<sub>4</sub>) Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% N as basal + 30% N at I<sup>st</sup> cut + 30% N at II<sup>nd</sup> cut were laid out in split plot design with three replications. As per the ~~results~~ results, the values of RO-19 obtained the highest Crude fibre (%), Digestible crude protein (%), Total Digestible Nutrient (%), and Watersoluble carbohydrates (%) as compared to JHO-851 and UPO-212. As regarded ~~to~~ cutting and splitting of nitrogen management on Crude fibre (%), Digestible crude protein (%), Total Digestible Nutrient (%) and Watersoluble carbohydrates (%) were found maximum under Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% N as basal + 25% N at I<sup>st</sup> cut + 25% N at II<sup>nd</sup> cut.

*Keyword: Fodder cutting, Crude fibre, Digestible crude protein, Total Digestible Nutrient, Watersoluble carbohydrates.*

## INTRODUCTION

Oat (*Avena sativa* L.) is a member of the Poaceae family. The genus *Avena* has diploid, tetraploid, and hexaploid species, all of which have an  $x=7$  chromosomal number. It is the common name for oat. It is a major cereal fodder crop that originated in the Western Mediterranean region and is typically produced during the *Rabi* season. It is used as green fodder and grain for animal nutrition. In many regions of the world, it is a major winter is a multifunctional crop that can be used for grain, grazing, fodder, or as a rotational crop. The oat plant has a great growth habit, recovers quickly after harvesting, and produces high-quality herbage.

The amount of nitrogen applied and when it is applied have a significant impact on the effectiveness of nitrogen level observation. Nitrogen should be administered to a crop at times when it will not cause substantial losses and will provide sufficient nitrogen when needed. Because of its productivity and profitability, splitting nitrogen application is a key nutrient management approach. Growers can improve nutrient efficiency by splitting the total nitrogen application into two or more splits. Splitting nitrogen can help to reduce leaching and volatilization losses while also increasing the effectiveness of nitrogen application. Cutting management is a critical aspect in determining the growth, productivity, and quality of fodder crops.

Cutting management is a key element that influences the growth, productivity, and quality of fodder crops. For increased yields, general cutting management may be abandoned in fodder crops. Cutting has an impact on the crop's nutritional and natural resource usage. Multi-cut crops absorb

**Comment [Ma1]:** Your Introduction part a lot of problem

- 1) References/citation
- 2) You haven't clearly stated the significance of the study
- 3) Indicate the gap you have filled
- 4) At end indicate clearly your objective/s

more nutrients than single-cut crops, ~~This~~ this has a direct effect on the crop's nitrogen content, protein content, and other qualitative markers. Because cutting is such an important part of biomass synthesis, it is among the most significant determinants of green and dry forage yields (Patel *et al.* 2013). Two cuts are regularly administered at various times in an oat crop, however the first cut at 55 Days after sowing and the second cut at 50 % flowering, on the other hand, provided the best growth and output. (Sharma *et al.* 2001) Cutting management may be used in fodder crops in general to increase yields.

Comment [Ma2]: Not listed in the references

Comment [Ma3]: Are you sure this a right reference? Please revisit it.

## MATERIALS AND METHODS

The experiment was carried out during the ~~Rabi~~ the season 2019-20 at the Genetics and Plant Breeding farm, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.). The field was well leveled having good soil condition. Geographically, Ayodhya (Kumarganj) falls in a subtropical climate and is situated at 26° 47' North latitude, 82° 12' East longitudes with an altitude of 113 meters above mean sea level. The ~~S~~ soil of the experiment field was classified as silt loam in texture with alkaline reaction (pH 8.8) Low in organic carbon (0.25%) and available nitrogen (190 kg ha<sup>-1</sup>), medium in phosphorus (16.5 kg ha<sup>-1</sup>) and potash (238.1 kg ha<sup>-1</sup>). The experiment was laid out in a split plot design with three replications. The main plot consisted of three oat varieties (RO-19, JHO-19 and UPO-212), whereas the sub-plot had three nitrogen levels (Two cuttings (55 DAS & 50% flowering) + 60% N as basal + 40% N at I<sup>st</sup> cut, Two cuttings (55 DAS & 50% flowering) + 50% N as basal + 50% N at I<sup>st</sup> cut, Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% N as basal + 25% N at I<sup>st</sup> cut + 25% N at II<sup>nd</sup> cut and Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% N as basal + 30% N at I<sup>st</sup> cut + 30% N at II<sup>nd</sup> cut). The sowing was done on 18 November 2019 in opened furrows ~~at~~ 25 cm apart using the seed rate of 100 kg/ha. Quality parameters were determined at a half kilogram (500g) of fresh plant samples (stem and leaf) ~~was~~ were taken from each treatment during each cutting and dried in an oven at 80°C for 24 hours then made powder then 10 gm weight of the sample for analysis of quality parameters;

### Crude fibre (%)

To determine crude fibre (%), 1.0 g of ~~oven dried~~ oven-dried plant material was taken in 250 ml beaker, added 1.25% H<sub>2</sub>SO<sub>4</sub> and distilled water and made the volume up to 200 ml then placed ~~it~~ it on flam for 30 minutes filtered and washed. Then again added 1.25% NaOH and ~~distillated~~ distilled water and made the volume up to 200 ml. Heated again for 30 minutes and residues were washed and filtered again. The residues were put in a pre-weighted crucible and it was placed in an oven at 105°C for drying for 24 hours. After recording the dry weight (W<sub>1</sub>) the samples were placed in a muffle furnace at 600 °C till ~~gray~~ grey or white ash was obtained. Then cool it and the weight of ash (W<sub>2</sub>) was recorded and the crude fibre (%) was calculated following the formula.

Comment [Ma4]: Be consistent with British English

$$\text{Crude fibre} = \frac{(\text{Weight of crucible} + \text{dry matter}) \times \text{Weight of crucible ash}}{\text{Amount of substance taken}} \times 100$$

### Digestible Crude Protein

To find out the digestible crude protein (DCP) content of a feed, multiply the digestibility coefficient to

the crude protein ~~with~~by the crude protein content of the feedstuff.

$$\text{DCP} = \text{Digestibility Co-efficient} \times \text{CP}$$

#### Water soluble carbohydrates

Water soluble carbohydrate was determined by AOAC (1975).

#### Total Digestible Nutrient

Moore *et al.* (1953) as follows:

$$\text{TDN}(\%) = \text{Digestible crude protein}(\%) + \text{Digestible crude fibre}(\%) + \text{Digestible Nitrogen free extract}(\%) + \text{Digestible ether extract}(\%) \times 2.25$$

### RESULT AND DISCUSSION

#### Crude fibre (%)

The crude fibre (%) was non-significant at all cutting stages of the crop. The maximum crude fibre (25.92, 25.52 and 26.30%) was recorded with variety RO-19 at all cutting stages, while minimum crude fibre (24.88, 24.55 and 25.30%) was found with variety UPO-212 during an experiment.

The crude fibre as cutting and splitting of nitrogen at first cut was found not affected significantly. The maximum crude fibre (25.95%) as first cut at two cuttings (55 DAS & 50% flowering) + 60% nitrogen as basal + 40% nitrogen at I<sup>st</sup> cut whereas minimum crude fibre (25.14%) at two cuttings (55 DAS & 50% flowering) + 50% nitrogen as basal + 50% nitrogen at I<sup>st</sup> cut and at second cut crude fibre (25.89%) was recorded at two cuttings (55 DAS & 50% flowering) + 50% nitrogen as basal + 50% nitrogen at I<sup>st</sup> cut significantly higher at par with crude fibre (23.99) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30% nitrogen at I<sup>st</sup> cut + 30% nitrogen at II<sup>nd</sup> cut and at third cut stage of crop as crude fibre (25.36%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% nitrogen as basal + 25% nitrogen at I<sup>st</sup> cut + 25% nitrogen at II<sup>nd</sup> cut, where recorded minimum crude fibre (24.74%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30% nitrogen at I<sup>st</sup> cut + 30% nitrogen at II<sup>nd</sup> cut. The result might be found due to soil moisture ~~had~~ having a direct relationship with crude fibre content. Bhilare and Joshi, (2007), also obtained the similar results.

#### Digestible Crude protein (%)

The digestible Crude protein (%) was observed as significantly not affected by varieties at all cut stages of crop and cutting and splitting of nitrogen at 35 Days after the first cut (II<sup>nd</sup> cut) and 50% flowering (III<sup>rd</sup> cut) affected significantly except 55 DAS are represented in Table- 1. The maximum digestible Crude protein (7.35 and 7.14 and 6.68%) was recorded by variety RO-19 at all cut stages whereas minimum digestible Crude protein (7.08, 6.87 and 6.61%) by variety UPO-212 at all cut stages.

The digestible crude protein was observed as cutting and spitting of nitrogen at the first cut was not affected significantly. The maximum digestible crude protein (7.27%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30% nitrogen at 1<sup>st</sup> cut + 30% nitrogen at 2<sup>nd</sup> cut whereas minimum digestible crude protein (7.16%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% nitrogen as basal + 25% nitrogen at 1<sup>st</sup> cut + 25%

nitrogen at 2<sup>nd</sup> cut and at second cut was found digestible crude protein (7.06%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 40 % nitrogen as basal + 30 % nitrogen at 1<sup>st</sup> cut + 30% nitrogen at 2<sup>nd</sup> cut significantly higher at par with digestible crude protein (6.94%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 50 % nitrogen as basal + 25% nitrogen at 1<sup>st</sup> cut + 25% nitrogen at 2<sup>nd</sup> cut while in third cut digestible crude protein (7.28%) was recorded maximum at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 50 % nitrogen as basal + 25 % nitrogen at 1<sup>st</sup> cut + 25 % nitrogen at 2<sup>nd</sup> cut over of the three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30% nitrogen at 1<sup>st</sup> cut + 30% nitrogen at 2<sup>nd</sup> cut.

### Total Digestible Nutrient (%)

The data observed was on total digestible nutrients (%) as significantly was not affected by varieties and cutting and splitting of nitrogen at 35 Days after the first cut (I<sup>nd</sup> cut) and 50% flowering (III<sup>nd</sup> cut) affected significantly except 55 DAS are presented in Table- 1.

The data presented maximum total digestible nutrient (73.64, 71.43 and 73.56%) by variety RO-19 whereas minimum total digestible nutrient (70.84, 68.71 and 73.13%) by variety UPO-212 at all cut stages of crop. The total digestible nutrient was observed as cutting and splitting of nitrogen at first cut not affected significantly. The maximum total digestible nutrient (72.80%) was found at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30 % nitrogen at I<sup>st</sup> cut + 30 % nitrogen at II<sup>nd</sup> cut whereas recorded minimum total digestible nutrient (71.68%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 50% nitrogen as basal + 25% nitrogen at I<sup>st</sup> cut + 25% nitrogen at II<sup>nd</sup> cut at the first cut stage and at the second cut stage was found that the highest total digestible nutrient (72.62%) significantly at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 40% nitrogen as basal + 30% nitrogen at I<sup>st</sup> cut + 30% nitrogen at II<sup>nd</sup> cut at par with total digestible nutrient (70.53%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 50 % nitrogen as basal + 25 % nitrogen at I<sup>st</sup> cut + 25% nitrogen at II<sup>nd</sup> cut and in third cut under was found total digestible nutrient (74.26%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 40 % nitrogen as basal + 30 % nitrogen at I<sup>st</sup> cut + 30 % nitrogen at II<sup>nd</sup> cut at par with total digestible nutrient (73.11%) at Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 50% nitrogen as basal + 25% nitrogen at I<sup>st</sup> cut + 25% nitrogen at II<sup>nd</sup> cut. The result Similar finding was reported by Bhilare and Joshi, Joshi (2007).

### Water soluble carbohydrates (%)

The water-soluble carbohydrates (%) was-were recorded at various cut stages of the crop *i.e.* 55 DAS to at each cut as influenced by different cutting and splitting of nitrogen by different varieties of oat is summarized in Table-1.

The data present water-soluble carbohydrates on different varieties not affected significantly. The water soluble carbohydrates was-were recorded as maximum (16.55, 16.07 and 12.44%) by variety RO-19 at all cut stages where recorded minimum (15.94, 15.48 and 12.14%) by variety UPO-212 at all cut stages. The water-soluble carbohydrates as cutting and splitting of nitrogen at the first cut were not affected significantly. The maximum water-soluble carbohydrates (16.38%) as first cut at

three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30 % nitrogen at I<sup>st</sup> cut + 30 % nitrogen at II<sup>nd</sup> cut whereas minimum crude fibre (16.10%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 50 % nitrogen as basal + 25 % nitrogen at I<sup>st</sup> cut + 25 % nitrogen at II<sup>nd</sup> cut and at second water soluble carbohydrates (15.90%) was recorded at Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 40 % nitrogen as basal + 30 % nitrogen at I<sup>st</sup> cut + 30 % nitrogen at II<sup>nd</sup> cut significantly higher at par with water soluble carbohydrates (15.66%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% nitrogen as basal + 25% nitrogen at I<sup>st</sup> cut + 25% nitrogen at II<sup>nd</sup> cut and at third cut stage of crop as water soluble carbohydrates (12.56%) found maximum at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50 % flowering) + 40 % nitrogen as basal + 30 % nitrogen at I<sup>st</sup> cut + 30 % nitrogen at II<sup>nd</sup> cut, where recorded minimum water soluble carbohydrates (12.31%) at three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 40% nitrogen as basal + 30% nitrogen at I<sup>st</sup> cut + 30% nitrogen at II<sup>nd</sup> cut. ~~The R~~ result might be found due to soil moisture ~~had~~ having a direct relationship with crude fibre content. Bhilare and Joshi, (2007), also obtained the similar results.

## CONCLUSION

Conclusively as per the results, the values of RO-19 obtained the highest Crude fibre (%), Digestible crude protein (%), Total Digestible Nutrient (%) and Water soluble carbohydrates (%) significantly. ~~cutting~~ Cutting and splitting of nitrogen management, Crude fibre (%), Digestible crude protein (%), Total Digestible Nutrient (%) and Water soluble carbohydrates (%) were found maximum under Three cuttings (55 DAS, 35 days after 1<sup>st</sup> cut & 50% flowering) + 50% N as basal + 25% N at I<sup>st</sup> cut + 25% N at II<sup>nd</sup> cut.

**Table-1: Crude fibre(%) and digestible crude protein(%) at subsequent cuttings influenced by the splitting of nitrogen on different varieties of oat. (*Avena sativa* L.).**

Treatments	Crude fibre(%)			Digestible crude protein(%)		
	(I <sup>st</sup> cut)	(II <sup>nd</sup> cut)	(III <sup>rd</sup> cut)	(I <sup>st</sup> cut)	(II <sup>nd</sup> cut)	(III <sup>rd</sup> cut)
<b>Varieties</b>						
RO-19	25.92	25.52	26.30	7.35	7.14	6.68
JHO-851	25.84	25.04	25.84	7.22	7.01	6.61
UPO-212	24.88	24.55	25.30	7.08	6.87	6.00
SEm±	0.53	0.50	0.44	0.12	0.17	0.18
CD at 5%	NS	NS	NS	NS	NS	NS
<b>Cutting and splitting of nitrogen management</b>						
Two cuttings (55 DAS & 50% flowering) + 60% Nasbasal + 40% N at I <sup>st</sup> cut	24.95	25.69	0.00	7.20	6.32	0.00
Two cuttings (55 DAS & 50% flowering) + 50% Nasbasal + 50% N at I <sup>st</sup> cut	25.14	25.89	0.00	7.25	6.05	0.00
Three cuttings (55 DAS, 35 days after 1 <sup>st</sup> cut & 50% flowering) + 50% N asbasal + 25% Nat I <sup>st</sup> cut + 25% Nat II <sup>nd</sup> cut	24.85	24.59	25.36	7.16	6.94	7.28
Three cuttings (55 DAS, 35 days after 1 <sup>st</sup> cut & 50% flowering) + 40% N asbasal + 30% Nat I <sup>st</sup> cut + 30% Nat II <sup>nd</sup> cut	25.24	23.99	24.74	7.27	7.06	6.43
SEm±	0.61	0.58	0.26	0.43	0.19	0.10
CD at 5%	NS	1.72	0.78	NS	0.57	0.30

**Table-2: Total Digestible Nutrient(%) and Watersoluble carbohydrates(%) at subsequent cuttings influenced by the splitting of nitrogen on different varieties of oat. (*Avena sativa* L.).**

Treatments	Total Digestible Nutrient(%)			Watersoluble carbohydrates(%)		
	(I <sup>st</sup> cut)	(II <sup>nd</sup> cut)	(III <sup>rd</sup> cut)	(I <sup>st</sup> cut)	(II <sup>nd</sup> cut)	(III <sup>rd</sup> cut)
<b>Varieties</b>						
RO-19	73.64	71.43	74.84	16.55	16.07	12.44
JHO-851	72.24	70.07	73.56	16.25	15.77	12.29
UPO-212	70.84	68.71	73.13	15.94	15.48	12.14
SEm±	1.28	1.42	1.34	0.28	0.28	0.21
CD at 5%	NS	NS	NS	NS	NS	NS
<b>Cutting and splitting of nitrogen management</b>						
Two cuttings (55 DAS & 50% flowering)+60% Nasbasal+40% N at I <sup>st</sup> cut	71.96	67.89	0.00	16.20	14.32	0.00
Two cuttings (55 DAS & 50% flowering)+50% Nasbasal+50% N at I <sup>st</sup> cut	72.52	68.13	0.00	16.30	14.86	0.00
Three cuttings (55 DAS, 35 days after I <sup>st</sup> cut & 50% flowering) + 50% N asbasal+25% Nat I <sup>st</sup> cut+25% Nat II <sup>nd</sup> cut	71.68	70.53	73.11	16.10	15.66	12.31
Three cuttings (55 DAS, 35 days after I <sup>st</sup> cut & 50% flowering) + 40% N asbasal+30% Nat I <sup>st</sup> cut+30% Nat II <sup>nd</sup> cut	72.80	72.62	74.26	16.38	15.90	12.56
SEm±	1.48	1.64	0.78	0.33	0.32	0.18
CD at 5%	NS	4.39	2.34	NS	0.94	0.54

## REFERENCE

- Bhilare, R. L. and Joshi, Y. P. (2007). Productivity and quality of oat (*Avena sativa* L.) in relation to cutting management and nitrogen levels. *Indian Journal of Agronomy* **52**:247-250.
- Dabhi, M.S., Patel, M.R., Chaudhari, C. R., Patel, V. N. and Patel, P.M. (2017) Response of oat (*Avena sativa* L.) varieties to methods of sowing and nitrogen levels on forage yield and quality. *International Journal of Chemical Studies*. **5**(4):683-686.
- Godara, A. S., Satpal, Neelam, Y., Jindal and Phogat, D. S. (2019). Performance of dual-purpose oat, wheat and barley under different cutting management system. *Forage Res.*, **44** (4):pp. 286-290.
- Kaur, G. and Goyal, M. (2017). Effect of growth stages and fertility levels on growth, yield and quality of fodder oats (*Avena sativa* L.). *Journal of Applied and Natural Science* **9**(3):1287 -1296.
- Kumar, S.G.V., Sukanya, T.S., Jayaramaiah, R., Somashekar, K.S. and Morab, P.N. (2020). Green fodder yield and quality of fodder oats (*Avena sativa* L.) as influenced by dates of sowing and nitrogen levels in Southern transition zone of Karnataka. *Journal of Pharmacognosy and Phytochemistry*. **10**(1):1475-1479.
- Midha, L. K., Duhan, B. S. and Arya, S. (2015). Performance of promising entries of oat (*Avena sativa* L.) under different nitrogen levels. *Forage Res.*, **41** (2):pp. 122-125.
- Sharma, S.K.; Bhunia, S.R. (2001). Response of oat (*Avena sativa* L.) to cutting management, method of sowing and nitrogen. *Indian Journal of Agronomy*. **46**(3): 563-567.
- Shikha and S Singh. (2018) Varietal evaluation of oats (*Avena sativa* L.) varieties under different nutrient management. *International Journal of Chemical Studies* **6**(3):1268-1271.

**Comment [Ma5]:** From where you have listed here? Almost all of the references were not cited in the document.....

**Comment [Ma6]:** Please check in the document....