

Response of Silkworm (*Bombyx mori* L.) Breeds to Temperature and *BmNPV* Stress

ABSTRACT

Silkworm *Bombyx mori* L. is one of the most important domesticated insects, which produces luxuriant silk thread in the form of cocoon by consuming mulberry leaves during larval period. The growth and development of silkworm is greatly influenced by environmental conditions. The seasonal differences in the environmental components considerably affect the genotypic expression in the form of phenotypic output of silkworm crop. *Bombyx mori* nucleopolyhedrovirus (*BmNPV*) is a pathogen that causes great economic losses in sericulture. In the present investigation, twelve bivoltine silkworm breeds (WM, ND₅, NB₄D₂, U-4, PO₁, ND₃, U-6, CSR₂, SH₆, SPO, U-3 and NSP) were studied for temperature and *BmNPV* stress. After third and fourth moult silkworm larvae were treated with *BmNPV* inoculum at 25±1^o C and 30±1^o C temperature respectively. Increased mortality was observed at 30±1^o C when compared to 25±1^o C. However, with respect to breeds the highest total mortality was recorded in breed PO₁ and least in breed U-3. Silkworm breed U-3 may have the ability to tolerate *BmNPV* incidence and can be exploit for evolving disease-tolerant silkworm hybrids.

Keywords: *BmNPV*, Silkworm, Breeds, Larval, Mortality

1. INTRODUCTION

Silkworm, (*Bombyx mori* L.) is sensitive to environmental, nutritional and microbial factors which results in outbreak of various diseases leading to silkworm mortality and cocoon crop loss. The most prevalent and serious diseases in the silkworm are grasserie, flacherie, muscardine and pebrine caused by virus, bacteria, fungi and microsporidia, respectively Jiang and Xia [1]. Among the viral diseases, nuclear polyhedrosis is most severe and contagious. NPV can affect *B. mori* in different stages of its life cycle but the infected silkworm expresses disease symptoms during the final stage of larval growth and dies without producing a cocoon resulting in the wastage of time and labour for the farmers thereby causing significant financial losses thus posing a serious threat to the global sericulture Brancalhao [2]. In India, more than 50 per cent of silk cocoon crop losses are attributed to *BmNPV* infection Khurad *et al.* [3]. The temperature fluctuations directly influence the incidence of viral diseases. The optimum conditions of temperature (25±1^oC) and humidity (75%) favours the best production and least incidence of nuclear polyhedrosis although with the increase in both factors, the incidence of nuclear polyhedrosis also increases even in uncontaminated batches Basavarajappa and Savanurmath [4].

Among many measures of silkworm disease control and prevention, the utilization of disease tolerant silkworm breeds along with the disinfection would be the most effective step in the direction of the disease prevention Sivaprasad and Chandrashekharaiyah [5]. In order to obtain high and stable cocoon yield it is necessary to reduce the incidence which in turn decrease pathogen quantity and pathogenicity and inturn strengthen the larval health by increasing their disease resistance ability Singh *et al.* [6]. The production and use of relatively tolerant silkworm breeds can be a best approach to prevent an infectious disease such as nuclear polyhedrosis. The screening of breeds for their relative tolerance for *BmNPV* would be helpful in identifying the silkworm breeds that are less susceptible and

they might be exploited commercially to evolve tolerant breeds through breeding plans for increased silk productivity.

The cocoon crop loss due to disease incidence in silkworm breeds is utmost, thus in order to minimize the loss there is a need to evolve tolerant breeds which can perform well even under adverse eco-climatic conditions to get sustainable cocoon yield. Therefore, the present experiment was undertaken to screen the available breeds for their relative tolerance to *BmNPV* and to identify and utilize the comparatively tolerant breeds for evolving certain crosses for use in future breeding programme.

2. MATERIALS AND METHOD

Twelve bivoltine silkworm breeds namely; WM, ND₅, NB₄D₂, U-4, PO₁, ND₃, U-6, CSR₂, SH₆, SPO, U-3 and NSP maintained at Division of Sericulture, SKUAST-J, Chatha, Jammu were utilized for conducting experiment. The silkworm eggs of above said breeds were surface sterilized by putting eggs in 2% formaldehyde solution for 2-3 minutes. After sterilization, the eggs were washed in running water till smell of formalin goes off and then dried the eggs in shade. The eggs were then incubated for twelve days at optimum temperature (25±1°C) and 80±5 per cent relative humidity, followed by black boxing. On Thirteen day, the silkworm larvae were hatched out and the newly hatched worms were then shifted to rearing beds and reared as per rearing method suggested by Krishnaswami [7].

2.1 Isolation and purification of polyhedra

Silkworm larvae showing symptoms of grasserie disease like intersegmental swelling and shiny body were collected from the rearing room. Then the haemolymph was collected from each diseased larvae in sterilized glass tubes by outward bending the worm in left hand and cutting the first pair of prolegs followed by gentle press. Test tubes containing the haemolymph subjected for bacterial degradation for 24 hours and centrifuged at 5000 rpm for 10 minutes for pelleting the *BmNPV* polyhedra. The polyhedra were washed thrice with distilled water and then with 1M NaCl by centrifugation at 5000 rpm for 10 minutes. The polyhedra were then suspended in distilled water and stored at 4°C in the refrigerator. The polyhedral concentration was determined using the Neubauer's haemocytometer.

2.2 Infectivity technique

Silkworm larvae of each breed were separated into two batches after third moult. Each batch having two treatments and each treatment consists of three replications with 50 larvae in each replication. In the first treatment- silkworm breeds after third moult were kept at 25±1°C temperature and then inoculated, in second treatment - silkworm breeds after third moult were kept at 30±1°C temperature and inoculated, in third treatment - silkworm breeds after fourth moult were kept at 25±1°C temperature followed by inoculation and in fourth treatment - silkworm breeds after fourth moult were kept at 30±1°C temperature and then inoculated. The silkworm larvae of fourth and fifth instar were selected because the silk gland is well developed during these instars and occurrence of disease is maximum at these stages. The silkworm larvae were starved during the temperature treatment period for six hours. The infection was carried out per orally by feeding silkworm with virus suspension smeared leaves. The leaf bits (10×12 cm size) were prepared, washed in running water and shade dried. These mulberry leaves were smeared evenly with the virus suspension at 1×10³ POB per ml by using non-absorbent cotton. After shade dried for five minutes, the leaf fed to the silkworm larvae. For subsequent feeding inoculum free leaves were used for treated and untreated batches. The observation on mortality due to *BmNPV* was recorded up to pupal stage.

3. RESULTS AND DISCUSSION

3.1 Temperature variation and *BmNPV* inoculum after third moult

Exposure of silkworm breeds to room temperature ($25\pm 1^{\circ}\text{C}$) and high temperature ($30\pm 1^{\circ}\text{C}$) for six hours followed by *BmNPV* (1×10^3 POB/ml) inoculation on first day of fourth instar recorded significant variation in total mortality. At $25\pm 1^{\circ}\text{C}$ temperature, the significant variations were observed in total mortality and the maximum total mortality was recorded in breed PO_1 (40.33 ± 1.45) followed by breeds SPO (35.67 ± 1.73) and NSP (33.67 ± 1.15). The lowest total mortality was observed in U-3 (17.33 ± 1.20) (Table 1). At $30\pm 1^{\circ}\text{C}$ temperature, the total mortality varied significantly and the highest total mortality was recorded in breed PO_1 (52.00 ± 1.00) followed by breed SPO (48.67 ± 0.66) and NSP (47.67 ± 0.66), whereas least total mortality was recorded in breed U-3 (21.00 ± 1.20) (Table 1). With respect to control batch the maximum total mortality was recorded in breed PO_1 (5.33 ± 0.33) followed by breed NSP (5.00 ± 0.57) and NB_4D_2 (4.66 ± 0.88), whereas least total mortality was recorded in breed U-3 (2.00 ± 0.57) (Table 1).

Table 1: Mortality percentage of silkworm breeds due temperature exposures and *BmNPV* inoculum (1×10^3 POB/ml) after third moult

Breeds	Total mortality		
	25±1° C	30±1° C	CONTROL
WM	25.33±1.15	36.00±0.33	4.00±0.57
ND ₅	20.67±1.45	27.33±0.88	3.66±0.88
NB ₄ D ₂	26.67±1.20	39.33±0.88	4.66±0.88
U-4	18.00±1.15	25.00±0.57	3.00±0.57
PO ₁	40.33±1.45	52.00±1.00	5.33±0.33
ND ₃	32.33±0.57	40.33±1.20	4.66±0.33
U-6	23.00±1.15	30.67±0.57	4.00±0.57
CSR ₂	33.00±0.88	39.67±0.88	4.00±0.57
SH ₆	32.00±1.45	41.00±1.20	4.33±0.33
SPO	35.67±1.73	48.67±0.66	4.00±0.57
U-3	17.33±1.20	21.00±1.20	2.00±0.57
NSP	33.67±1.15	47.67±0.66	5.00±0.57

Each value is a mean±standard error of three replications.

3.2 Temperature variation and *BmNPV* inoculum after fourth moult

Larvae of Silkworm breeds exposed to room temperature ($25\pm 1^{\circ}\text{C}$) and high temperature ($30\pm 1^{\circ}\text{C}$) for six hours followed by *BmNPV* (1×10^3 POB/ml) inoculation on first day of fifth instar recorded significant variation in total mortality. At $25\pm 1^{\circ}\text{C}$ temperature, the highest total mortality was recorded in breeds NSP (31.67 ± 1.15) that was statistically at par with PO_1 (31.67 ± 1.15). However, the lowest total mortality was found in breed U-3 (12.00 ± 0.88). At $30\pm 1^{\circ}\text{C}$ temperature, the total mortality varied significantly and the maximum larval mortality was observed in breed PO_1 (60.00 ± 0.66) followed by breed SPO (55.67 ± 1.15), while the least total mortality was observed in U-3 (27.00 ± 0.88) (Table 2). With respect to control batch the maximum total mortality was recorded in breed PO_1 (5.33 ± 0.33) followed by breed NSP (5.00 ± 0.57) and NB_4D_2 (4.66 ± 0.88), whereas least total mortality was recorded in breed U-3 (2.00 ± 0.57) (Table 2).

Table 2: Mortality percentage of silkworm breeds due temperature exposures and *BmNPV* inoculum (1×10^3 POB/ml) after fourth moult

Breeds	Total mortality		
	$25\pm 1^{\circ}\text{C}$	$30\pm 1^{\circ}\text{C}$	CONTROL
WM	20.67 ± 1.15	46.00 ± 0.88	4.00 ± 0.57
ND_5	16.67 ± 0.66	34.33 ± 0.57	3.66 ± 0.88
NB_4D_2	21.67 ± 1.45	45.33 ± 0.88	4.66 ± 0.88
U-4	15.00 ± 0.57	31.33 ± 0.57	3.00 ± 0.57
PO_1	31.67 ± 1.15	60.00 ± 0.66	5.33 ± 0.33
ND_3	26.67 ± 0.88	45.33 ± 0.57	4.66 ± 0.33
U-6	18.67 ± 1.15	36.00 ± 0.33	4.00 ± 0.57
CSR_2	27.67 ± 1.20	49.00 ± 1.15	4.00 ± 0.57
SH_6	25.33 ± 1.45	50.00 ± 0.88	4.33 ± 0.33
SPO	31.00 ± 0.57	55.67 ± 1.15	4.00 ± 0.57
U-3	12.00 ± 0.88	27.00 ± 0.88	2.00 ± 0.57
NSP	31.67 ± 1.15	49.33 ± 1.20	5.00 ± 0.57

Each value is a mean \pm standard error of three replications.

Success of silkworm breeds depends largely on their adaptability to the environment in which it is destined to be reared. It is clear from the results, under BmNPV inoculated conditions the survival rate of silkworm breeds decline to that of control. Extensive work has been done on the induction of viral diseases. Among them the important physical agents which induce diseases are temperature and pathogenic virulence. Aruga *et al.* [8] found that temperature higher or lower than 25^o C act as a stress and tends to increase the larval susceptibility to viral infections. Sowmyshree *et al.* [9] studied the influence of temperature on resistance of the breed was much higher when the host was infected by the pathogen even at low dosage. Himeno *et al.* [10] concluded that virus multiplication at 25^o C temperature in fifth instar larvae went hidden but were reactivated by cold temperature treatment and observed that the incidence of nuclear polyhedrosis virus was higher in sudden changes from room temperature to low temperature. Matsubara [11] studied the changes in the resistance to NPV by subjecting fourth and fifth instar larvae to high temperature (33, 35 and 37^o C) treatments at various periods of time immediately after the ecdysis and concluded that temperature and pathogenic virulence are the most important physical agent for induction of diseases in silkworms. Temperature acts as an important external physical factor for both larval susceptibility and multiplication of viruses in the host. Silkworms are adapted to rearing at 25^o C and temperature higher and lower than this limit increases the susceptibility of silkworm to the virus and this susceptibility decreases with the larval stages. Steinhaus [12] reported that environmental factor such as temperature activates the latent virus and once the occult virus reaches the infective stage, the virus multiplication proceeds, in the same manner as in case of natural infections. Wojda [13] reviewed that heat-shock proteins are induced by infection as well as heat shock, and also play an important role in the immune response of insects. Such stress components function as immune modulators (Palmer [14]). Lakshmi *et al.* [15] studied the stability ten hybrids under fluctuating tropical environmental conditions with BmNPV inoculation at room temperature (25^o C), low temperature (5^o C) and high temperature (32^o C). They recorded maximum mortality at 5^o C and minimum mortality at 32^o C. In this study, the larvae when exposed to 30±1^o C temperature, the least mortality was observed in U-3 breed that indicates its high level of tolerance to temperature and BmNPV at fourth and fifth instar.

4. CONCLUSION

Observations from the current study, such as total larval mortality of twelve silkworm breeds (ND₅, NB₄D₂, U-4, PO₁, ND₃, U-6, CSR₂, SH₆, SPO, U-3 and NSP) after temperature variation and inoculation with BmNPV at fourth and fifth instar showed that silkworm breed U-3 recorded the least

total larval mortality at both $25\pm 1^{\circ}$ C and $30\pm 1^{\circ}$ C. Thus, silkworm breed U-3 may have the ability to tolerate BmNPV incidence and can be exploit for evolving disease-tolerant silkworm hybrids.

REFERENCES

1. Jiang L, Xia Q. The progress and future of enhancing antiviral capacity by transgenic technology in the silkworm *Bombyx mori*. Insect biochemistry and molecular biology. 2014;48:1-7.
2. Brancalhão RM. Vírus entomopatogênicos no bicho-da-seda: taxonomia e citopatologia causada por nucleopolyhedrovirus em células de *Bombyx mori*. Biotecnologia. 2002;24:54-8.
3. Khurad AM, Kanginakudru S, Qureshi SO, Rathod MK, Rai MM, Nagaraju J. A new *Bombyx mori* larval ovarian cell line highly susceptible to nucleopolyhedrovirus. Journal of invertebrate pathology. 2006;92(2):59-65.
4. Basavarajappa S, Savanurmah CJ. Effect of different mulberry varieties during late-age silkworm *Bombyx mori* on incidence of grasserie and the cocoon characters. Uttar Pradesh Journal of Zoology. 1996:104-8.
5. Sivaprasad V and Chandrashekharaiiah. Strategies for breeding disease resistance silkworms. Mulberry Silkworm Breeders Summit, APSSRDI, Hindupur, India. 2003.
6. Singh GP, Yuyin C, Datta RK, Mengkui X. Development of resistance to *Bombyx mori* Densonucleosis virus into a susceptible silkworm breed. International Journal of Industrial Entomology. 2003;6(2):145-9.
7. Krishnaswami S. New Technology of Silkworm Rearing, Central Silk Board, Bangalore. 1978. Pp- 17-21.
8. Aruga H, Watanabe H, Nagano H. Interference by the heat-inactivated virus on the active virus of the cytoplasmic polyhedrosis in the silkworm, *Bombyx mori* L. The Journal of Sericultural Science of Japan. 1963;32(2):51-7.
9. Sowmyashree, T. S., Sowmya, P., Sudhakara Rao, P. and Nataraju, B. Determination of virulence in the selected bivoltine silkworm breeds through *BmNPV* inoculation and low temperature induction. Indian Journal of Animal Research. 2013; 47(6), pp. 486-491.
10. Himeno M, Matsubara F, Hayashiya K. The occult virus of nuclear polyhedrosis of the silkworm larvae. Journal of Invertebrate Pathology. 1973;22(2):292-5.
11. Matsubara F, Wu Yi, Mori H, Oogashira H. Changes in the resistance to the infection with NPV induced by low and high temperature treatment in the germ-free silkworm, *Bombyx mori*. The Journal of Sericultural Science of Japan. 1984;53(6):538-42.

12. Steinhaus E A. Stress as factor in insect diseases. In: Proceedings of Tenth International Congress Entomology. 1958. pp 725-730.
13. Wojda, I. Temperature stress and insect immunity. *Journal of Thermal Biology*. 2017; 68, pp. 96-103. <https://doi.org/10.1016/j.jtherbio.2016.12.002>
14. Palmer, C.V. Immunity and the coral crisis. *Communications Biology* 2018: 1, pp. 91 <https://doi.org/10.1038/s42003-018-0097-4>
15. Lakshmi LV, Sivaprasad V, Sujathamma P. Studies on seasonal performance of newly developed bivoltine silkworm (*Bombyx mori* L) hybrids tolerant to *BmNPV* and effect of temperature on disease induction. *Animal Review*. 2014;1(4):57-64.

UNDER PEER REVIEW