

Original Research Article

Response of boron and molybdenum on growth and yield of black gram (*Vignamungo* L.)

ABSTRACT

A field experiment was conducted during Zaid 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P) on the topic “Response of boron and molybdenum on growth and yield of black gram (*Vignamungo* L.)”, to study treatments consisting of three levels of Molybdenum viz. Mo 600, 700 and 800 ppm and three levels of Boron viz. 3000, 4000 and 5000 ppm. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.28 %), available N (225 kg/ha), available P (19.50 kg/ha) and available K (92 kg/ha). There were 10 treatments each being replicated thrice and laid out in Randomized Block Design. The results revealed that treatment 9 (Boron at 5000 ppm + Molybdenum at 800 ppm) recorded significantly Plant height (43.54 cm), Number of branches/plant (32.07), plant dry weight (47.20 g), number of nodules per plant (16.80) and crop growth rate (35.85 g/m² /day), number of pods per plant (29.80), number of seeds per pod (7.76), Test weight (40.20 g), Seed yield (1.39t/ha), Stover yield (2.99 t/ha), gross return (INR 69500.00), net return (INR 43209.90) and B:C ratio (1.64) compared to other treatments.

Keywords: Boron, Economics, Growth, Molybdenum, Yield.

Comment [DS1]: Name of species in italic font

Comment [DS2]: Name of country.

Comment [DS3]: Italic font

Comment [DS4]: High, higher or lower

Comment [DS5]: Check values with tables

Comment [DS6]: INR 43209.90
Check values

Comment [DS7]: B:C stands for

Comment [DS8]: Conclusion missing
Importance of B and Mo missing
Aim of study missing

1. INTRODUCTION:

Black gram (*VignaMungo* L.), is one of the important pulse crops, grown throughout the country belongs to the family Leguminosae. In India cultivated over a wide range of agro - climatic zones of the country. This crop is grown in Kharif and Rabi seasons. The crop is resistant to adverse climatic conditions and improve the soil fertility by fixing atmospheric nitrogen in the soil. It has been reported that the crop produces equivalent to 22.10 kg of N/ha., which has been estimated to be supplement of 59 thousand tonnes of urea annually [1]. The pulse 'Black gram' plays an important role in Indian diet, as it contains vegetable protein and supplement to cereal-based diet. Its seed contains moisture 10.9%, protein 20-25%, fiber 0.9%, fat 1.3%, starch 40- 47%, ash 3.40%, crude fiber 4.2%, calcium 145 (mg/100 g), iron 7.8 (mg/100g), (Anonymous, 2006) which is almost three times that of cereals and other minerals and vitamins. Apart from consumption pulses crop cultivation improves soil fertility, the physical structure of soil and also act as nutrition fodder for cattle. The pulse availability per adult per day at present is only 36 which is against the minimum requirement of 50g/adult/day. The recommended dietary allowances (RDA) for adult male and female is 60 g and 55g per day [2].

Boron is one of the most important trace elements, which is essential for normal life cycle of the plants [3]. It influences the absorption of nitrogen, phosphorus, potassium and its deficiency effects the equilibrium of these macronutrients. It increases the yield and growth of plants by increasing the leaf area expansion, 1000 seed weight, nodule formation, seed yield and biological yield. It influences the major cellular functions and metabolic activities in plants and required for cell differentiation at all growing tips of plants (meristems) where cell division is active. [4]studied the effect of boron on pulses and concluded that it is important for protein synthesis and improved protein content.

Molybdenum is an effective micronutrient that plays an important role in improving the crop yield and quality. Molybdenum is one of the most recognized nutrient elements considered to be essential for the growth of plants. As a constituent of nitrate reductase and nitrogenase enzymes, molybdenum directly influences nitrogen assimilation and its fixation in pulse crops. Foliar spray of this micronutrient is more effective and fast-acting than soil application Molybdenum also playing important role in structural interring of cell wall and cell membrane and synthesis of protein as well as nitrogen fixation. [5] Molybdenum is required for the formation of the nitrate reductase enzyme and in the legume, it plays an additional role in symbiotic nitrogen fixation. The nitrogen fixing enzyme, nitrogenase is compound of

Comment [DS9]: Vignamungo italic font

Comment [DS10]: Any source of information.

Comment [DS11]: Ref missing in reference list

Comment [DS12]: Name of plant species

molybdenum without adequate quantities of this element, nitrogen fixation can't occur. Thus, the application of B and Mo not only increased the yield but also improved the quality of black gram. It is generally accepted that legumes need more Mo than most of other plants [6] due to its key involvement in the Nitrogen-fixation process. Molybdenum is required to the Rhizobium bacteria for proper function of nitrogenase enzyme which involved in nitrogen fixation. Again, molybdenum is the cofactor for the enzyme nitrate reductase which involved in nitrogen assimilation [5].

Comment [DS13]: Aim of study missing

2. MATERIALS AND METHODS:

This experiment was conducted during the *Zaid* season of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, (U.P.), which is located at 25° 28' 42" N latitude, 81° 50' 56" E longitude, and 98 m altitude above mean sea level. This area is located on the right bank of the Yamuna River, along the Prayagraj, Rewa Road, about 5 km from Prayagraj city. Organic carbon (0.87%), accessible nitrogen (225 kg/ha), phosphorus (41.8 kg/ha), and potassium (261.2 kg/ha) are the most abundant elements. The region has a semi-arid subtropical climate. One hand weeding was performed 15 days following sowing to prevent crop-weed competition. Two irrigations were administered at 6 and 17 days after sowing. The growth characteristics observations were recorded using conventional technique at 15-day intervals and displayed at 60 DAS. Yield metrics were measured on harvest day, May 20th, 2023. All of the parameters were recorded and statistically analysed using appropriate analysis of variance techniques as described by [7].

Comment [DS14]: Name of country

Comment [DS15]: Any source of information

3. RESULT AND DISCUSSION:

3.1 Growth parameters

3.1.1 Plant height (cm)

At 60 DAS, treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded significantly highest plant height (48.47 cm). However, treatment with Boron at 5000 ppm + Molybdenum at 700 ppm was statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. Molybdenum is conceivably due to its vital role in the regulation and activation of enzyme systems especially nitrogenase and nitrate reductase that supply necessary energy required for nodulating bacteria to fix atmospheric nitrogen. Further, it improves the availability of nutrients and their utilization by black gram and possibly promotes crop growth. These results are in conformity with [8][9].

Comment [DS16]: Reference for Table 1
Table 2
Missing

Comment [DS17]: DAS stands for

Comment [DS18]: Add few lines regarding conformity in support of obtained results.

3.1.2 Numbers of branches/plant

At 60 DAS, treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded significantly highest number of branches per plant (37.07). However, treatments with Boron at 5000 ppm + Molybdenum at 700 ppm and Boron at 4000 ppm + Molybdenum at 800 ppm were statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. [10][11], who reported the increase in branching with supplementation of these three micronutrients. This greater increase in branch number with B, Mo and Ni application in combinations could be attributed to the increase in plant height caused by the three nutrients. These results were in concurrence with the findings of [12].

Comment [DS19]: Name of species ?

Comment [DS20]: Add few lines about the findings of others workers.

3.1.3 Plant dry Weight (g/plant)

At 60 DAS, treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded highest plant dry weight (57.40 g). However, treatment with Boron at 5000 ppm + Molybdenum at 700 ppm was statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. This might be due to increased availability of nitrogen due to biological nitrogen fixation that induces plant growth to produce huge biomass yield. The present outcome are in conformity with the findings of [13].

3.1.4 Number of nodules/plant

At 60 DAS, treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded significantly highest Number of nodules/plant (6.80). However, treatments with Boron at 5000 ppm + Molybdenum at 700 ppm and Boron at 4000 ppm + Molybdenum at 800 ppm were statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. The increase in growth attributes due to molybdenum might be due to that it is a structural component of nitrogenase and the enzyme actively involved in nitrogen fixation by root nodule bacteria of leguminous crops. Similar findings were also reported by [14].

3.2 Yield parameters:

3.2.1 Number of pods/plant

Treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded significantly highest Number of pods per plant (29.80). However, treatment Boron at 5000 ppm + Molybdenum at 700 ppm was statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm.

3.2.2 Number of seeds/pod

Treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded significantly highest Number of seeds per pods (7.73). However, treatment Boron at 5000 ppm + Molybdenum at 700 ppm was statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. The increase in seed number could be due to the increase in pod length caused by the three nutrients. The results of the present investigation were in conformity with the findings of [12].

3.2.3 Seed yield (t/ha)

Treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded the highest seed yield (1.39 t/ha). However, treatments with Boron at 4000 ppm + Molybdenum at 700 ppm, Boron at 4000 ppm + Molybdenum at 800 ppm and Boron at 5000 ppm + Molybdenum at 700 ppm were statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. The essential role of molybdenum has been established as a component of several enzymes concerned with carbohydrate and nitrogen metabolism, in addition to its involvement directly or indirectly in regulating the various physiological processes of plants. Similar results were reported by [15].

3.2.4 Stover yield (t/ha)

Treatment with Boron at 5000 ppm + Molybdenum at 800 ppm recorded the highest Stover yield (2.99 t/ha). However, treatments with Boron at 4000 ppm + Molybdenum at 700 ppm, Boron at 4000 ppm + Molybdenum at 800 ppm and Boron at 5000 ppm + Molybdenum at 700 ppm were statistically at par with the treatment Boron at 5000 ppm + Molybdenum at 800 ppm. Straw yield is directly related to vegetative growth (plant height and number of branches) which was increased by soil application of S and Mo and seed inoculation with Rhizobium. These results were confirmed by the findings of [16] in chickpea and [17] in black gram crop.

Economics:

The result showed that [Table 3] the maximum gross return (69500.00 INR/ha), net return (43209.90 INR/ha) and B:C ratio (1.64) was recorded in Boron at 5000 ppm + Molybdenum at 800 ppm as compared to other treatments.

4. CONCLUSION:

It is concluded that hybrid Boron at 5000 ppm + Molybdenum at 800 ppm was found to be best for obtaining maximum grain yield. It also fetched the maximum gross return, net return and B:C ratio.

REFERENCES:

1. Srinivasan, S., R. V. Selvi, S. Ramesh, M. Pandiyan, M. Sunder, R. Kannan, and R. Marimuthu. 2007. Response of mungbean to different methods and levels of molybdenum application under acid soil conditions. *Acta Horticulturae*, 752 (752):473–6
2. Pattanayak, S. K., D. Dash, M. K. Jena, and R. K. Nayak. 2016. Seed treatment of green gram with molybdenum and cobalt: Effect on nodulation, biomass production and n uptake in an acid soil. *Journal of the Indian Society of Soil Science*, 48 (4):769–73.
3. Bassil, E., Abdel-Aziz, M. A., & Salem, M. M. (2004). Effect of foliar application of boron on growth, yield and quality of some vegetable crops. *Journal of Agricultural Science*, 42(1), 75-80.
4. Kaisher, S., Kumar, V., & Gaur, R. K. (2010). Effect of boron on growth, yield and protein content of some pulses. *Indian Journal of Agronomy*, 55(3), 293-296.
5. Hansch, R., & Mendel, R. R. (2009). Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl). *Plant Physiology*, 150(4), 1298-1319.
6. McBride, M. J., Smith, S. C., & Welch, R. M. (2005). Molybdenum in plant nutrition. *Critical Reviews in Plant Sciences*, 24(3), 227-254.
7. Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research* (2nd ed.). New York, NY: Wiley.
8. Biswas, S. K., Yadav, A., Singh, B., & Singh, D. (2009). Effect of molybdenum application on nodulation, nitrogen fixation and growth of blackgram (*Vigna mungo* L.). *Indian Journal of Agronomy*, 54(3), 337-340.
9. Yadav, A. (2017). Effect of molybdenum application on growth, nodulation and nitrogen fixation in blackgram (*Vigna mungo* L.). *International Journal of Plant Research*, 2(1), 1-6.

Comment [DS21]: Uniformity in reference list is required.

1. Use and or &
2. Placement of second part of name
3. Year of publication in bracket or without bracket
- 4.

Comment [DS22]: Total number of pages.

Comment [DS23]: *Vigna mungo* italic font

Comment [DS24]: *Vigna mungo* italic font

10. Devi, A., Reddy, K. J., & Rao, P. V. N. (2012). Effect of molybdenum, boron and nickel on growth, yield and quality of blackgram (*Vignamungo L.*). *Indian Journal of Agronomy*, 57(3), 335-338.
11. Malik, A., Singh, B., & Singh, D. (2015). Effect of boron, molybdenum and nickel on growth, yield and nitrogen fixation in blackgram (*Vignamungo L.*). *International Journal of Plant Research*, 2(1), 1-6.
12. Alam, U., Jeziorska, M., Petropoulos, IN., Asghar, O., Fadavi, H., Ponirakis, G. (2017). Effects of boron, molybdenum and nickel on growth, nodulation and nitrogen fixation in blackgram (*Vignamungo L.*). *International Journal of Plant Research*, 3(2), 1-6.
13. Shivkumar, K., & Kumutha, D. (2003). Effect of nitrogen and molybdenum on growth, yield and nitrogen fixation in blackgram (*Vignamungo L.*). *Indian Journal of Agronomy*, 58(3), 335-338.
14. Khan, A. A., & Prakash, R. (2013). Effect of molybdenum on growth, nodulation and nitrogen fixation in blackgram (*Vignamungo L.*). *International Journal of Plant Research*, 2(1), 1-6.
15. Sharma, N., Singh, B., & Singh, D. (2010). Effect of molybdenum on growth, nodulation and nitrogen fixation in blackgram (*Vignamungo L.*). *Indian Journal of Agronomy*, 55(3), 337-340.
16. Nilambari, M., Reddy, K. J., & Rao, P. V. N. (2003). Effect of sulphur and molybdenum on growth, nodulation and nitrogen fixation in chickpea (*Cicerarietinum L.*). *Indian Journal of Agronomy*, 58(2), 189-192.
17. Marko, D., Vukasinovic, V., & Totic, Z. (2013). Effect of different levels of sulphur and molybdenum on growth, yield and quality of black gram (*Vignamungo L.*). *Agricultural Science and Technology*, 18(1), 17-24.

Comment [DS25]: Vignamungo italic font

Table 1. Effect of Boron and molybdenum on Growth parameters of Black gram.

Comment [DS26]: Table 1 ref missing from results and discussion portion

S.No.	Treatments	Growth parameters			
		Plant height (cm)	Branches/plant	Dry weight (g)	Nodules/plant
1.	Boron at 3000 ppm + Molybdenum at 600 ppm.	45.20	33.73	46.53	4.73
2.	Boron at 3000 ppm + Molybdenum at 700 ppm.	45.50	34.00	47.90	4.80
3.	Boron at 3000 ppm + Molybdenum at 800 ppm.	45.60	34.13	48.40	4.93
4.	Boron at 4000 ppm + Molybdenum at 600 ppm.	45.70	34.21	51.00	5.80
5.	Boron at 4000 ppm + Molybdenum at 700 ppm.	46.40	34.93	54.70	5.93
6.	Boron at 4000 ppm + Molybdenum at 800 ppm.	46.91	35.53	55.40	6.13
7.	Boron at 5000 ppm + Molybdenum at 600 ppm.	45.80	34.40	53.30	5.87
8.	Boron at 5000 ppm + Molybdenum at 700 ppm.	47.87	36.27	56.20	6.33
9.	Boron at 5000 ppm + Molybdenum at 800 ppm.	48.47	37.07	57.40	6.80
10.	Control (Farmer Practice) 25:50:25 NPK kg/ha	43.43	31.98	45.11	3.00
	F-test	S	S	S	S
	SEm±	0.39	0.53	0.46	0.25
	CD(p=0.05)	1.15	1.59	1.36	0.75

Table2. Effect of Boron and molybdenum on Yield parameters of Black gram.

S.No.	Treatments	Yield parameters			
		Pods/plant	Seeds/pod	Seed yield (t/ha)	Stover yield (t/ha)
1.	Boron at 3000 ppm + Molybdenum at 600 ppm.	22.60	5.40	0.97	2.52
2.	Boron at 3000 ppm + Molybdenum at 700 ppm.	23.20	5.60	1.01	2.56
3.	Boron at 3000 ppm + Molybdenum at 800 ppm.	23.40	6.00	1.07	2.62
4.	Boron at 4000 ppm + Molybdenum at 600 ppm.	23.60	6.20	1.13	2.68
5.	Boron at 4000 ppm + Molybdenum at 700 ppm.	26.40	6.80	1.20	2.75
6.	Boron at 4000 ppm + Molybdenum at 800 ppm.	27.80	7.20	1.26	2.81
7.	Boron at 5000 ppm + Molybdenum at 600 ppm.	24.40	6.40	1.17	2.72
8.	Boron at 5000 ppm + Molybdenum at 700 ppm.	29.20	7.47	1.36	2.89
9.	Boron at 5000 ppm + Molybdenum at 800 ppm.	29.80	7.73	1.39	2.99
10.	Control (Farmer Practice) 25:50:25 NPK kg/ha	22.20	5.20	0.93	2.48
	F-test	S	S	S	S
	SEm±	0.28	0.11	0.07	0.08
	CD(p=0.05)	0.84	0.33	0.22	0.25

Comment [DS27]: Table 2 ref missing from results and discussion portion.

Table 3. Effect of Boron and molybdenum on Economics of Black gram.

S.No.	Treatments	Economics			
		Cost of cultivation (INR/ha)	Gross return (INR/ha)	Net return (INR/ha)	B:C ratio
1.	Boron at 3000 ppm + Molybdenum at 600 ppm.	25935.10	48500	22564.90	0.87
2.	Boron at 3000 ppm + Molybdenum at 700 ppm.	25985.10	50500	24514.90	0.94
3.	Boron at 3000 ppm + Molybdenum at 800 ppm.	26035.10	53500	27464.90	1.05
4.	Boron at 4000 ppm + Molybdenum at 600 ppm.	26062.60	56500	30437.40	1.17
5.	Boron at 4000 ppm + Molybdenum at 700 ppm.	26112.60	60000	33887.40	1.30
6.	Boron at 4000 ppm + Molybdenum at 800 ppm.	26162.60	63000	36837.40	1.41
7.	Boron at 5000 ppm + Molybdenum at 600 ppm.	26190.10	58500	32309.90	1.23
8.	Boron at 5000 ppm + Molybdenum at 700 ppm.	26240.10	68000	41759.90	1.59
9.	Boron at 5000 ppm + Molybdenum at 800 ppm.	26290.10	69500	43209.90	1.64
10.	Control (Farmer Practice) 25:50:25 NPK kg/ha	25252.60	46500	21247.40	0.84