

## Original Research Article

### **Influence of Integrated Nutrient Management (INM) on Bulb Yield and Profitability of Onion (*Allium cepa* L.) Crop in Western Uttar Pradesh**

#### **Abstract**

**Aim:** A field experiment was carried out at Horticulture Research Centre, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) 250110 during the Rabi seasons of year 2021-22 and 2022 to fulfil various objectives of yield and profitability of bulbs of onion for western Uttar Pradesh climatic conditions.

**Methodology:** A total of eleven treatments were **exasperated** in Randomized Complete Block Design (RCBD) with three replications.

Comment [W1]: ?

**Result:** Out of the eleven treatments, T<sub>7</sub> - 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each) had a beneficial effect on maximum bulb yield (337.49 q/ha), net return (Rs. 208438.14 ha<sup>-1</sup>) and cost-benefit ratio (4.29) followed by 75% RDF + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each). Whereas, the lowest bulb yield (122.16 q/ha) was observed in T<sub>1</sub> – Control treatment followed by T<sub>8</sub>- 50% RDF + FYM @ 12 t/ha. While the minimum net return (Rs. 48307.20 ha<sup>-1</sup>) and cost-benefit ratio (1.97) were recorded in T<sub>1</sub> – Control treatment followed by treatment T<sub>4</sub>- 75% RDF + FYM @ 6 t/ha.

**Keywords:** Onion, NPKS, FYM, Vermicompost, yield and Economics.

#### **Introduction**

Onion (*Allium cepa* L.) is a widely cultivated and consumed vegetable crop. It is a member of the Amaryllidaceae family and is known for its pungent flavour and characteristic aroma. Onions are grown for their bulbs, and are used in an extensive variety of dishes, including soups, stews, salads, and sauces, and are a staple ingredient in many cuisines around the world. Although they may be produced in a variety of regions and are a cool-season crop, onions demand fertile, well-drained soils with a pH of 6.0 to 6.8. They are typically grown from seeds, sets (small bulbs), or transplants. Onions can be stored for several months under proper conditions of temperature and humidity (Thamburaj and Singh, 2022).

Onion bulbs are low in calories but rich in vitamins and minerals and are a good source of dietary fibre. Onions are also a good source of antioxidants and sulphur-containing

compounds that may have health benefits. They have been proven to assist in reducing inflammation, lowering blood sugar, and enhancing heart health. However, it's important to note that onions may cause digestive discomfort in some people, particularly when consumed in large amounts. (Fageria *et al.*, 2019).

Proper nutrient management is crucial for achieving high yield in onion crops. The goal of integrated nutrient management (INM), a comprehensive strategy for crop nutrition, is to maximise nutrient use effectiveness while lowering environmental impact, boosting profitability, and enhancing soil health (Singh *et al.*, 2018). Onion crop has a high nutrient demand and requires balanced nutrition for optimum growth and bulb yield. INM provides a balanced combination of different organic sources of nutrients and inorganic fertilizers, which ensures the supply of all essential nutrients to the crop.

INM promotes the use of organic matter, which improves soil health and enhances the soil's ability to retain moisture and nutrients. INM is a cost-effective approach to crop nutrition as it reduces the reliance on expensive chemical fertilizers and promotes the use of locally available organic resources. INM reduces the environmental impact of agriculture by promoting the use of organic resources and minimizing the use of chemical fertilizers. INM provides a balanced supply of all essential nutrients, which promotes healthy plant growth and maximizes yield (Singh *et al.*, 2020).

In summary, INM is essential for the sustainable production of onion crops by ensuring a balanced supply of nutrients, improving soil health, maximum profitability and reducing environmental pollution.

## Materials & Methods

A field trial was conducted at Horticultural Research Centre, College of Horticulture, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) 250110 during the Rabi seasons of the year 2021-22 and 2022 to fulfil various objectives of the study. The experiment was laid out in Randomized Complete Block Design (RCBD) and replicated thrice. Total 11 treatments have been tried i.e., T<sub>1</sub> - Control, T<sub>2</sub> - 100% RDF (NPKS @ 120:60:80:40 kg/ha), T<sub>3</sub> - 100% RDF + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each), T<sub>4</sub> - 75% RDF + FYM @ 6 t/ha, T<sub>5</sub> - 75% RDF + VC @ 2 t/ha, T<sub>6</sub> - 75% RDF + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each), T<sub>7</sub> - 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each), T<sub>8</sub> - 50% RDF + FYM @ 12 t/ha, T<sub>9</sub> - 50% RDF + VC@ 4 t/ha, T<sub>10</sub> - 50% RDF + Biofertilizers (*Azospirillum* + PSB @ 5

Comment [W2]: ?

kg/ha each) and T<sub>11</sub> - 50% RDF + FYM @ 6 t/ha + VC @ 2 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each). The soil nature of the trial field was sandy loam with a pH level of 7.70 and available nitrogen (149.80 kg ha<sup>-1</sup>), phosphorous (29.18 kg ha<sup>-1</sup>), potassium (153 kg ha<sup>-1</sup>) and sulphur (10.78 kg ha<sup>-1</sup>). The onion variety NHRDF Red -4 was selected for research reasons. Organic manure, such as FYM, was added into the experimental field two weeks after seedling transplanting, as recommended by the protocols.

Comment [W3]: ?

However, vermicompost was also used in experimental plots during seeding. Nitrogen, phosphorus, potassium, and sulphur were added as urea, di-ammonium phosphate (DAP), muriate of potash (MOP), and sulphur. At the time of transplanting onion seedlings, 50% of nitrogen and the 100% of phosphorus, potassium, and sulphur have been applied in experimental plots. The remaining half-dose of nitrogen has been given in two equal doses 30 and 60 days adhering to transplantation. During the experiment, all cultural practises were carried out at regular intervals based on the crop's needs. The collected data were statistically examined using standard statistical procedures recommended by Gomez and Gomez (1996).

## Result & Discussion


### Effect of Integrated Nutrient Management on Yield of Onion

The results of the bulb yield per hectare during two years apart from pooled mean data have been given in Table 2 & Figure 2. Based on pooled mean data of both of the years clearly shows that the maximum bulb yield (337.49 q/ha) was observed in T<sub>7</sub> - 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each), followed by T<sub>6</sub> - 75% RDF + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each) treatment (317.00 q/ha). While lowest bulb yield (112.16 q/ha) was recorded under control subsequently T<sub>8</sub>- 50% RDF + FYM @ 12 t/ha (234.33 q/ha). The defined release of nutrients in the soil caused by the mineralization of organic manures may have contributed to the higher bulb production with the simultaneous use of farm yard manure, vermicompost, biofertilizers, and inorganic fertilizers. The onion crop obtains a balanced supply of nutrients throughout each stage of its growth as a result of a combination of chemical fertilizers, organic manures, and biofertilizers. This can result in increased bulb formation, larger-sized onions, and overall improved yield. The above outcomes are correspondingly in nearby conformism by the result of Thangasamy *et al.*, (2015), Sharma *et al.* (2018), Rathod *et al.* (2020) and Verma *et al.*, (2021) in *Allium cepa* L.

Comment [W4]: ?

**Table 1: Bulb Yield of different treatments in Onion during both the cropping years 2021-22 and 2022-23.**

Treatments	Bulb Yield (q/ha) 2021-22	Bulb Yield (q/ha) 2022-23	Pooled Yield (q ha <sup>-1</sup> )
T <sub>1</sub> - Control	119.00	125.33	122.16
T <sub>2</sub> - 100% RDF (NPKS @ 120:60:80:40 kg/ha)	261.00	252.00	256.60
T <sub>3</sub> - 100% RDF + Biofertilizers ( <i>Azospirillum</i> + PSB @ 5 kg/ha each)	313.00	301.00	307.00
T <sub>4</sub> - 75% RDF + FYM @ 6 t/ha	222.00	229.66	225.83
T <sub>5</sub> - 75% RDF + VC @ 2 t/ha	249.66	246.66	248.16
T <sub>6</sub> - 75% RDF + Biofertilizers ( <i>Azospirillum</i> + PSB @ 5 kg/ha each)	321.00	313.00	317.00
T <sub>7</sub> - 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers ( <i>Azospirillum</i> + PSB @ 5 kg/ha each)	342.33	332.67	337.49
T <sub>8</sub> - 50% RDF + FYM @ 12 t/ha	231.66	237.03	234.33
T <sub>9</sub> - 50% RDF + VC@ 4 t/ha	241.00	264.00	252.50
T <sub>10</sub> - 50% RDF + Biofertilizers ( <i>Azospirillum</i> + PSB @ 5 kg/ha each)	276.33	286.67	281.49
T <sub>11</sub> - 50% RDF + FYM @ 6 t/ha + VC @ 2 t/ha + Biofertilizers ( <i>Azospirillum</i> + PSB @ 5 kg/ha each)	304.33	295.67	299.99
CD at 5%	26.57	26.90	26.73
SE(m)	9.01	9.12	9.06



**Fig. 1: Bulb Yield of different treatments in Onion during both the cropping years 2021-22 and 2022-23.**

#### **Effect of Integrated Nutrient Management on the Economics of Treatments-**

The data (Table-2 & Figure- 2) indicated that the uppermost cost of cultivation (Rs. 61553 ha<sup>-1</sup>) was noted in 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each). Nevertheless, the lowermost cost of cultivation was recorded (Rs. 49420 ha<sup>-1</sup>) in control. The highest cost input was observed, which might be attributed to the usage of 75% of the required dose of inorganic fertilizers in addition to FYM, vermicompost, and biofertilizers. i.e., *Azospirillum* and PSB. The highest gross return (Rs. 269992 ha<sup>-1</sup>), net return (Rs. 208438 ha<sup>-1</sup>) and cost: benefit ratio (4.36) were recorded in 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each), while the minimum gross return (Rs. 97728 ha<sup>-1</sup>), net return (Rs. 48307 ha<sup>-1</sup>) and minimum cost: benefit ratio (1.97) were observed under control treatment during investigation. The higher gross return under 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each) was primarily due to a greater yield, while the higher net return and benefit: cost ratios were due to lower expenses for cultivation underneath 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each). These outcomes are also

in conformity with the findings of Kumar *et al.*, (2017), Prusty *et al.*, (2019), Nirala *et al.*, (2019) and Chaudhary *et al.*, (2020).

**Table 2: Economics of different treatments in Onion during both the cropping years 2021-22 and 2022-23.**

Treatments	Yield (q ha <sup>-1</sup> ) Pooled	Gross Return (Rs ha <sup>-1</sup> ) @ 800/q	Cost of Treatment (Rs ha <sup>-1</sup> )	Total cost of cultivation (Rs ha <sup>-1</sup> )	Net Return (Rs ha <sup>-1</sup> )	B: C Ratio
T <sub>1</sub>	122.16	97728	00.00	49420.80	48307.20	1.97
T <sub>2</sub>	256.60	205280	10411.02	59831.82	145448.18	3.43
T <sub>3</sub>	307.00	245600	12461.02	61881.82	183718.18	3.96
T <sub>4</sub>	225.83	180664	10883.06	60303.86	120360.14	2.99
T <sub>5</sub>	248.16	198528	10883.06	60303.86	138224.14	3.29
T <sub>6</sub>	317.00	253600	9933.06	59353.86	194246.14	4.27
T <sub>7</sub>	337.49	269992	12433.06	61853.86	208138.14	4.36
T <sub>8</sub>	234.33	187464	11355.28	60776.08	126687.92	3.08
T <sub>9</sub>	252.50	202000	11355.28	60776.08	141223.92	3.32
T <sub>10</sub>	281.49	225192	7405.28	56826.08	168365.92	3.96
T <sub>11</sub>	299.99	239992	13405.28	62826.08	177165.92	3.81




Fig. 2: Economics of different treatments in Onion during both the cropping years 2021-22 and 2022-23.

### Conclusion

In conclusion, integrated nutrient management positively influences onion yield by ensuring a balanced and sustained supply of nutrients, improving soil health and fertility, enhancing weed and pest management, promoting environmental sustainability, and providing cost-effective solutions. Adopting INM practices can lead to higher onion yields, better crop quality, and increased profitability for farmers. In accordance with the study's findings, it concluded that onion bulb production was shown to be higher with treatment T<sub>8</sub>- 75% RDF + FYM@ 2 t/ha + VC@ 1 t/ha + Biofertilizers (*Azospirillum* + PSB @ 5 kg/ha each). It was also noted that the above treatments had the best cost-benefit ratio.

### Future Scope

However, it's important to note that the successful implementation of Integrated Nutrient Management for onion bulb yield and economic feasibility will depend on various factors, such as regional climate, soil characteristics, crop management practices, and market dynamics. Continuous research, field trials, and farmer education and training will play a crucial role in realizing the full potential of Integrated Nutrient Management for sustainable onion bulb production and Profitability for farmers.

### References

1. **Chaudhary SK, Yadav SK, Mahto DK, Singh SK, Singh BB and Sinha N.** Effect of Different Organic and Inorganic Sources of Nutrients on Growth and Yield of Rabi Onion (*Allium cepa* L.). *Int. J. Curr. Microbiol. App. Sci.* 2020; 9(10): 2851-2858.
2. **Fageria MS, Chaudhary BR, and Dhaka RS.** Vegetable Crops Production Technology. *Kalyani Publishers*, 2019; pp-135
3. **Gomez AK and Gomez AA.** Statistical procedure for Agriculture Research. John Willey and Sons Pnc, New York 1996.
4. **Kumar S, Garhwal OP and Sharma A.** Effect of Integrated nutrient management on growth and yield attributes of kharif onion (*Allium cepa* L.). *Int. J. Pure App. Biosci.* 2017; 5 (3): 854-857.
5. **Nirala K, Punetha S, Pant SC and Upadhaya S.** Effect of Various Organic Manures with Biofertilizers on Growth, Yield and Economics of Onion (*Allium cepa* L.). *Int. J. Curr. Microbiol. App. Sci.* 2019; 8(01): 3092-3099.
6. **Prusty M, Mishra N, Kar DS and Pal S.** Effect of Integrated Nutrient Management on Growth and Yield of Onion (*Allium cepa* L.) cv. Bhima Super. *International Journal of Agriculture Sciences.* 2019;11(4): 7910-7912.
7. **Rathod NB, Khandare RN and Karle AS.** Effect of organic, inorganic and integrated nutrient management on yield attributes, production of onion and soil properties, under vertisol of Maharashtra. *International Journal of Chemical Studies,* 2020; 8(1): 937-940.
8. **Sharma SK, Garhwal OP, Mahala P and Yadav TV.** Influence of integrated nutrient management on yield attributes and economics of kharif onion (*Allium cepa* L.) under loamy sand soils. *International Journal of Current Microbiology and Applied Sciences,* 2018;7(6): 2806-2811.
9. **Singh J, Singh MK, Kumar M, Gupta A and Singh KP.** Growth, Yield and Quality Parameters of Cucumber (*Cucumis sativus* L.) as Influenced by Integrated Nutrient Management Application. *Int. J. Curr. Microbiol. App. Sci.* 2020; 9(10): 1455-1462.
10. **Singh J, Singh MK, Kumar M, Kumar V, Singh KP and Omid AQ.** Effect of integrated nutrient management on growth, flowering and yield attributes of cucumber (*Cucumis sativus* L.). *International Journal of Chemical Studies* 2018; 6(4): 567-572.
11. **Thamburaj S and Singh N.** Textbook of Vegetables, Tuber crops and spices. *ICAR. New Delhi*, pp. 469; 2022.

12. **Thangasamy A and Lawande KE. (2015).** Integrated nutrient management for sustainable onion production. *Indian J. Hort.* **72**(3): 347-352.
13. **Verma N, Jaiswal RK, Ali SA and Dhakad RK.** Impact of Biofertilizers on Growth and Yield Attributes of Onion (*Allium cepa* L.) var. AFLR, *Int. J. Curr. Microbiol. App. Sci.* 2021;10(02): 931-938.

UNDER PEER REVIEW