

THE PERFORMANCE OF L-THEANINE AS AN IMMUNOSIMULANT TO REDUCE STRESS LEVELS IN MOTIVANT FISH (*Helostoma temminckii*)

ABSTRACT

Stressed animal fish are easily infected by several diseases which will affect their health, thus reducing their reproductive performance. Immunostimulant is a substance that can enhance or stimulate the fish's immune system by interacting directly with cells that activate the immune system. The aim of the study was to analyze the performance of L-theanine as an immunostimulant to reduce stress levels in monkfish. This research was carried out at the Wet Laboratory of the Faculty of Fisheries and Maritime Affairs, University of Lambung Mangkurat, South Kalimantan Province. The treatment that will be given is as follows, namely Treatment A was feed added with a dose of L-Theanine 0 mg/kg . Treatment B feed added with a dose of L-Theanine 300 mg/Kg . Treatment C feed added with a dose of L-Theanine 600 mg/Kg . Treatment D feed added with a dose of L-Theanine 900 mg/kg . The observed haematological conditions were hematocrit, erythrocytes and blood glucose. Substitution of feed with a dose of L-Theanine 900 mg/kg was able to maintain immunostimulants to reduce stress levels in animal fish according to hematological measurements in the form of hematocrit, erythrocytes and blood glucose.

Keywords: Biawan Fish, L-theanin, Immunistulants and Stress.

INTRODUCTION

Biawan fish (*Helostoma temminckii*) is a native Indonesian fish found in several rivers in Sumatra and Kalimantan. Biawan fish live in rivers, tributaries and inundation areas from upstream to downstream, even in river estuaries which are hilly and forested along the sides. This fish has high economic value, has prospects for developing aquaculture with great opportunities, the selling price is quite expensive, is an important commodity in the freshwater fish business but this fish is still rarely cultivated to date (Rohana, 2015).

Stress is defined as the effect of any environmental changes or encouragement to achieve homeostasis or changes in stability reaching normal limits. Biawan fish are very sensitive to any changes both externally and internally, this condition causes them to be susceptible to stress (Utomo, 2010). Stressed animal fish are easily infected by several diseases that will affect their health, thereby reducing reproductive performance (eg growth, yield, survival and feed efficiency). Immunostimulants are substances that can enhance or stimulate the immune system of fish by interacting directly with cells that activate the immune system (Prianto, 2006). The stress response of fish can be divided into primary, secondary and tertiary phases. The primary effect on stress is an endocrine response that stimulates metabolic and osmotic adaptation (secondary phase) (Bijaksana, 2010). The mechanism of action of immunostimulants in stimulating the immune system is by increasing the activity of phagocytic cells (Tang, 2001).

The L-theanine compound is known as one of the neutron conductors found in the brain. This compound is absorbed in the intestinal brush border membrane because it is fat soluble, while it is transported into the brain via the preferred transport system of leucine from the blood lining the brain. theanin also increases GABA levels in the brain which brings a feeling of comfort (Abdullah, 2008). Direct administration of L-theanine into the striatum of the brain by microinjection also causes a significant increase in DA release in a dose-dependent manner. As mentioned that the release of DA, one of the neurotransmitters, has a major effect on human emotions, these results suggest that L-theanine can affect metabolism and/or release of several neurotransmitters in the brain (Gustiana, 2013).

Irsadi (2022) the L-Theanine content in green tea can reduce stress levels in fish. L-Theanine can be a progressive muscle relaxant to reduce anxiety levels. This condition is the basis that giving β -glucan can manage stress and increase immunity. L Theanin as an immunostimulant can be used as an alternative in aquaculture during the rematuration process of broodfish. There is a need for more studies regarding the use of L Theanin as an immunostimulant to enhance the non-specific immune

system in fish which is very necessary in an effort to reduce stress levels in fish which can inhibit the rematuration process (Rejeki et al, 2014). The aim of the study was to analyze the performance of L-theanine as an immunostimulant to reduce stress levels in monkfish.

MATERIALS AND METHODS

LOCATION AND TIME OF RESEARCH

This research was conducted at the Wet Laboratory of the Faculty of Fisheries and Maritime Affairs, Lambung Mangkurat University, South Kalimantan Province. The biawan fish to be used in this study came from the Bincau Freshwater Fish Hatchery, South Kalimantan. The required number of broodstock is 60 female broodstock, body weight from 40 grams to 100 grams. Maintenance of brood fish during the 15 day study by providing artificial feed which was given pellet feed which was added with L – Theanine 3 times a day morning at 07:00, noon at 12:00 and evening 17:00 with a normal dose of 5% of the total weight of the fish .

EXPERIMENTAL DESIGNS

The treatment that will be given is as follows, namely:

- Treatment A was feed added with a dose of L-Theanine 0 mg/kg
- Treatment B feed added with a dose of L-Theanine 300 mg/Kg
- Perlakuan C pakan yang ditambahkan dengan dosis L-Theanine 600 mg/Kg
- Perlakuan D pakan yang ditambahkan dengan dosis L-Theanine 900 mg/kg

Adapun kondisi hematologis yang diamati adalah hematokrit, eritrosit dan glukosa darah.

DATA ANALYSIS

The parameter data observed in this study were analyzed descriptively and the results were presented in graphs and tables. Covariance analysis (ANCOVA) was applied to examine differences between treatments in terms of growth patterns. One-way ANOVA test was used to ascertain whether there was a difference in the length-weight measurement of snakehead fish reared in indoor containers between the four treatments. If there is, then the test (HSD) is applied. All tests were analyzed at the 0.05 significance level using the Microsoft excel.

RESULTS AND DISCUSSION

Analysis of the performance of L-theanine as an immunostimulant to reduce stress levels in monkfish was carried out by haematological testing. Hematology is a branch of medical science that studies blood , blood-forming organs and their diseases . Hematology studies disorders, diagnosis, treatment, recovery and prevention of diseases that attack blood and its components (Astuti, 2003). The hematological conditions observed in this study were the hematocrit, erythrocytes and blood glucose of the biawan fish.

A: Hematocrit

Hematocrit is the percentage of the volume of erythrocytes in fish blood or the ratio between the volume of blood cells and blood plasma. The hematocrit can provide an indication of the health of the fish and help determine the occurrence of abnormalities due to the use of immunostimulants. The hematocrit value can be calculated by the number of red blood cells contained in fish blood (January, 2012).

The hematocrit content also depends on nutritional factors, age, sex, body size. The results of the calculation of the hematocrit of the monkfish during the study can be seen in the following figure:




Figure 1. Hematocrit Value of Biawan Fish

Information:

- Treatment A feed substitution with a dose of L-Theanine 0 mg/kg
- Treatment B with feed substitution at a dose of L-Theanine 300 mg/Kg
- Treatment C feed substitution with a dose of L-Theanine 600 mg/Kg
- Feed substitution D treatment with a dose of L-Theanin 900 mg/kg

Based on Figure 1 it can be seen that the percentage of hematocrit for each treatment was different. But overall, the hematocrit value of the Bawan fish is still within normal limits. The value of hematocrit analysis of monkfish that was given treatment with feed substitution with a dose of L-Theanine, namely A as a control (42.40%), Treatment B (29.37%), Treatment C (38.70%) and Treatment D (45.73%). The results of the normality test for the hematocrit of the biawan fish obtained a Li Max value of $0.91 < Li \text{ Table } 5\% \text{ } 0.242$, so it can be concluded that the data is normally distributed. The results of the Barlett variance homogeneity test obtained the value $X^2 \text{ count } -7.284 < X^2 \text{ table } 5.991$, which means homogeneous. The results of the ANOVA analysis of diversity showed that the calculated F value was $1,900 < F \text{ table } 5\%$, namely 4.07, which meant that there was no difference between treatments.

The highest hematocrit value during the study was in treatment D (45.73%). This condition is due to the influence of L-Theanin contained in the feed. According to Abdullah (2008) in Faizah (2013) fresh water fish are said to be healthy if their hematocrit levels range from 22-60%. A hematocrit concentration of $<22\%$ indicates that the fish is anemic and the fish may be sick or stressed. Giving L-Theanine to feed can help increase the immunity of the animal fish.

Hematocrit shows the percentage of solids (red blood cell levels, etc.) to the amount of blood fluids. The higher the hematocrit percentage, the thicker the blood concentration. This happens because there is seepage (leakage) of fluid out of the blood vessels while the amount of solids remains, so the blood becomes thicker. A decrease in hematocrit occurs when the body experiences acute blood loss, sudden blood loss, for example during anemia, leukemia, chronic kidney failure, malnutrition, deficiency of vitamins B and C. An increase in hematocrit above normal occurs in dehydration, severe diarrhea, eclampsia, the effects of surgery, burns and others.

Hematocrit is the ratio of the number of red blood cells to the total volume of blood calculated as a percentage. If your hematocrit is known to be 20%, it means that there are 20 milliliters of red blood cells per 100 milliliters. The description of the blood of an organism can be used to determine the health condition being experienced by fish, one of which is seen from the percentage of hematocrit which has an important role as a defense against bacterial attacks that enter the fish's body. When fish are stressed, the hematocrit value will tend to decrease. Stress in fish also causes physiological deviations and hormonal imbalances, causing blood components to also change.

The results of the analysis provide an illustration that at each sample point the Hematocrit levels varied in each treatment group. With an average value of initial treatment = 41.5%, Treatment A = 42.4% Treatment B = 29.3% Treatment C = 38.7% Treatment D = 45.7% which is spread in each observation group above the minimum recommended level. Hematocrit can be influenced by season, food and hormonal factors. According to Fange (1992), in tolerant fish, hematocrit levels are generally relatively constant between 20-40%. The normal value of tilapia hematocrit ranges from 27-37% (Farouq, 2011). Tilapia hematocrit values range from 28.00 – 35.13%. Thus the Hct value of tilapia is classified as healthy.

Calculation of the percentage of hematocrit is one of the approaches used to be able to recognize the fish's body when stressed. Hematocrit is the percentage of the volume of red blood cells (erythrocytes) in fish blood. If the hematocrit value is less than 25%, it indicates anemia (Kuswardani,

2007). The mechanism for changing the percentage of hematocrit during stress starts from receiving information on the causes of stress factors by the receptor organs. Furthermore, the information is conveyed to the hypothalamus part of the brain through the nervous system. The hypothalamus instructs the chromaffin cells to secrete catecholamines via sympathetic nerve fibers. The presence of these catecholamines will activate lipopolysaccharide which attacks blood components whose function can reduce the hematocrit in fish. The results of examination of the hematocrit can be used as an indicator to determine the health condition of fish, a hematocrit value of less than 25% indicates anemia. A low hematocrit can also indicate contamination, lack of food for fish, low protein feed, vitamin deficiency, stress or infection (Aryani, 2014).

Stress in fish also causes physiological deviations and hormonal imbalances, causing blood components to also change. Changes in blood picture and blood chemistry, both qualitatively and quantitatively, can determine the state of health. A decrease in the hematocrit value indicates the uncomfortable condition of an organism and causes anemia (Nirmala et al., 2012). A hematocrit value below 30% indicates a deficiency of erythrocytes. The number of red blood cells, hematocrit, and hemoglobin decreased with increasing concentration of mercury in the rearing medium. The decrease in hematocrit and hemoglobin levels in fish blood due to mercury is affected by contamination, absorption and accumulation of mercury which can cause anemia in fish.

A decrease in the value of the hematocrit can occur if, at the time of stress, the fish cannot maintain homeostatic conditions where at the General Adaptation Syndrome (GAS) stage a defense reaction occurs against the stressor which causes changes in heart rate, changes in respiration, and blood supply capacity. When a defense reaction occurs, the body will provide a primary response in the form of increased secretion of corticosteroids and catecholamines.

The measured hematocrit value is directly related to the number of erythrocytes and hemoglobin levels. When fish experience stress, the contraction of the spleen will decrease so that the circulation of red blood cells becomes weak, so that the hematocrit value decreases. The occurrence of stress on fish will cause an increase in heart rate, blood pressure, increase in blood glucose and release of cortisol. At times of stress, cortisol suppresses the immune system, causing lymphocytes to increase and increased cortisol secretion can also cause a decrease in monocytes and basophils. In the body of fish, monocytes play a role in phagocytosis of foreign bodies. These cells have the ability to kill various types of pathogenic agents, including bacteria and worm larvae.

The function of basophils is related to antigen sensitivity, stress symptoms and phagocytosis. If fish are exposed to a stressor for a long time, then chronic stress will occur in fish which results in a decrease in the immune system so that fish are susceptible to attack by bacteria, fungi and parasites. A decrease in the immune system is usually followed by mortality in fish (2019).

Stress consists of 3 components, namely stressor, process and response. The term stress does not only refer to the source of stress but the interrelationships between the three. A stressor is an event, situation, or object that causes stress and causes a stress reaction as a result. There are four kinds of stress factors, namely chemical stressors, which can be in the form of water quality, pollution, the presence of nitrogen compounds and metabolic waste substances. Biological stressors, can be density, parasites, microbes, fungi and bacteria. Physical stressors, in the form of sound, and light intensity. Procedural stressors include handling, transportation and disease management (Linder et al., 2013).

B. Erythrocytes

Erythrocytes are also known as red blood cells. The red color of the erythrocytes is due to the presence of hemoglobin. Hemoglobin is composed of the iron compound heme and a type of protein, namely globin. The main role of erythrocytes is to transport oxygen from the lungs throughout the body of the fish. The results of the analysis of the monkfish's erythrocytes during the study period can be seen in the following figure:




Figure 2. Erythrocyte Value of Biawan Fish ($\times 10^6 / \mu\text{L}$)

Information:

- Treatment A feed substitution with a dose of L-Theanine 0 mg/kg
- Treatment B with feed substitution at a dose of L-Theanine 300 mg/Kg
- Treatment C feed substitution with a dose of L-Theanine 600 mg/Kg
- Feed substitution D treatment with a dose of L-Theanin 900 mg/kg

Based on Figure 2, it can be seen that the percentage of erythrocytes in each treatment was different. But overall, the value of the red blood cells is still within normal limits. The value of the erythrocyte analysis of animal fish treated with feed substitution with a dose of L-Theanine, namely A as a control ($3.83 \times 10^6 / \mu\text{L}$), Treatment B ($3.67 \times 10^6 / \mu\text{L}$), Treatment C ($3.71 \times 10^6 / \mu\text{L}$) and Treatment D ($3.18 \times 10^6 / \mu\text{L}$). The results of the normality test for the liliefors erythrocytes of the biawan fish obtained a Li Max value of $0.438 < \text{Li Table } 5\% \text{ } 0.242$, so it can be concluded that the data is normally distributed. The results of the Barlett variance homogeneity test obtained the value $X^2 \text{ count } 1.685 < X^2 \text{ table } 5.991$, which means homogeneous. The results of the ANOVA analysis of diversity showed that the calculated F value was $0.541 < F \text{ table } 5\%$, namely 4.07, which meant that there was no difference between treatments.

The highest erythrocyte value during the study was given the animal feed containing L-Theanine, namely in treatment A ($3.83 \times 10^6 / \mu\text{L}$). In teleost fish, the normal number of normal red blood cells or erythrocytes ranges from $1.05 \times 10^6 - 3.0 \times 10^6 \text{ cells/mm}^3$ (Royan *et al.*, 2014). The value of the erythrocytes in this study ranged from $3.17-3.83 \times 10^6 / \mu\text{L}$. This amount is still within the reasonable limits of normal fish erythrocytes. This condition means that the addition of L-Theanine in the feed has a positive impact on the red blood cells' erythrocytes.

Hidayah (2019) the value of erythrocytes given to the feed means that the addition of the plant extract has a positive impact on the erythrocytes of the biawan fish. Erythrocytes ranged from $2.07-3.05 \times 10^6 / \mu\text{L}$. The red *blood cell count* (RBC) is the number of **erythrocytes** per cubic millimeter or microliter. Robert (1978) As with hematocrit, low erythrocyte levels indicate anemia. Meanwhile, high levels indicate that fish are under stress (Wedemeyer and Yasutake, 1977).

The shape and small size of erythrocytes is an adaptive value for oxygen and carbon dioxide, namely as a carrier that can quickly spread throughout the network. The normal range for the number of fish erythrocytes in general is $20,000-3,000,000 \text{ cells/mm}^3$, thus the number of fish erythrocytes studied is normal with a healthy category (Oktavia, 2011), animals that actively move will have a lot of erythrocytes because they will consume a lot of oxygen, Erythrocytes function as oxygen transport in the blood. The number of erythrocytes varies depending on age, sex, hormones and environment (Sarkiah *et al.*, 2016).

C. Blood Glucose

Blood glucose is glucose found in the blood which is formed from carbohydrates in food and stored as glycogen in the liver and skeletal muscles (Joyce, 2007). Blood glucose is the main source of energy and an important element to support the metabolism of fish cells, especially brain cells. The results of the blood glucose analysis of the monkfish during the study period can be seen in the following figure:




Figure 3. Biawan Fish Blood Glucose Value (mg/dL)

Information:

- Treatment A feed substitution with a dose of L-Theanine 0 mg/kg
- Treatment B with feed substitution at a dose of L-Theanine 300 mg/Kg
- Treatment C feed substitution with a dose of L-Theanine 600 mg/Kg
- Feed substitution D treatment with a dose of L-Theanine 900 mg/kg

Based on Figure 3, it can be seen that the percentage of blood glucose in the monk fish was different for each treatment. But overall the blood glucose value of the monkfish is still within normal limits. Value of blood glucose analysis of animal fish treated with feed substitution with a dose of L-Theanine, namely A as a control (48.67 mg/dL), Treatment B (46.00 mg/dL), Treatment C (44.00 mg/dL) and Treatment D (65.33 mg/dl). The results of the normality test for the blood glucose liliefors of monkfish obtained a Li Max value of $0.303 < Li \text{ Table } 5\% 0.242$, so it can be concluded that the data is normally distributed. The results of the Barlett variance homogeneity test obtained the value $X^2 \text{ count } -9.422 < X^2 \text{ table } 5.991$, which means homogeneous. The results of the ANOVA analysis of diversity showed that the calculated F value was $1.370 < F \text{ table } 5\%$, namely 4.066, which meant that there was no difference between treatments.

The highest blood glucose value during the study was given to the fish feed containing L-Theanine, namely in treatment D (65.33 mg/dL). Generally, fish blood glucose levels that are considered normal range from 40-90 mg/dL. If the blood glucose state of the fish is not normal, it will interfere with the life of the fish and can even cause death. Giving L-Theanine to feed can help maintain and increase the immunity of the cattle fish.

The content of L-Theanine has been shown to be able to reduce blood glucose levels in Biawan fish that experience stress due to environmental changes during transportation. From the test results it can be concluded that treatment B with L-Theanine substitution 300 mg/L is the best treatment. This is based on the efficiency of the ingredients and the results which show normal levels in fish and the stability of the measurement results. Normal fish blood glucose levels contain 40-90 mg/dL, the blood glucose content is almost the same as blood glucose in humans, namely 70-110 mg/dl (Rahardjo et al., 2011).

High blood glucose levels stimulate the thyroid gland and increase the production of thyroxine. High thyroxine can trigger lymphocytopenia (low lymphocytes) in the blood. Then the sympathetic nervous system overreacts, which causes lymphatic contraction, increasing the respiratory rate and blood pressure. Stress in fish is defined as a number of physiological responses that occur when fish are trying to maintain homeostasis. When a fish experiences stress, it responds by developing a new homeostatic state by changing its metabolism.

Stress can increase blood glucose levels. Physically, stress can be seen from the behavior of fish, such as less aggressive movements, decreased fish appetite, and fish body color becomes dark. Changes in fish blood glucose can also be an indication of stress in fish caused by external factors in the aquatic environment such as drastic environmental changes, domestic and industrial waste pollution. Changes in environmental conditions will cause a high demand for blood glucose supply.

CONCLUSION

Substitution of feed with a dose of L-Theanine 900 mg/kg was able to maintain immunostimulants to reduce stress levels in animal fish according to hematological measurements in the form of hematocrit, erythrocytes and blood glucose.

REFERENCES

1. Abdullah, Y. 2008. The effectiveness of paci-paci leucas lavandulaefolia leaf extract for the prevention and treatment of mas motile aeromonad septicemia in terms of macro pathology and hematology of African catfish *Clarias* sp. Thesis. Faculty of Fisheries and Marine Science. Bogor Agricultural Institute.
2. Ari Hepi Yanti, ELTRS (2019). Hematology Profile of Snakehead Fish (*Channa striata* Bloch, 1793). *Protobiont Journal* , 8 (2), 283–289. <https://doi.org/10.26418/protobiont.v8i2.32474>
3. Aryani, N., Efawani & A. Nur. 2014. Enrichment of artificial feed with vitamin E for gonadal maturation of mali fish (*Labeobabbarbus festivus*). *J. Fisheries and aquatic*. 2: 126-129.
4. Astuti, Asrini Budi. 2003. Interaction of Pepticide and Bacterial Infection (*Aeromonas hydrophyla*) in Doumbo Catfish (*Clarias* sp). Thesis Research Report. Faculty of Fisheries and Marine Science. IPB. Bogor
5. Astuti . 2003 . Relationship of Communication Quality and Stress Tolerance in. Marriage. *Journal of Sukma* Volume 2 No. 1 Pgs 52-60. Yogyakarta
6. Bijaksana, U. 2010. Reproductive Physiological Study of Snakehead Fish (*Channa Striata* Blkr) in Containers and Swamp Waters as a Domestication Effort. Dissertation. Bogor Agricultural Institute.
7. Faizah, Hana. 2013. Effectiveness of noni leaf extract (*Morinda citrifolia* L) against MAS (Motile *Aeromonas* Septicemia) attacks on sangkuriang catfish (*Clarias* sp). JTAM. Faculty of Arts, University of Lambung Mangkurat. Banjarbaru.
8. Farouq, A. 2011. Application of Probiotics, Prebiotics and Synbiotics in Feed to Increase Immune Response and Survival of *Tilapia Oreochromis niloticus* infected with *Streptococcus agalactiae*.
9. Gustiana , M. 2013. Length-Weight Relationship and Condition Factors for Mulletts (*Liza subviridis*) in the Opak Estuary, Bantul Regency. Thesis. Gadjah Mada University 2013. 46 pages.
10. Irshad Hillman (2022). Provision of Green Tea and Progressive Muscle Relaxation Techniques Have an Influence on Reducing the Anxiety Level of UAS Pre-Test Nursing Students at STIKIM Jakarta in 2021. *Indonesian Journal of Health Interprofession*. 2(1): 234-242
11. January, Wiwik. 2012. The Effectiveness of Bioactive Cashew Leaves of Guava (*Psidium gunjava* L) With Different Doses To Treat MAS Disease (Motile *Aeromonas* Septicaemia) in Dumbo Catfish (*Clarias gariepinus*). Thesis of the Faculty of Fisheries. Lambung Mangkurat University. Banjarbaru.
12. Prianto , Eko, Husnah, Syarifah Nurdawaty, and Asyari. 2006. Eating Habits of Biawan Fish (*Helostoma teminckii*) in Lake Sababila Watershed Barito, Central Kalimantan. *Eating Habits of Biawan Fish*, 12(1):161-166.
13. Rejeki, S., Royan, F., & Haditomo, AHC (2014). The Effect of Different Salinities on the Blood Profile of *Tilapia (Oreochromis niloticus)*. *Journal of Aquaculture Management and Technology* , 3 (2), 109–117. <https://ejournal3.undip.ac.id/index.php/jamt/article/view/5239>
14. Robert, RJ 1978. *The Bacteriology of Teleostei in Fish Pathology*. Ballier Tindall London. 205-308 pp.
15. Rohana. 2015. Hatchery Techniques for Biawann Fish (*Helostoma temminckii*) at the Central Fish Cultivation Center (BBIS) Pavilion, Mempawah Regency. Faculty of Fisheries and Marine Science. Aquatic Culture Department. Pontianak Muhammadiyah University.
16. Sarkiah , Rimalia, A. and Iskandar, R. 2016 . Health of *Tilapia Gift (Oreochromis niloticus)* in Cage Business in Masta Village, Tapin, South Kalimantan
17. Tang, UM and Affandi, R. 2001. Reproductive Biology of Fish. Research Center for Coastal and Water Areas, University of Riau, Pekanbaru. 153 p.
18. Utomo , AD, S. Adjie., SN Aida, and K. Fatah. 2010 . *Fish Resources Potential* . In the Musi River Basin, South Sumatra. Palembang: Research Center.
19. Wedemeyer, GA and Yasutke. 1977. Clinical Methods for The Assessment on The Effect of Environmental Stress on Fish Health. Technical Paper of The US Department of The Interior Fish and the Wildlife Service, 89 : 1-17.