

Seasonal photosynthesis variations of dominant tree species used in different urban landscapes

Abstract

Photosynthesis is the only natural process that converts photon energy into chemical energy, and it is responsible for 90–95% of plant biomass accumulation and hence growth. Environmental factors such as light, temperature, and CO₂ concentration in the atmosphere are major factors that influence photosynthesis, and excess of these can cause stress on plants. The photosynthetic response of tree species is found to vary to these factors. Therefore, the photosynthetic response of tree species can be a good indicator to assess these environmental factors that are changing due to urbanization and can be used to mitigate these factors as well by cultivating suitable tree species. The present study provides information on the seasonal variations in photosynthetic activity in six tree species. In the present study during the month of May (late summer to early monsoon) all the tree species showed higher photosynthetic rates $30.64 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. It is also noticed that transpiration rate and water use efficiency was more in the month of May compared to other months. A decrease in net photosynthesis and transpiration is observed among the tree species under low rainfall conditions 285.2 mm. It is also observed that photosynthesis decreased where the evaporative demand is very high.

Keywords: Tree species, Photosynthesis; Urbanization, Temperature;

1. INTRODUCTION

The changes in climate at a global scale are predicted to have serious implications, especially in tropical countries in terms of increased natural perturbations that influence food security, health, and the environment. By the end of this century, average global temperatures are predicted to rise by 1.8–4.0 °C [1]. The rate of respiration and photosynthetic capacity of leaves are two gas exchange traits that are temperature sensitive [2, 3]. In recent decades, there has been a significant increase in studies to understand the pros and cons of global warming on photosynthesis [4, 5].

Plants have many complex mechanisms that allow them to adapt to environmental conditions and survive despite the impacts of various stressors. Photosynthetic activity, as one of the

Formatted: Portuguese (Brazil)

Formatted: Highlight

Comment [A1]: It is not true, some values are bigger in April.

Comment [A2]: This affirmative is confusing. The months with very low precipitation were February and March, and photosynthesis rates were more than in June and July.

Formatted: Highlight

Comment [A3]: It is not true due comment before.

Formatted: Highlight

most important metabolic processes, is a good indicator for the assessment of the influence of various unfavorable environmental conditions on their physiological condition and development in the terrestrial habitat [6, 7]. Photosynthetically active radiation (PAR) is referred to as solar radiation with a wavelength between 400 and 700 nm [8, 9]. Radiation absorption is carried out by the photosynthetic pigments chlorophyll and carotenoids, which are found in light-harvesting complexes (LHCs). They only expend a portion of the energy they have gathered. Some of it is released as chlorophyll fluorescence or lost as heat. The Kautsky kinetics is used to describe the induction of chlorophyll fluorescence [10]. Fluorescence is emitted by chloroplasts and is connected to all other physiological and metabolic activities taking place inside the plant cell. Therefore, any environmental change that has an impact on these processes will likewise have an impact on photosynthesis [11, 12]. Typically, topographic differences in species composition and tree growth are observed. A slope's topography affects a variety of environmental factors, such as the chemical composition of the soil, its moisture regime, and how it is exposed to weather and sunlight. The connection between various environmental factors and the development or physiology of trees has been the subject of numerous studies [13], but little is known about how topographic position affects plant photosynthetic activity in actual field conditions.

Photosynthesis is the mechanism that takes place naturally which converts photon energy into chemical energy and it is responsible for 90–95% of plant biomass accumulation [14]. About 40% of a plant's dry mass consists of carbon which is fixed only through photosynthesis. The net photosynthetic rate is often influenced by environmental factors such as irradiance, temperature, and water supply, and also by leaf age, leaf position, and leaf developmental stage (15, 16, 17). Trees act as a CO₂ sink which captures carbon from the atmosphere and stores the same in the form of biomass while releasing oxygen to the air through the process of photosynthesis [18]. Conservation and restoration of green spaces comprising trees, therefore, is an important approach to improving the environmental quality which plays a very important role in sustaining a clean environment not only by sequestering atmospheric carbon and releasing oxygen.

The present study explains the seasonal variations in the photosynthetic capacity of six tree species grown under controlled conditions which were investigated in this research to clarify the impact of summer, winter, and rainy seasons on the photosynthetic activity of plants.

2. METHODOLOGY

In the present study sapling of tree species namely *Grevillea robusta*, *Spathodea campanulata*, *Swietenia mahagoni*, *Bauhinia purpurea*, *Pongamia pinnata*, and *Peltophorum pterocarpum* were grown under controlled conditions in the greenhouse facility of the Department of Forestry & Environment Sciences, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India. One-year-old saplings were planted in grow bags with a capacity to hold 25kg of the potting mixture. The recommended package of practices was followed and seedlings were irrigated every day twice during the summer months and once in the rainy season. The saplings were allowed to stabilize for one month after transplanting before recording the observations. [What? \(Various\)](#) physiological parameters were assessed periodically. The six species studied in this experiment are the dominant tree species out of 44 tree species present across the urban landscapes of Bengaluru as avenue trees, in the parks, lakes, and home gardens.

2.1 Climate and weather conditions

University of Agricultural Sciences, Bangalore, is located at 13° 05" N latitude and 77° 34" E longitude and an altitude of 924 meters. The annual rainfall ranges from 528 mm to 1374.4 mm with a mean of 915.8 mm.

The weather parameters of different seasons were procured from the Department of Agrometeorology, University of Agricultural Sciences, Bangalore, Karnataka. The rainfall during the first three months of the study period (from February to April) was negligible and for the rest of the three months, it was almost uniform. Rainfall had no effect on the plants used in this study as they were irrigated but influenced the surrounding temperature (Table 1). [\(The influence of the rainfall is the moisture of the air that controls the opening and closing of stomata\)](#) The maximum temperature varied from 27.30 to 33.50 °C and the minimum temperature varied from 16 to 20.10 °C during the study period (Table 1).

Table 1: Distribution of rainfall (mm), maximum temperature, and minimum temperature during the experimental period.

Months (2022)	Minimum Temperature (°C)	Maximum Temperature (°C)	Rainfall (mm)
---------------	--------------------------	--------------------------	---------------

Comment [A4]: The quantity of the irrigation water was controlled per grow bag? The description of the experiment was not written. What is the quantity of samples per species?

February	16.0	29.3	0.01
March	18.3	32.0	0.60
April	20.1	33.5	64.8
May	19.4	29.5	285.2
June	19.0	28.9	216.8
July	18.7	27.3	149.8
Mean	18.58	30.08	119.53

2.3 Photosynthesis measurements

Measurements of photosynthesis and allied parameters were made simultaneously using a portable photosynthesis system (CIRAS-3, PP-Systems, USA). Photosynthesis was measured on fully matured and well-exposed, at monthly intervals for a period of six months from February 2022 to July 2022. The measurements were made from 09:00 AM–12:00 PM in the second week of every month. Five to six observations per plant were recorded for each plant and the average is presented as the photosynthetic rate for a given month of a species. The other related parameters that were simultaneously recorded are stomatal conductance, transpiration rate, and water use efficiency.

Comment [A5]: Was it used one plant per species? The five to six repetitions were measured at different times or in different places of the plant?

2.4 Data analysis

The variation in the photosynthesis among the tree species and over time was analyzed based on the Analysis of Variance Technique (single-factor analysis) using the EXCEL-STAT statistical package.

Comment [A6]: This analysis without a mean test informs that there is at least one difference between treatments, it does not inform what they are. I suggest that, instead of tables with F results at the end, the analysis will be of trends and correlation, more comprehensive, including to present the results, as follows in the text.

3. RESULTS


A significant difference in photosynthetic rate was observed among the tree species as well as with time over months (Table 2). In the month of February *Grevillea robusta* was found to have the highest photosynthetic rate $29.18 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ followed by *Bauhinia purpurea* $27.92 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, *Pongamia pinnata* $22.62 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, *Peltophorum pterocarpum* $21.74 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, *Swietenia mahagoni* $19.94 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and the lowest photosynthetic rate was observed in *Spathodea campanulata* $16.15 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Table 2). Similarly, in the month of March and April *Bauhinia, purpurea* was found to have the highest photosynthetic rate whereas, *Swietenia mahagoni* and *Peltophorum pterocarpum* had the lowest photosynthetic rate. In the month of May, June, and July *Grevillea robusta* was found to have the highest photosynthetic rate although, *Peltophorum*

Comment [A7]: These comparisons should not be done because statistical comparison did not occur. These results already in table, no need to repeat them

pterocarpum showed the lowest photosynthetic rate in the month of May and *Spathodea campanulata* showed the lowest photosynthetic rate in the month of June and July. The photosynthetic ability of dominant tree species is found to be good.

Comment [A8]: This affirmation was done with what references of photosynthetic rates of tree species?

Formatted: English (United States)



Photosynthesis on six tree species in different seasons in the months of April and May (early monsoon season) all tree species showed a higher photosynthetic rate as compared to June and July (in February and March this happened too). *Grevillea robusta* was und?? to have the highest photosynthetic rate of 30.64 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in the month of May and the lowest photosynthetic rate was observed in the month of July at 30.64?? $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and no much difference was found in the month of summer (April) and winter (February) (Table 2). The same trend was followed in the rest of the species where the highest photosynthetic rate was observed in the early rainy season and the lowest was in July. The photosynthetic rates especially during the summer months (drought period) indicate the drought tolerance of these species, which is very significant considering the poor management practices these species provided.

Comment [A9]: This affirmation is wrong, the experiment was irrigated.

Table 2: Average monthly response of Photosynthesis rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) of six tree species

Species	Months						Mean
	February	March	April	May	June	July	
<i>Grevillea robusta</i>	29.18	27.58	29.18	30.64	24.60	21.50	27.11
<i>Spathodea campanulata</i>	16.15	18.66	23.66	22.47	14.28	14.50	18.37
<i>Swietenia mahagoni</i>	19.94	16.55	23.24	21.78	17.28	15.40	19.03
<i>Bauhinia purpurea</i>	27.92	28.08	29.30	29.13	22.44	21.11	26.28
<i>Pongamia pinnata</i>	22.62	21.26	25.81	24.57	22.18	19.15	22.58
<i>Peltophorum pterocarpum</i>	21.74	22.54	21.54	20.53	19.45	19.22	20.83
SEM	2.105	1.698	1.609	1.556	1.727	1.604	
CD (<0.05)	6.717	5.42	5.135	4.966	5.513	NS	

Note: NS - Non Significant

Table 3: Changes in the transpiration rates over time ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$) among six dominant tree species of urban landscapes.

Species	Months						Mean
	February	March	April	May	June	July	
<i>Grevillea robusta</i>	10.41	8.15	8.65	11.21	7.47	7.12	8.83
<i>Spathodea campanulata</i>	7.59	9.68	11.86	11.19	6.85	5.45	8.77
<i>Swietenia mahagoni</i>	7.17	4.55	8.12	9.21	5.97	5.65	6.77
<i>Bauhinia purpurea</i>	8.56	11.10	8.45	9.12	7.65	7.24	8.68

<i>Pongamia pinnata</i>	6.26	5.96	6.87	8.15	6.46	8.49	7.03
<i>Peltophorum pterocarpum</i>	7.98	6.45	9.15	7.10	9.92	7.26	7.97
SEM	0.799	0.867	0.615	0.738	0.676	0.349	
CD (<0.05)	NS	2.767	1.962	2.354	2.158	1.114	

Note: NS = Non Significant

The above results are reiterated from the transpiration rates (Table 3). There is a significant difference in the transpiration rate among the tree species except in the month of February. In the month of February *Grevillea robusta* recorded the highest transpiration rate at 10.41 mmol H₂O m⁻² s⁻¹ followed by *Bauhinia purpurea* at 8.56 mmol H₂O m⁻² s⁻¹, *Peltophorum pterocarpum* 7.98 mmol H₂O m⁻² s⁻¹, *Spathodea campanulata* 7.59 mmol H₂O m⁻² s⁻¹, *Swietenia mahagoni* 7.17 mmol H₂O m⁻² s⁻¹ and the lowest transpiration rate was observed in *Pongamia pinnata* 6.26 μmol H₂O m⁻² s⁻¹ (Table 3). Similarly in the month of March, *Bauhinia purpurea* and *Spathodea campanulata* had the highest transpiration rate and *Swietenia mahagoni* had the lowest transpiration rate. Similarly, *Spathodea campanulata* in the month of April, *Grevillea robusta* in the month of May, *Peltophorum pterocarpum* in June, and *Pongamia pinnata* in July showed the maximum transpiration rate. Likewise, *Pongamia pinnata* in the month of April and May, *Swietenia mahagoni* in June, and *Spathodea campanulata* in July showed the lowest transpiration rate. It can be observed that transpiration rates are complementary to the photosynthetic rates because both these functions are mutually exclusive (I think that the concept does not explain this affirmation. Mutually exclusive means independence among events. The opening stomata occur in photosynthesis and transpiration, no? Although with little data, an analysis to verify can be the correlation) [19].

Comment [A10]: The F was not significant. Why was this comparison made?

Formatted: Highlight

Pearson correlations among photosynthesis, transpiration, water efficient

Species	Photosy x Trans	Corr Pearson	
		Photosy x water effic	Trans x water effic
<i>Grevillea robusta</i>	0.851976217	0.493089	0.567483
<i>Spathodea campanulata</i>	0.971320381	0.975663	0.959073
<i>Swietenia mahagoni</i>	0.886871327	0.88639	0.875932
<i>Bauhinia purpurea</i>	0.659087038	0.441274	0.765072
<i>Pongamia pinnata</i>	-0.136066037	0.980682	0.002283
<i>Peltophorum pterocarpum</i>	-0.345227192	0.324642	-0.40456

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted Table





Table 3: Changes in the transpiration rates over time (mmol H₂O m⁻² s⁻¹) among six dominant tree species of urban landscapes.

Species	Months						Mean
	February	March	April	May	June	July	
<i>Gravillea robusta</i>	10.41	8.15	8.65	11.21	7.47	7.12	8.83
<i>Spathodea campanulata</i>	7.59	9.68	11.86	11.19	6.85	5.45	8.77
<i>Swietenia mahagoni</i>	7.17	4.55	8.12	9.21	5.97	5.65	6.77
<i>Bauhinia purpurea</i>	8.56	11.10	8.45	9.12	7.65	7.24	8.68
<i>Pongamia pinnata</i>	6.26	5.96	6.87	8.15	6.46	8.49	7.03
<i>Peltophorum pterocarpum</i>	7.98	6.45	9.15	7.10	9.92	7.26	7.97
SEM	0.799	0.867	0.615	0.738	0.676	0.349	
CD (<0.05)	NS	2.767	1.962	2.354	2.158	1.114	

Note: NS- Non Significant



[Continued... all observations above](#)

It is noticed that in the month of April (Late summer) and May (Early rainy season), all tree species had a higher transpiration rate compared to June and July except for *Pongamia pinnata*, which showed more transpiration rate in July. *Gravillea robusta* was found to have the highest transpiration rate of $11.21 \mu\text{mol H}_2\text{O m}^{-2} \text{s}^{-1}$ in the month of May and the lowest transpiration rate was observed in the month of July at $7.12 \mu\text{mol H}_2\text{O m}^{-2} \text{s}^{-1}$ and no much difference was found during summer months (Table 3). The same trend was followed in the rest of the species where the highest transpiration rate was observed in the early rainy season and the lowest was in the month of July.

~~Table 4: Average monthly response of Water Use Efficiency (WUE) $\text{mmol CO}_2/\text{mol H}_2\text{O}$ of six tree species~~

Species	Months						Mean
	February	March	April	May	June	July	
<i>Gravillea robusta</i>	2.50	2.42	2.50	3.86	3.24	2.00	2.67
<i>Spathodea campanulata</i>	2.27	3.64	4.70	4.19	2.26	2.29	3.22
<i>Swietenia mahagoni</i>	3.30	3.25	4.55	4.82	3.3	2.98	3.70
<i>Bauhinia purpurea</i>	2.05	5.09	2.53	5.08	2.07	2.77	3.26

<i>Pongamia pinnata</i>	3.66	2.96	5.51	5.26	3.93	2.24	3.92
<i>Peltophorum pterocarpum</i>	2.92	5.29	3.12	4.7	4.03	2.85	3.81
SEM	0.407	0.366	0.494	0.3	0.517	0.42	
CD (<0.05)	NS	1.169	1.578	NS	NS	NS	

Note: NS - Non Significant

A significant difference in the WUE was found in the month of March and April whereas, remaining months there was no difference among the tree species were noticed (Tabela 4). In the month of February, April and May *Pongamia pinnata* found to have the highest WUE likewise *Peltophorum pterocarpum* in the month of March and June, and *Swietenia mahagoni* in the month of July had a higher WUE (Table 4). *Spathodea campanulata* recorded lower WUE in the month of February, while *Grevillea robusta* in the month of March, April, and May, *Bauhinia purpurea* in the month of June, and *Pongamia pinnata* in the month of July had a lower WUE among all the species. The WUE in the month of April and May was found to be highest as compared to other months. Since photosynthesis and transpiration are mutually exclusive, WUE depends on the efficiency with which carbon is assimilated to a unit of transpiration.

Formatted: Highlight

Table 4: Average monthly response of Water Use Efficiency (WUE) mmol CO₂/mol H₂O of six tree species

Species	Months						Mean
	February	March	April	May	June	July	
<i>Grevillea robusta</i>	2.50	2.42	2.50	3.86	3.24	2.00	2.67
<i>Spathodea campanulata</i>	2.27	3.64	4.70	4.19	2.26	2.29	3.22
<i>Swietenia mahagoni</i>	3.30	3.25	4.55	4.82	3.3	2.98	3.70
<i>Bauhinia purpurea</i>	2.05	5.09	2.53	5.08	2.07	2.77	3.26
<i>Pongamia pinnata</i>	3.66	2.96	5.51	5.26	3.93	2.24	3.92

<u><i>Peltophorum pterocarpum</i></u>	<u>2.92</u>	<u>5.29</u>	<u>3.12</u>	<u>4.7</u>	<u>4.03</u>	<u>2.85</u>	<u>3.81</u>
SEM	0.407	0.366	0.494	0.3	0.517	0.42	
CD (<0.05)	NS	1.169	1.578	NS	NS	NS	

Note: NS- Non Significant

4. DISCUSSION

In the present study, it was found that six tree species growing in the temperatures range of 19.4 - 29.5 0C. temperatures Temperatures showed variation in the photosynthetic rates with a maximum rate recorded in the month of May (Table 1). The temperature range reported here is shown to be the optimal range for other tree species [24, 25, 26]. The temperatures around 40 0C are shown to reduce the photosynthetic rates critically because the process of photosynthesis is a chain of biochemical reactions in which many enzymes are involved. These enzymes are highly temperature sensitive and get denatured at higher temperatures. One of the studies found that plants growing at 8, 12, 14, 19, 23, and 30 0C showed similar gross leaf photosynthetic rates when measured at ambient temperatures [27]. When plants were incubated at temperatures 200C higher than ambient, the leaf photosynthetic rates increased to essentially the same value, irrespective of growth or measurement temperature. To better understand the effects of temperature on photosynthesis it is important to know the effect of temperature on the enzymes involved in photosynthesis. Enzymes may be affected a great deal by temperature. If the temperature is too low or too high the enzymes may move around too slowly to meet the substrate and for a reaction to occur. These findings suggest that temperature is an important factor in determining leaf photosynthetic potential in many terrestrial trees (all trees are terrestrial) [28, 29]. The different effects of temperature history on short-term photosynthesis in terrestrial species may be due to the lack of stomata and the rapid temperature equilibration of leaves with the surrounding environmental conditions.

Photosynthesis is particularly sensitive to moisture stress because the stomata will tend to close to conserve water as the soil water status decreases [30]. Indeed, stomatal regulation is vital in maintaining the hydraulic status of the plant [31]. In the present study during the month of May (late summer to early monsoon) all the tree species showed higher photosynthetic rates (Table 2). T-this is because, during the month of May, the study area received more rainfall as compared previous months (Table 1) as it alleviates the moisture

Comment [A11]: A temperatura não mostrou variação na taxa fotossintética, senão a correlação teria sido alta.

Comment [A12]: Introdução

Comment [A13]: Crescendo na mesma taxa fotossintética que temperatura ambiente de 20-25oC?

Formatted: Highlight

Comment [A14]: Não compreendido

Formatted: Highlight

Comment [A15]: O que o(s) autor(es) quiseram explicar?

stress the plant was under. So, from this study, it is also noticed that transpiration rate and WUE were found more in the month of May compared to other months (Table 3 & 4).

Here it used the average of the photosynthesis, transpiration, water efficient with minimal, maximal temperature, and rainfall.

	corr Pearson
<u>phot_aver x MinTemp</u>	<u>0.201035</u>
<u>phot_aver x Rainfall</u>	<u>0.733883</u>
<u>phot_aver x MaxTemp</u>	<u>-0.11138</u>
<u>trans_averx MinTemp</u>	<u>0.322865</u>
<u>trans_aver x Rainfall</u>	<u>0.511619</u>
<u>trans_averx MaxTemp</u>	<u>0.224934</u>
<u>Wateff_aver x MinTemp</u>	<u>0.529999</u>
<u>Wateff_aver x Rainfall</u>	<u>0.52509</u>
<u>wateff_aver x MaxTemp</u>	<u>0.349276</u>

Formatted: English (United States)

Formatted Table

A decrease in the net photosynthesis rate and transpiration is usually observed in plants under low rainfall conditions [32, 33, 34] and is also observed in environments where the evaporative demand is very high [35]. The present study observed that the tree species showed a photosynthetic depression at fewer rainfall events. According to a study, the depression of photosynthesis during dry days or non-rainy days probably results from too much light during this period, which can inhibit photosynthesis [23]. Other possible causes of depression at less rainfall include an increase in the atmospheric vapor pressure deficit [36, 37, 38] and high temperature [39, 40]. The closure of stomata during the event may also decrease the intercellular CO₂ concentration, inhibiting photosynthesis [34]. Thus, the effective regulation of the stomatal aperture is a critical plant functional attribute of plants for optimal development of plants.

The ratio between carbon assimilation and the **throughput** of water by transpiration results in water use efficiency [20]. The WUE, therefore, indicates the efficiency of carbon assimilation taking place to the unit amount of water lost in the transpiration. Most often WUE is attributed to a higher rate of photosynthesis [21] and not a lower transpiration rate [22].

Formatted: Highlight

Hence, **Photosynthesis, transpiration, and WUE are interdependent.** (Here ok) Extreme environmental conditions, such as high temperatures, rainfall, and drought, can affect

Formatted: Highlight

photosynthetic activity and WUE in plants [23]. Therefore, these or any one of these can be good indicators of a plant's performance under stressful conditions.

5. CONCLUSION (need revision after revision of all text)

This study provides an understanding of how photosynthetic activities in trees occur over time. Photosynthesis among the tree species was found to be highest in *Grevillia robusta* and in *Spathodea campanulate* and was least in *Bauhinia purpurea*. Among the six-tree species *Grevillea robusta* recorded the highest photosynthetic rates in pre and post-monsoon seasons too. The highest rate of photosynthesis was observed in the month of May among all the species during which plants experienced a minimum temperature of 19.4⁰C and a maximum of 29.5⁰C and received a rainfall of 285.2 mm after a prolonged dry spell of the previous months. Though the plants were watered and not subject to moisture stress during the experimental period the surrounding environment prevailed during the experimental period has created atmospheric stress for the plants. Transpiration most often changes in tune with photosynthesis. However, the WUE indicates the efficiency with which photosynthesis operates in relation to transpiration. **These findings suggest that all these species have good photosynthetic ability to sustain atmospheric stress and grow.** The presence of these tree species in more numbers across different landscapes of urban regions is also a **testimony to their pollution tolerance** in the urban ecosystem.

Formatted: Highlight

Formatted: Highlight

6. REFERENCE

1. IPCC (2007) Climate changes the physical science basis. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds) Contribution of Working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. 2007: Cambridge University Press, Cambridge.
2. Hikosaka K, Ishikawa K, Borjigidai A, Muller O, Onoda Y. Temperature acclimation of photosynthesis: mechanisms involved in the changes in the temperature dependence of photosynthetic rate. J Exp Bot. 2006; 57:291–302.

3. Sage RF, Kubien DS. The temperature response of C3 and C4 photosynthesis. *Plant Cell Environ.* 2007; 30:1086–1106.
4. Sage RF, Way DA, Kubien DS. Rubisco, Rubisco activase, and global climate change. *J Exp Bot.* 2008; 59:1581–1595.
5. Smith NG, Dukes JS. Plant respiration and photosynthesis in global-scale models: incorporating acclimation to temperature and CO₂. *Glob Change Biol.* 2013; 19:45–63.
6. Lichtenthaler, H.K. Vegetation stress: An introduction to the stress concept in plants. *J. Plant Physiol.* 1996; 148, 4–14.
7. Hofmann, G.E.; Todgham, A.E. Living in the now: Physiological mechanisms to tolerate a rapidly changing environment. *Annu. Rev. Genet.* 2010; 72, 127–145.
8. McCree, K.J. The measurement of photosynthetically active radiation. *Solar Energy* 1973; 15, 83–87.
9. McCree, K.J. Photosynthetically active radiation. In *Physiological Plant Ecology I*; Springer: Berlin/Heidelberg, Germany, 1981; pp. 4–155.
10. Kautsky, H. Hirsch, A. Neue versuche zur kohlen säureassimilation. *Naturwissenschaften* 1931; 19, 964
11. Kalaji, H.M.; Schansker, G.; Ladle, R.J.; Goltsev, V.; Bosa, K.; Allakhverdiev, S.; Brestic, M.; Bussotti, F.; Calatayud, A.; Dąbrowski, P.; et al. Frequently asked questions about in vivo chlorophyll fluorescence: Practical issues. *Photosynth. Res.* 2014; 122, 121–158.
12. Chukhutsina, V.U.; Holzwarth, A.R.; Croce, R. Time-resolved fluorescence measurements on leaves: Principles and recent developments. *Photosynth. Res.* 2019; 140, 355–369.
13. Farquhar GD, von Caemmerer S, Berry JA. A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta.* 1980; 149:78–90.
14. Gomez, L. F., Lopez, J. C., Riano Y, Mantoya, E. C., Diurnal changes in leaf gas exchange and validation of a mathematical model for coffee canopy photosynthesis. *Photosynthesis.*, 2005; 43(4): 575-582.
15. Zhang, D. Y., Wang, X. H., Chen, Y. And Xu, D. Q., Determinant of photosynthetic capacity in rice leaves under ambient air conditions. *Photosynthetica.*, 2005; 43: 273–276.
16. Ephrath, J. E., Timlin, D. J., Reddy, V. And Baker, J., Irrigation and elevated carbon dioxide effects on whole canopy photosynthesis and water use efficiency in cotton (*Gossypium hirsutum* L.). *PlantBiosyst.*, 2012; 145: 202–215.

17. Lindeman, Nishihara, G. N, Noro, T. And Terada, R., Effect of temperature and light on the photosynthesis as measured by chlorophyll fluorescence of cultured *Eucheuma denticulatum* and *Kappaphycus sp.* (Sumba strain) from Indonesia. J.Appl. Phycol., 2013; 25: 399– 406.
18. Watson, R. T., Noble, I. R., Bolin, B., Ravindranath, N. H., Verardo, J. D. And Dokken, D. J., Land Use, Land-Use Change, and Forestry, IPCC Special Report. Cambridge Univ. Press, Cambridge 2000; 388.
19. Farquhar, G.D., Models describing the kinetics of ribulose biphosphate carboxylase oxygenase. Arch. Biochem. Biophys. 1979; 193: 456–468.
20. Mortlock, M. Y., Transpiration and Water Use Efficiency. Office of Economic and Statistical Research., 2018; pp: 1250-1252.
21. Venter, W. O., Snyman, H. A., Van, R., And W. L. J., Photosynthetic response to water stresses in Themeda Iriandra and Era- growl is lehmanniana. S. Afr. J. Eor. 1997; 63: 37-41.
22. Agata, W., Kawamitsu. Y., Hakoyama. S. And Kaufman. P. B., Characteristics of photosynthesis, transpiration and water use efficiency of C3 and C4-grasses leaves, Kyoto, Japan. Prol. Int. Gnusland Collg., 1985; 347- 349.
23. Yu, D. J., Kim, S. J, Lee, H. J., Stomatal and non-stomatal limitations to photosynthesis in field-grown grapevine cultivars. Biol. Plant., 2009; 53: 133–137.
24. Biebl, R., McRoy, C.P., Plasmatic resistance and rate of respiration and photosynthesis of *Zostera marina* at different salinities and temperatures. Mar. 1971; Biol. 8, 48-56.
25. Drew, E.A., Physiological aspects of primary production in seagrasses. Aquat. Bot. 1979; 7, 139-150.
26. Bulthuis, D.A., Effects of in-situ light reduction on density and growth of the seagrass *Heterozostera Tasmania* in western Port Victoria, Australia. J. Exp. Mar. Biol. Ecol. 1983; 67, 91-103.
27. Evans, A.S., Webb, K.L., Penhale, P.A., Photosynthetic temperature acclimation in two coexisting seagrasses *Zostera marina* and *Ruppia maritima*. Aquat. Bot.1986; 24, 185-198.
28. Berry J, Bjorkman O. Photosynthetic response and adaptation to temperature in higher plants. Annu Rev Plant Physiol. 1980; 31:491– 543.
29. Sall T, Pettersson P. A model of photosynthetic acclimation as a special case of reaction norms. J Theor Biol, 1994; 166:1–8.

Formatted: Highlight

30. Lawlor, D. W. And Tezara, W., Causes of decreased photosynthetic rate and metabolic capacity in water-deficient leaf cells: a critical evaluation of mechanisms and integration of processes. *Ann. Bot.*, 2009; 103: 561–579.
31. Guo, X. Y., Zhang, X. S. And Huang, Z. Y., Drought tolerance in three hybrid poplar clones submitted to different watering regimes. *J. Plant Ecol.*, 2010; **3**: 79–87.
32. Endres, L., Daily and seasonal variation of water relationship in sugar apple (*Annona squamosa L.*) under different irrigation regimes at semi-arid Brazil. *Sci. Hortic.*, 2007;113: 149–154.
33. Hessini, K., Martinez, J. P., Gandour, M., Albouchi, A., Soltani, A. And Abdelly, C., Effect of water stress on growth, osmotic adjustment, cell wall elasticity and water-use efficiency in *Spartina alterniflora*. *Environ. Exp. Bot.*, 2009; 67: 312–319.
34. Saibo, N. J. N., Lourenc, O, T. And Oliveira, M. M., Transcription factors and regulation of photosynthetic and related metabolism under environmental stresses. *Ann. Bot.*, 2009; 103: 609–623.
35. Feng, Y. L, Cao, K. F., Photosynthesis and photoinhibition after night chilling in seedlings of two tropical tree species grown under three irradiances. *Photosynthetica.*, 2005; 43: 567–574.
36. Shirke, P. A. And Pathre, U.V., Influence of leaf to air vapor pressure deficit (VPD) on the biochemistry and physiology of photosynthesis in *Prosopis juliflora*. *J. Exp. Bot.*, 2004; 55: 2111–2120.
37. Ohsumi, A., Hamasaki, A., Nakagawa, H., Homma, K., Horie, T. And Shiraiwa, T., Response of leaf photosynthesis (*Oryza sativa L.*) varieties to vapor pressure difference in rice relation to stomatal and leaf internal conductance. *Plant Prod. Sci.*, 2008;11: 184–191.
38. Oliver, R. J., Finch, J. W. And Taylor, G., Second generation bioenergy crops and climate change: a review of the effects of elevated atmospheric CO₂ and drought on water use and the implications for yield. *GCB Bioenergy.*, 2009; 1: 97–114.
39. Day, M., Influence of temperature and leaf to air vapor pressure deficit on net photosynthesis and stomatal conductance in red spruce (*Picea rubens*). *Tree Physiol.*, 2000; 20: 57–63.
40. Eamus, D., Taylor, D. T., Macinnisng, C. M. O., Shanahan, S. And Silva, L., Comparing model predictions and experimental data for the response of stomatal conductance and guard cell turgor to manipulations of cuticular conductance, leaf-to-air vapor pressure difference, and temperature. *Plant, Cell Environ.* 2008; 31: 269–277.

Photosynthetic rate, transpiration rate, and water use efficiency of cotton canopy in oasis edge of Linze]

Formatted: Font: 24 pt, Bold, English (United States), Kern at 18 pt

[Article in Chinese]

Ting-Ting Xie¹, Pei-Xi Su, Song Gao

Formatted: English (United States)

Affiliations expand

- PMID: 20873616

Abstract

The measurement system of Li-8100 carbon flux and the modified assimilation chamber were used to study the photosynthetic characteristics of cotton (*Gossypium hirsutum* L.) canopy in the oasis edge region in middle reach of Heihe River Basin, mid Hexi Corridor of Gansu. At the experimental site, soil respiration and evaporation rates were significantly higher in late June than in early August, and the diurnal variation of canopy photosynthetic rate showed single-peak type. The photosynthetic rate was significantly higher ($P < 0.01$) in late June than in early August, with the daily average value being (43.11 ± 1.26) micromol CO₂ x m⁻² x s⁻¹ and (24.53 ± 0.60) micromol CO₂ x m⁻² x s⁻¹, respectively. The diurnal variation of canopy transpiration rate also presented single-peak type, with the daily average value in late June and early August being (3.10 ± 0.34) mmol H₂O x m⁻² x s⁻¹ and (1.60 ± 0.26) mmol H₂O x m⁻² x s⁻¹, respectively, and differed significantly ($P < 0.01$). The daily average value of canopy water use efficiency in late June and early August was (15.67 ± 1.77) mmol CO₂ x mol⁻¹ H₂O and (23.08 ± 5.54) mmol CO₂ x mol⁻¹ H₂O, respectively, but the difference was not significant ($P > 0.05$). Both in late June and in early August, the canopy photosynthetic rate was positively correlated with air temperature, PAR, and soil moisture content, suggesting that there was no midday depression of photosynthesis in the two periods. In August, the canopy photosynthetic rate and transpiration rate decreased significantly, because of the lower soil moisture content and leaf senescence, but the canopy water use efficiency had no significant decrease.

Formatted: Font: 12 pt, English (United States)

Formatted: English (United States)

UNDER PEER REVIEW

