

## Original Research Article

# Influence of Different Temperatures and Sexes on the Survival and Stress Responses of Nile Tilapia (*Oreochromis niloticus* L.)

### ABSTRACT

This study was conducted to assess the influence of different water temperature ranges and sexes on the stress response of Nile tilapia. Sixteen proactive males and 16 proactive female fish were used in the study. There were eight combinations of factors, and each combination was replicated four times. Changes in eye color pattern (ECP), ventilation rate (VR), and skin color of the fish were monitored after reaching the treatment temperature of the environment. Mortality of fish was also monitored. Results showed that the mean ECP of the fish was not affected by the interaction between water temperature and sex of fish. However, mean ECP was significantly affected by water temperature and sex of fish. Mean VR of the fish was affected by the interaction between water temperature and sex of fish. Furthermore, mean VR was significantly affected by water temperature and sex of fish. Decreased dissolved oxygen (DO) was observed in increasing water temperature which may have also contributed to the stress response of Nile tilapia besides the increasing water temperature. Two out of four fish in T4M died after 6-12 hours of exposure while in T4F three out of four fish died after 2-4 hours. There was a decrease in the final weight of fish in all combinations of factors. However, there were no significant differences in the initial and final weights and lengths of fish in all combinations of factors. On body color, both male and female fish exposed to increasing water temperatures exhibited an intense body color (presence of vertical stripes). From the results of this study, male fish had significantly lower mean ECP scores and mean VR values than female fish when exposed to different ranges of water temperature. Likewise, male fish exposed to high temperatures (39-40°C) survived longer than female fish. Therefore, it can be concluded that male fish are more tolerant to increasing water temperature up to 37°C than female fish since males can easily cope with stress. The ECP, VR, and body color are good indicators that can be used in assessing the level of stress of Nile tilapia.

*Keywords: eye color pattern, ventilation rate, stress responses, Nile tilapia*

### 1. INTRODUCTION

"Tilapia" is the generic name of a group of cichlids endemic to Africa [1]. They are second only to carp as the most cultured and widely farmed fish species [1,2]. Tilapia possess a large variety of adaptation responses to match the vast array of its environment [3]. Among the tilapiines, the Nile tilapia (*Oreochromis niloticus*) is the most important cultured species [2]. The Nile tilapia has a potential for aquaculture due to its rapid growth at warm temperatures [1]; ability to adapt to high-density confinement and various systems of farming

Comment [WU1]: Too monotonous way of speaking

Comment [WU2]: add some of the results of this research study that are easily understood by the reader

[4]; high resistance to diseases, viruses, and parasites than other cultured fish [5]; tolerance to wide range of water salinities [6] and high tolerance to poor water and crowding [7].

Temperature is one of the major environmental factors influencing reproduction and a key modulator of early gonadal ontogenesis in fish [8]. Other factors are photoperiod, density, sex ratio, size and weight of breeders, food, stress, oxygen, salinity, and pH [3]. The rise of water temperature due to climate change puts additional stress on freshwater ecosystems [9].

The majority of Asia's tropical and subtropical regions account for over 65 percent of the world's aquaculture production, which is produced inland. Global warming's effects on the climate are probably not going to have much of an impact on aquaculture techniques. Contrarily, climate change will affect the availability of water, and weather patterns like severe rain that may lead to eutrophication, and stratification in static (lentic) waters [10].

Temperature is one of the environmental factors with the greatest influence on the growth performance of animals [11]. Fish is affected by the temperature of the surrounding water which influences the body temperature, growth rate, food consumption, feed conversion, and other body functions [12,13]. Growth and livability in fish are optimum within a defined temperature range [14]. This study assessed the fish response to different water temperatures above the ideal range for the growth of the fish. The optimum temperature range in order for the fish to perform its best is 27-32°C [11]. The morphological changes of Nile tilapia to different ranges of temperature will increase our knowledge of how the fish reacts and copes with a given temperature.

Comment [WU3]: good review

## 2. MATERIAL AND METHODS

Comment [WU4]: made more systematic

### 2.1 Experimental Fish

Thirty-two pieces of male and 32 pieces of female Nile tilapia, with an individual weight of about 25 grams were used in this study. The male and female Nile tilapia were obtained from the Freshwater Aquaculture Center, Central Luzon State University, Science City of Muñoz, Nueva Ecija. Freshwater Aquaculture Center Selected Tilapia (FaST) strain was used as the strain of Nile tilapia. Both sexes of Nile tilapia were similar in weight, length, and age so there are no phenotypic and genetic variations that may affect the growth of the fish. The fish were individually weighed and measured for their length.

### 2.2 Experimental Treatments and Design

There were two factors that were evaluated on their influence on fish mortality and stress responses. These factors were: Factor A - temperature ranges (30-31°C; 33-34°C; 36-37°C and 39-40°C) and Factor B - sexes of fish (male and female). Each combination of factors is shown in Table 1. The study used a 4 x 2 factorial experiment using randomized complete block design (RCBD). Each factor combination had four replicates.

Table 1. Experimental treatments

Factor A (Temperatures, °C)	Factor B (Sexes)	
	Male (M)	Female (F)
30 – 31 (T1)	T1M	T1F
33 – 34 (T2)	T2M	T2F
36 – 37 (T3)	T3M	T3F
39 – 40 (T4)	T4M	T4F

## 2.3 Experimental Unit

The fish underwent conditioning for seven days in the aquarium prior to the study being conducted. The experimental set-up was composed of thirty-two (32) units of glass aquaria measuring 30 x 30 x 30 cm. An electric heater was installed in each aquarium to attain constant treatment temperature throughout the experimental period. An aeration system was provided in each aquarium to provide sufficient dissolved oxygen (DO) for the fish. A glass thermometer was used to monitor the water temperature. DO meter was used to monitor dissolved oxygen in the water. One fish was stocked in each aquarium.

## 2.4 Experimental Procedures

Thirty-two male and 32 female fish were isolated individually in glass aquaria for five days. Three sides of glass aquaria were covered with white paper to avoid the fish from seeing each other during the isolation period. The eye color pattern (ECP) of the fish was monitored daily. Sixteen males and 16 females were selected based on their ECP. Fish with a mean ECP score of less than 4 during the isolation period were selected.

Selected fish were transferred from the isolation set-up to the experimental set-up. The water temperature in each aquarium of the experimental set-up was gradually adjusted to the treatment temperature within 8 hours to avoid thermal shock to the fish. Changes in ECP, ventilation rate (VR), and body color of the experimental fish were monitored at 0, 15, 30, 45 min; 1, 2, 6, 12, 24, 48, 72, 96, and 120 h after reaching the treatment temperature of the environment. Mortality was also observed in the 7-day experimental period. Feed was given once per day at 1% of the body weight [15].

## 2.5 Monitoring of the Eye Color Pattern and Ventilation Rate

The ECP was monitored daily during the entire duration of the experiment. The ECP was quantified as the darkened area of both the iris and sclera. The circular area of the eye is divided into 8 equal parts using 4 imaginary diameter lines (Figure 1)[15,16]. The ECP values observed ranged from zero (no darkening) to eight (total darkening).




Figure 1. Quantification of ECP

The VR was visually estimated by counting the time (i.e. seconds) for 20 successive opercular or buccal movements [17]. It was measured daily at three times per fish in a period of 3 consecutive minutes. The readings were averaged for each fish and were used in analyzing the relationship between the ECP and VR.

## 2.6 Statistical Analysis

The relationship between ECP and VR was assessed using linear regression and Pearson correlation coefficient. Data were analyzed using two-way Analysis of Variance and the treatment means were compared using Duncan's Multiple Range Test. The statistical analyses were performed using SPSS software version 23 for Windows.

### 3. RESULTS AND DISCUSSION

#### 3.1 Eye Color Pattern

The mean ECP and mean VR of male and female fish subjected to different temperature ranges for 7 days are shown in Table 2. The mean ECP of the fish was not affected by the interaction between water temperature and sex of fish ( $P=0.495$ ). However, mean ECP was significantly affected by water temperature ( $P<0.01$ ) and sex of fish ( $P<0.01$ ). The mean ECP of female fish at 39-40°C (T4F) of  $5.81\pm0.84$  and of male fish at 39-40°C (T4M) of  $5.41\pm0.85$  were statistically comparable to each other but were significantly higher than ( $P<0.05$ ) those in the rest of the treatments except that those in T4M and female fish in 36-37°C (T3F) ( $4.83\pm2.02$ ). The mean ECP of male fish at 36-37°C (T3M) ( $3.70\pm1.94$ ), female fish at 33-34°C (T2F) ( $3.40\pm2.04$ ), male fish at 33-34°C (T2M) ( $3.04\pm1.71$ ) were not significantly different from each other but were significantly higher ( $P<0.05$ ) than those in female fish in 30-31°C (T1F) with  $2.02\pm1.86$  and male fish in 30-31°C (T1M) with  $1.44\pm1.60$ . The mean ECP in the latter two treatments was not significantly different from each other.

In this study, it was observed that the ECP of male and female fish changed from pale to dark after reaching the treatment temperature ranges. ECP is a stress-coping style in Nile tilapia, and it is a good, easy, and inexpensive tool for assessing both alertness and stress levels in the fish [15].

Comment [WU5]: the discussion is very good

**Table 2. Mean eye color pattern ( $\pm$ SD) of male and female fish per treatment temperature**

SEX	Mean ECP of fish per Treatment Temperature			
	T1	T2	T3	T4
Male	$1.44\pm1.60^{ax}$	$3.04\pm1.71^{ay}$	$3.70\pm1.94^{by}$	$5.41\pm0.85^{bz}$
Female	$2.02\pm1.86^{aw}$	$3.40\pm2.04^{ax}$	$4.83\pm2.02^{ay}$	$5.81\pm0.84^{az}$

**Note:** Means having a common superscript within a column (a, b), or within a row (w, x, y, z) are not significantly different from each other at 5% ( $P<0.05$ ) probability level by DMRT. T1 = 30-31 °C; T2 = 33-34 °C; T3 = 36-37 °C; T4 = 39-40 °C.




Figure 2. Mean eye color pattern of all fish per combination of factors

Figure 2 shows the daily ECP values of male and female fish during the 7-day experimental period. It was observed that there was a decreasing trend of ECP values in all combinations of factors except T4M and T4F. It may be because fish subjected to different ranges of water temperatures (T1, T2, and T3) were able to gradually cope in the stressful environment. However, fish in T4M and T4F exhibited darker ECP throughout the study because of very high temperature of 39-40°C. It was observed that fish in T4M and T4F refused to eat the food given to them and dedicated their energy to escaping the stressful environment by swimming aggressively and trying to jump out of the water. Similar results were also reported in red tilapia (*Oreochromis* spp.) possibly due to the need to maintain their energy consumption during stressful conditions [18].

It also shows that males can more easily cope with stress than females when subjected to different ranges of water temperature. Male fish had lower mean ECP scores when exposed to different ranges of water temperature than female fish. According to Figueiredo-Fernandes et al. [19], male fish subjected to higher temperature exhibits significantly higher Glutathione S-transferase (GST) activity than females. Apart from its essential functions in different intracellular activities, a crucial role of GST is the defense against stress [19].

Similar studies were conducted wherein ECP was used as an indicator of stress. Decano et al. [20], subjected red tilapia to high temperature and confinement stress. They found that an increased value of ECP was recorded in an increased temperature and confinement stress. Apino et al. [21], assessed changes in the ECP of red tilapia exposed to blue and red-light emitting diode (LED). Results of the study showed that darker ECP was observed in fish exposed to a blue and red LED. Likewise, Bernandino et al. [22] used red tilapia as experimental fish where they subject the fish to electric shock and handling stress. Darker ECP was also observed in this study. The stress response of red tilapia to high ammonia levels was also reported [23]. Again, darker ECP was observed in fish subjected to high ammonia levels.

Comment [WU6]: poor explanation development

### 3.2 Ventilation Rate

Table 3 shows the mean ventilation rate of male and female fish per treatment temperature. It was observed that the mean VR of the fish was affected by the interaction between water temperature and sex of fish ( $P < 0.01$ ). Furthermore, mean VR was significantly affected by water temperature ( $P < 0.01$ ) and sex of fish ( $P < 0.01$ ). Female fish in T4 had the highest mean VR of  $2.87 \pm 0.70$  buccal movements/sec which was significantly different in all combinations of factors ( $P < 0.05$ ). The mean VR of T4M ( $2.04 \pm 0.44$  buccal movements/sec) and T3F ( $2.12 \pm 0.76$  buccal movements/sec) were statistically comparable but both were significantly higher than those of the rest of the remaining treatments. Also, there was no significant difference among T3M ( $1.54 \pm 0.46$  buccal movements/sec), T2F ( $1.61 \pm 0.57$  buccal movements/sec), and T2M ( $1.39 \pm 0.40$  buccal movements/sec) but they were significantly higher ( $P < 0.05$ ) than those in female fish in T1F with  $1.07 \pm 0.26$  buccal movements/sec and male fish in T1M with  $1.02 \pm 0.25$  buccal movements/sec. The VR in the latter two combinations of factors were not significantly different from each other.

Comment [WU7]: add relevant explanations

**Table 3. Mean ventilation rate ( $\pm$ SD) of male and female fish per treatment temperature**

SEX	Mean VR of fish per Treatment Temperature			
	T1	T2	T3	T4
Male	$1.02 \pm 0.25^{ax}$	$1.39 \pm 0.40^{ay}$	$1.54 \pm 0.46^{by}$	$2.04 \pm 0.44^{bz}$
Female	$1.07 \pm 0.26^{aw}$	$1.61 \pm 0.57^{ax}$	$2.12 \pm 0.76^{ay}$	$2.87 \pm 0.70^{az}$

**Note:** Means having a common superscript within a column (a, b), or within a row (w, x, y, z) are not significantly different from each other at 5% ( $P < 0.05$ ) probability level by DMRT. T1 = 30-31 °C; T2 = 33-34 °C; T3 = 36-37 °C; T4 = 39-40 °C.




Figure 3. Ventilation rate of all fish per combination of factors

Figure 3 shows the daily VR values (buccal movement/sec) of males and females during the experimental period of 7 days. Results showed that there was an abrupt change in VR values when the temperature of water increased. It was observed that there was a decreasing trend of VR in all combinations of factors indicating that the fish can cope with increasing temperature. However, T4F had the highest significant mean which is statistically different ( $P < 0.05$ ) compared to the other combination of factors. It was observed that female fish exposed to 39-40°C died 2-4 hours after exposure while male fish died after 6-12 hours to the same range of water temperature. Ventilation rate is a sympathetic response (a response that indicates that someone is in a bad situation) that quickly increases in response to a stressor, thus it is particularly useful for measuring stress levels because it is easily quantifiable [15,24].

Similar studies [20,21,22,23] that used ECP as an indicator of stress also used VR to evaluate the stress responses of fish. Red tilapia subjected to high temperature and confinement stress, blue and red LED, electric shock and handling stress, and high ammonia levels had faster VR.

### 3.3 Relationship between Eye Color Pattern (ECP) and Ventilation Rate (VR)

The relationship between the mean VR and mean ECP of Nile tilapia is shown in Figure 4. It was observed that there was a significant strong positive correlation between the mean VR and mean ECP ( $n=8$ ;  $r=0.810$ ;  $P < 0.01$ ) of Nile tilapia when exposed to thermal stress. This indicates that as the VR increases, the ECP also increases.

**Comment [WU8]:** it's very good, the writing hopes to be more consistent and in accordance with existing scientific principles




Figure 4. Relationship between mean ventilation rate and mean eye color pattern of fish (n=8;  $r=0.810$ ;  $P<0.01$ )

### 3.4 Dissolved Oxygen (DO) and Temperature

Table 4 presents the mean DO and temperature in all combinations of factors during the 7-day experimental period. T1M and T1F were significantly different ( $P<0.05$ ) compared to those in the rest of the combination of factors. However, T1M with a mean DO of  $5.91\pm0.18$  mg/L, and T1F with a mean DO of  $5.92\pm0.28$  mg/L were statistically comparable. T2M, T2F, and T3M had a mean DO of  $5.36\pm0.09$  mg/L,  $5.34\pm0.04$  mg/L, and  $5.19\pm0.15$  mg/L, respectively, which were significantly different in all combinations of factors. Yet, T2M, T2F, and T3M were statistically comparable. T3F had a mean DO of  $4.94\pm0.19$  mg/L. T4M and T4F had statistically comparable means of  $4.40\pm0.21$  mg/L and  $4.40\pm0.06$  mg/L, respectively. T4M had the lowest mean DO in this study.

It was observed in this study that the DO concentration decreases as the temperature increases. The relationship between water temperature and DO is inverse; that is, cold water is able to "hold" more dissolved oxygen than warm water [25]. Decreased DO concentration could be predicted by increasing water temperature [26]. It was observed that DO in T4M and T4F was below the required DO concentration, which may contribute to the observed fish mortalities. Since water temperature affects the oxygen solubility in the water, it affects the behavioral responses and morphological changes of fish [28]. Thus, the high range of water temperature and low DO concentration imposed stress on the fish [32].

Comment [WU9]: it's very good, the writing hopes to be more consistent and in accordance with existing scientific principles

**Table 4. Mean dissolved oxygen (mg/L) and temperature (°C)**

Combination of factors	Mean ( $\pm$ SD)	
	Temperature	Dissolved Oxygen
T1M	$30.54\pm0.06^a$	$5.91\pm0.18^a$
T1F	$30.48\pm0.09^a$	$5.92\pm0.28^a$
T2M	$33.51\pm0.08^b$	$5.36\pm0.09^b$
T2F	$33.48\pm0.09^b$	$5.34\pm0.04^b$
T3M	$36.41\pm0.08^c$	$5.19\pm0.15^b$
T3F	$36.47\pm0.17^c$	$4.94\pm0.19^c$

T4M	39.55±0.10 <sup>d</sup>	4.40±0.21 <sup>d</sup>
T4F	39.57±0.18 <sup>d</sup>	4.40±0.06 <sup>d</sup>

**Note:** Means in a column superscripted with different letters are significantly different at 5% ( $P < 0.05$ ); T1 = 30-31 °C; T2 = 33-34 °C; T3 = 36-37 °C; T4 = 39-40 °C; M= males; F = females

### 3.5 Mortality

There were no mortalities in T1M, T1F, T2M, T2F, T3M, and T3F in all replicates. Since all fish used in this study were proactive (fish with ECP values of less than 4 after the isolation period), they can easily adapt to the stressful environment. However, in T4M, two out of four male fish subjected to the water temperature of 39-40°C died after 6-12 hours of exposure. In T4F, three out of four female fish died after 2-4 hours of exposure to the water temperature range of 39-40°C. Factors affecting these results can be the stress experienced by the fish due to the high-temperature ranges and low concentration of DO.

**Comment [WU10]:** explanation is quite good, but it needs innovation in how to synthesize

### 3.6 Weight and Length

On initial weight, T1M with a mean weight of 25.50±2.33 g was statistically comparable to T1F with 27.28±3.65 g, T2M with 25.93±2.49 g, T2F with 23.60±1.01 g, T3M with 27.38±2.45 g, T3F with 27.80±2.42 g, T4M with 25.03±5.10 g and T4F with 24.88±4.31 g. After the study, T2F had the lowest final mean weight of 22.95±0.98 g followed by T4F and T4M with a final mean weight of 23.53±3.45 g and 23.48±4.81 g, respectively. T1F had the highest final mean weight of 27.05±3.84 g, followed by T3M with 27.03±2.41 g. All combinations of factors were not significantly different from each other.

**Comment [WU11]:** explanation is quite good, but it needs innovation in how to synthesize and describe scientifically

On initial length, T1F had a mean length of 11.73±0.49 cm, followed by T3F with 11.65±0.40 cm, T3M with 11.50±0.41 cm, T1M with 11.50±0.45 cm, T2M with 11.43±0.30 cm, T4F with 11.20±0.54 cm, T2F with 11.10±0.27 cm and lastly T4M with 11.10±0.42 cm. On final length, T2M had the lowest final mean length with 11.43±0.30 cm. While T1F and T3M had the highest final mean length with 11.75±0.50 cm and 11.75±0.50, respectively. Yet, all combinations of factors were not significantly different from each other.

**Table 5. Changes in the final weight and length of fish in all combinations of factors**

Combination of factors	Final Weight	Final Length
T1M	-	*
T1F	-	+
T2M	-	*
T2F	-	+
T3M	-	+
T3F	-	+
T4M	-	+
T4F	-	+

**Note:** + = increase; - = decrease; \* = no changes

Table 5 presents the changes in the final weight and length of fish in all combinations of factors. It was observed that there was a decrease in the final weight of fish in all combinations of factors. Also, some fish had an increase while some had no changes in the final length. A possible factor affecting the results is the physical stress experienced by fish due to low dissolved oxygen concentration and high-temperature range [31]. Exposure to any type of stressors either environmental, chemical, or perceived stressors results in a series of physiological responses in animals, like growth depensation in fish [28].

### 3.7 Body color

Both male and female fish exposed to increasing water temperatures exhibited an intense body color (presence of vertical stripes) (Figure 5). It was observed that as the temperature was increased higher than the ideal range, the body color of the fish started to intensify from pale to grey. Color alteration is controlled by the brain through the pigment called chromatophore. Cichlid species, including the Nile tilapia, change their skin color when stressed and this was also observed in red Nile tilapia [29][30].




Figure 5. Fish exposed to 30-31°C (left) and 39-40°C (right)

### 4. CONCLUSION

Male fish had a significantly lower mean ECP score, mean VR values, and less intensified skin coloration than female fish when exposed to increasing water temperature. Likewise, male fish exposed to high temperatures (39-40°C) survived longer than female fish. Therefore, it only shows that the males can more easily cope with stress than females when subjected to different ranges of water temperature. These behavioral (VR) and morphological (ECP and body color) responses are good, easy, and inexpensive tools for assessing both alertness and stress levels in the fish subjected to thermal stress.

Comment [WU12]: good conclusion

### REFERENCES

1. Popma, T., Masser M. Tilapia life history and biology. Southern Regional Aquaculture Center. 1999; 4 p.
2. Ariadi, H., Fadjar, M., Mahmudi, M. The relationships between water quality parameters and the growth rate of white shrimp (*Litopenaeus vannamei*) in intensive ponds. Aquaculture, Aquarium, Conservation & Legislation. 2019;12(6): 2103-2116.
3. Baroiller, J.F., Toguyeni A. The tilapiini tribe: environmental and social aspects of reproduction and growth. Fisheries and Aquaculture. 1997;3, 1-10.
4. Rakocy, J.E. Tank culture of tilapia. Southern Regional Aquaculture Center. 1989; 4 p.
5. Popma, T.J., Lovshin LL. Worldwide prospects for commercial production of tilapia. 1996. Accessed 28 March 2023. Available: <https://aurora.auburn.edu/handle/11200/4157>
6. El-Sayed, A-FM. Tilapia culture in salt water: environmental requirements, nutritional implications and economic potentials. p. 95-106. In: L. Elizabeth Cruz Suarez, D. Ricque Marie, M.T. Salazar, M.G. Nieto Lopez, D.A. Villareal Covazos, A.C. Puello Cruz y Armando Garcia Ortega (Eds.). Advances on Aquaculture Nutrition. VIII Simposium International de

Nutricion Acuicola, Universidad Autonoma de Nuevo Leon Monterrey, Nuevo Leon Mexico. 2006.

7. Wurts, WA. Tilapia: a potential species for Kentucky fish farms. Kentucky Fish Farming. 1999. Accessed 28 March 2023. Available: <http://www2.ca.uky.edu/wkrec/tilapiakentucky.htm>

8. Strüssmann, CA, Nakamura, M. 2002. Morphology, endocrinology, and environmental modulation of gonadal sex differentiation in teleost fishes. Fish Physiology and Biochemistry. 2002;26:13–29.

9. Ariadi, H, Wafi, A. Water Quality Relationship with FCR Value in Intensive Shrimp Culture of Vannamei (*Litopenaeus vannamei*). Samakia: Jurnal Ilmu Perikanan. 2019;11(1): 44–50.

10. De Silva, SS, Soto, D. Climate change and aquaculture: potential impacts, adaptation and mitigation. p. 151-212. In: K. Cochrane, C. De Young, D. Soto and T. Bahri (eds.). Climate Change Implications for Fisheries and Aquaculture: Overview of Current Scientific Knowledge. FAO Fisheries and Aquaculture Technical Paper No. 530. Rome, Italy. 2009.

11. Ariadi, H, Fadjar, M, Mahmudi, M. Financial feasibility analysis of shrimp vannamei (*Litopenaeus vannamei*) culture in intensive aquaculture system with low salinity. ECSOFiM (Economic and Social of Fisheries and Marine Journal). 2019;7(1): 95-108.

12. Ariadi, H, Mahmudi, M., Fadjar, M. Correlation between density of vibrio bacteria with *Oscillatoria* sp. abundance on intensive *Litopenaeus vannamei* shrimp ponds. Research Journal of Life Science. 2019;6(2): 114-129.

13. Azevedo, PA, Cho, CY, Leeson, S, Bureau, DP. Effects of feeding level and water temperature on growth, nutrient and energy utilization and waste outputs of rainbow trout (*Oncorhynchus mykiss*). Aquat. Living Resour. 1998;11: 227-238.

14. Gadomski, DM, Caddell, SM. Effects of temperature on early life history stages of California halibut *Paralichthys californicus*. Fishery Bulletin. 1991;89: 567-576.

15. Vera Cruz, EM, Tauli MP. Eye color pattern during isolation indicates Stress-Coping Style in Nile Tilapia *Oreochromis niloticus* L. International Journal of Scientific Research in Knowledge. 2015;3(7): 181-186.

16. Volpato, GL, Luchiari, AC, Duarte, CRA, Barreto RE, Ramanzini, GC. Eye color as an indicator of social rank in the fish Nile tilapia. Braz. J. Med. Biol. Res. 2003; 36: 1659–1663.

17. Alvarenga, CMD, Volpato, GL. Agonistic profile and metabolism in alevins of the Nile tilapia. Physiology and Behavior. 1995;57: 75-80.

18. Ariadi, H, Mujtahidah, T. Analisis Permodelan Dinamis Kelimpahan Bakteri *Vibrio* sp. Pada Budidaya Udang Vaname, *Litopenaeus vannamei*. Jurnal Riset Akuakultur. 2022;16(4): 255-262.

19. Figueiredo-Fernandes, A, Fontainhas-Fernandes A, Peixoto, F, Rocha, E, Reis-Henriques, MA. Effects of gender and temperature on oxidative stress enzymes in Nile tilapia *Oreochromis niloticus* exposed to paraquat. Pesticide Biochemistry and Physiology. 2006;85: 97-103.

20. Wafi, A, Ariadi, H, Muqsith, A, Mahmudi, M, Fadjar, M. Oxygen consumption of *Litopenaeus vannamei* in intensive ponds based on the dynamic modeling system. *Journal of Aquaculture and Fish Health*. 2021; 10(1):17-24.
21. Apino, RM, Templonuevo, RMC, Vera Cruz, EM. Stress Responses of Red Tilapia (*Oreochromis* spp.) Exposed to Blue and Red-Light Emitting Diode (Led). *Egyptian Academic Journal of Biological Sciences, B. Zoology*. 2022;14(1): 159-167.
22. Bernardino, JT, Templonuevo, RMC, Vera, EM. Responses of red tilapia (*Oreochromis* spp.) subjected to electric shock and handling stress. *International Journal of Fisheries and Aquatic Studies*. 2020; 8(1): 287-290.
23. Constantino, RV, Templonuevo, RMC, Fajardo, LJ. Stress responses of red tilapia (*Oreochromis* spp.) to high ammonia levels. *International Journal of Fisheries and Aquatic Studies*. 2019;7(6): 89-93.
24. Barreto, RE, Volpato GL. Caution for using ventilator frequency as an indicator of stress in fish. *Behav. Process*. 2004;66: 43–51.
25. Wafi, A, Ariadi, H. Estimasi Daya Listrik Untuk Produksi Oksigen Oleh Kincir Air Selama Periode "Blind Feeding" Budidaya Udang Vaname (*Litopenaeus vannamei*). *Saintek Perikanan: Indonesian Journal of Fisheries Science and Technology*. 2022;18(1): 19-35.
26. Clipperton, GK, Koning, CW, Locke, AG, Mahoney, JM, Quazi, B. Instream flow needs determinations for the South Saskatchewan River basin, Alberta, Canada. *Alberta Environment*. 2003. Accessed 28 March 2023. Available: <https://open.alberta.ca/publications/0778530450>
27. Moyle, PB, Cech, J. J. *Fishes: An Introduction to Ichthyology*. Upper Saddle River: Prentice Hall. 2003. Accessed 28 March 2023. Available: [https://books.google.com.ph/books/about/Fishes.html?id=sZYWAQAIAAJ&redir\\_esc=y](https://books.google.com.ph/books/about/Fishes.html?id=sZYWAQAIAAJ&redir_esc=y)
28. Vera Cruz, EM, Brown, CL. 2007. The influence of social status on the rate of growth, eye color pattern and insulin-like growth factor-I gene expression in Nile tilapia, *Oreochromis niloticus*. *Hormones and Behavior*. 2007;51: 611-619.
29. Templonuevo, RMC, Vera Cruz EM. Responses of red Nile tilapia (*Oreochromis niloticus* L.) subjected to social and confinement stresses. *The CLSU International Journal of Science & Technology*. 2016;1(2): 7-14.
30. Ariadi, H, Khristanto, A, Soeprapto, H, Kumalasari, D, Sihombing, J.L. Plankton and its potential utilization for climate resilient fish culture. *Aquaculture, Aquarium, Conservation & Legislation*. 2022;15(4): 2041-2051.
31. Ariadi, H, Mujtahidah, T, Hidayati, S. Pelaksanaan Tradisi Petik Laut Nelayan Hindu dan Islam Dalam Korelasi Pengelolaan Sumber Daya Pesisir di Jembrana. *Jurnal Kebijakan Sosial Ekonomi Kelautan dan Perikanan*. 2022;12(2): 137-144.
32. Wafi, A, Ariadi, H, Muqsith, A, Madusari, B.D. Business feasibility of intensive vaname shrimp (*Litopenaeus vannamei*) with non-partial system. *ECOSOFIM (Economic and Social of Fisheries and Marine Journal)*. 2021;8(2): 226-238