

Original Research Article

Profiling the Contamination level and pollution load index of borehole water from Esako West LGA, Edo State, Nigeria

Abstract

Fresh water has continued to decrease both in quality and quantity with a wide variety of pollutants entering into fresh water resources. In this research the quality of fresh water was being accessed using standard method of analysis of water. Physicochemical parameters and heavy metal content were determined and data was assessed using contamination factors, degree of contamination and pollution load index result revealed that borehole water at Esako west LGA was slightly contaminated by phosphate (0.16mg/l) and over 5 times the WHO standard. The acidic water (pH6) was observed over 200mg/l of carbon dioxide the contamination factor for zinc were generally greater than 1, while other parameter had values of contamination factors less than 1. In conclusion, borehole water at Esako west contaminated and slightly polluted through the locals continued to consume this water, there is a need for treatment to remove phosphate whose source may be from phosphate rocks in the area or sipping from phosphate fertilizer in farms.

Key words: Anthropogenic, Chemometrics, Index Models Ground water, Pollutants, Physicochemical, Water quality.

Introduction

Water is one of the most abundant gifts of nature. It is an indispensable resource for the continued existence of all living things including man and adequate supply of fresh and clean drinking water is a basic need for all human beings¹. In nature, all water contains impurities; as water flows in streams, accumulates in lakes and filters through layers of soil and rock in the ground, it dissolves or absorbs substances it come in contact with, which may be harmful or harmless². One of the major and critical problems in most developing countries today is the provision of an adequate and safe drinking water to its populace³. Drinking water that is safe and aesthetically acceptable is a matter of high priority to National Agency for Foods and Drugs Administration and Control (NAFDAC) and other regulatory agencies in Nigeria and is expected to meet the Nigerian Industrial Standard. Furthermore, drinking water that is fit for human consumption is expected to meet the World Health Organization and be free from physical and chemical substances and microorganisms in an amount that can be hazardous to health⁴. It is a known fact that no single method of purification can eliminate 100% contaminants from drinking water. However, water can be and should be made safe for consumption within acceptable limits⁵.

Sachet water is any commercially treated water, manufactured, packaged and distributed for sale in sealed food grade containers and is intended for human consumption while bore hole water is a non-treated water that is readily consumed by humans. Water consumers are frequently unaware of the potential health risks associated with exposure to water borne contaminants which have often led to diseases like diarrhea, cholera, dysentery, typhoid fever, legionnaire's disease and parasitic diseases⁶. The continuous increase in the sale and indiscriminate consumption of both bore hole and packaged drinking water in Nigeria is of public health significance, as the prevalence of water related diseases in developing countries is determined by the quality of their drinking water^{7,8}.

The safety of drinking water in poor and deprived communities has in the last decade been in jeopardy as a result of the introduction of refuse and sewage into sources of water supply. The intake of unwholesome water could have devastating effects on our health as unsafe drinking water is a key determinant of many microbial diseases with serious complications in immune-competent and immune-compromised individuals. The introduction of sachet water was aimed at providing safe, hygienic and affordable instant drinking water to the public and to curb the magnitude of water related infections in the country^{9-116,14}.

Pure water is colorless, odorless and tasteless with high boiling and melting points as well as high heat of vaporization. Pure water can be slightly ionized reversibly to yield hydrogen and hydroxyl ions. Therefore, water is not just a solvent in which the chemical reactions of the living cell occur. It is very considered often in direct participation in those reactions. Quality of drinking water is evaluated on the basis of its chemical components. This is done by assessing the pH, hardness, total alkalinity, dissolved oxygen, carbon dioxide, heavy metals and organic constituents^[14].

Consumption of sachet water in Nigeria is on the increase irrespective of whether they have NAFDAC Certification or not. However, despite the strong effort by NAFDAC in the regulation and quality assessment of sachet water, there are a growing number of reported public illnesses after drinking sachet water. There are a number of reported cases of typhoid, diarrhea and other water borne diseases arising from consumption of sachet water¹⁵. Sachet water serves as a major source of potable/drinking water to communities around Imo State.

Borehole water is readily available water these days for human consumption, and the quality and source of the water are questionable. Anthropogenic activities such as the use of pesticides, fertilizers and the release of contaminated waste material could pollute the water aquifer. This borehole water is consumed by humans with the believe that it is coming directly from its source, hence it is clean. This research therefore was conducted to evaluate the quality of borehole water samples collected from Esako West LGA in Edo State.

Some of these bore hole and sachet water are contaminated, either chemically or microbial, and are toxic to man. When these contaminants accumulate beyond the recognized and recommended limits, they become toxic to living organism. The consumption or use of contaminated bore hole or poorly treated sachet water is capable of causing water or chemical related diseases. It is on these bases that an attempt is being made to determine the quality of both treated and untreated water in Esako west area of Edo State¹⁹.

The aim of this research work was to carry out quality assessment of bore hole water in Esako West area of Edo State. The specific objectives for achieving the aim were: to analyze borehole and water samples collected from Esako West area of Edo State LGA; to compare the result of the analysis with those of WHO acceptable standard for drinking quality water; to evaluate contamination factors and pollution load. This research work was limited to the following analysis; pH, conductivity, total dissolved solid, chloride ion, phosphate, nitrate, copper, iron, zinc, manganese and copper and CO₂ concentrations.

2.0 MATERIAL AND METHODS

PREPARATION OF REAGENT

1 gram of phenolphthalein salt was weighed with a weighing balance and dissolved in 100ml of ethanol to obtain 1% phenolphthalein indicator.

0.02N sodium hydroxide solution was prepared by dissolving 0.8 grams of sodium hydroxide pellet in 1000ml of distilled water in a 1000ml volumetric flask.

SAMPLE COLLECTION

The water samples used for this research were collected from Esakowest Local government area of Edo State Nigeria according to standard methods for water analysis.

2.2.1 DETERMINATION OF pH

The pH values of the water samples were determined by the method of national agency for food and drug, administration and control (NAFDAC), using (Standard operating procedure for water analysis). 200ml beaker was washed, rinsed dried in an oven at 105°C and allowed to cool in a desiccator. 100ml of the water sample was introduced into the beaker and a pH meter of the model JENWAY 3510 pH meter previously calibrated with buffer 7.0 is used to determine the pH of the water samples.

2.2.2 DETERMINATION OF CONDUCTIVITY

The method of NAFDAC for determination of pH of water samples was followed and the conductivity was determined using Lasany microprocessor conductivity meter.

2.2.3 DETERMINATION OF TOTAL DISSOLVED SOLID

The total dissolved solid for the water samples was estimated from the values obtained from the conductivity using a mathematical manipulation as shown below

$$1000\mu\text{s}/\text{cm} = 500\text{mg}/\text{L}$$

2.2.4 DETERMINATION OF THE CO₂ CONTENT The carbon dioxide content of the water samples was determined by the method of NAFDAC as stated in the standard operating procedure for water analysis. The maximum acceptable limit for NAFDAC is 50mg/L. 100ml of the water samples was measured using a measuring cylinder and introduced into a previously washed, rinsed and oven dried conical flask. 2-3 drops of phenolphthalein indicator was added to the sample and the resulting mixture was titrated using 0.02N sodium hydroxide solution until a faint permanent pink colour was observed. Agitation must be done to avoid dissolving more carbon dioxide in the samples from the atmosphere.

The titre value is recorded and the concentration of the CO₂ in the water samples is calculated using the relation 1ml of 0.02NaOH = 10mg/L of CO₂

2.2.5 MINERAL ANALYSIS OF WATER SAMPLE

The mineral analysis of water samples was determined by the method used and described by Verla et al.,2022 using Atomic Absorption Spectrometer. The concentration of each of the minerals was obtained in mg/L following the procedures described by the instrument manual.⁴¹

RESULTS

The results from 8 different borehole water samples collected in Esako West LGA of Edo state are as shown in the tables below.

Table 1: Results of physicochemical analysis of the bore hole water samples.

S/ N	Parameter	S1	S2	S3	S4	S5	S6	S7	S8	M	SD	Range	WHO
a	PO ₄ ³⁻ (mg/l)	0.08	0.10	0.18	0.05	0.21	0.20	0.24	0.12	0.15	0.07	0.08-0.24	0.03
b	NO ₃ ⁻ (mg/l)	29	7	59	32	55	41	12	05	30	20.94	7-59	50
c	Cl(mg/l)	4	3	9	10	5	2	3	0.10	4.51	3.40	2-10	5
d	Fe (mg/l)	0.10	0.12	0.41	0.21	0.27	0.09	0.12	0.13	0.18	0.11	0.09-0.41	0.3

E	Zn (mg/l)	3.5	2	4	0.57	3.2	1.10	2.1	0.96	2.18	1.27	0.57-3.5	2
F	Mn(mg/l)	0.02	0.03	0.01	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.01-0.03	0.4
g	Cu (mg/l)	0.02	0.001	0.80	0.03	0.01	0.02	0.01	0.01	0.11	0.27	0.001-0.80	3
h	Ph	6.08	6.83	5.19	6.29	4.36	6.40	5.97	5.99	5.89	0.77	4.36-6.83	6.5- 9.0
i	EC (µS/cm)	27.75	29.50	124	23.5	105	34.55	20.30	18.10	47.83	41.7	18.1-104.5	100
j	TDS (mg/l)	13.87	14.65	276	20.0	256	17.27	17.15	17.05	79.02	115.61	13.87-276	250
11	CO ₂ mg/L	280	175	125	75	270	116	200	153	174.25	72.83	75-280	50

Contamination and Pollution Index Models: The C_f is an indicator used to assess the presence and intensity of anthropogenic contaminants in groundwater. The C_f was determined mathematically using equation 1

$$C_F = \frac{C_m}{C_b} \quad 1$$

Where C_m is the concentration of a particular parameter in the water and C_b is the reference concentration of that parameter. WHO standards for drinking water were taken as the reference concentrations. The degree of contamination degree (CD), is simply the summation of contamination factors and is employed in comparing sites

$$CD = \sum C_F \quad 2$$

The PLI is a potent tool that provides a simple and comparative means for assessing the level of pollution. The PLI gives a summative indication of the overall level of toxicity in a sample. The PLI value greater than 1 is polluted, less than 1 indicates no pollution whereas values equal to 1 indicates contaminant loads close to the reference concentration. The pollution load index was determined mathematically using Equation 2.

Generally, PLI is estimated using the following equation;


$$PLI = [Cf_1 \times Cf_2 \times Cf_3 \times Cf_4 \dots \dots \dots Cf_n]^{1/n} \quad 3$$

where n is the number of parameters considered in the study and Cf_n is the contamination factor for each individual parameter. This PLI provides a comparative means for assessing a site quality.


Table 2: CONTAMINATION FACTOR

Parameters	1	2	3	4	5	6	7	8	ΣCF
PO ₄ ³⁻	2.66	3.33	6.00	1.66	7.00	6.66	8	4.00	39.31
NO ₃ ⁻	0.58	0.14	1.18	0.64	1.10	0.82	0.24	0.10	4.8
Cl ⁻	0.80	0.60	1.80	2.00	1.00	0.40	0.60	0.02	7.22
Fe	0.33	0.024	1.36	0.70	0.90	0.30	0.40	0.03	4.044


Zn	1.75	1.00	2.00	0.28	1.60	0.55	1.05	0.48	8.71
Mn	0.05	0.08	0.03	0.05	0.03	0.03	0.03	0.05	0.35
Cu	0.01	0.001	0.04	0.02	0.01	0.01	0.01	0.01	0.111
Ph	0.42	1.05	0.03	0.97	0.67	0.92	0.92	0.92	5.9
EC	0.28	0.30	1.24	0.24	1.05	0.35	0.20	0.18	3.84
TDS	0.06	0.06	1.10	0.08	1.03	0.07	0.07	0.07	2.54
CO ₂	5.65	3.5	2.5	1.5	5.4	2.32	4.0	3.06	27.93
PLI	1.09	0.89	1.57	0.74	1.80	1.13	1.41	0.81	
ΣCF	12.59	10.085	17.28	8.14	19.79	12.43	15.52	8.92	




a




b




c




d





e




f




g h



i



j



k

Fig. a to k: Charts for Comparisons of mean values of physicochemical parameters with NAFDAC and WHO standard values and Standard deviation

4.1 DISCUSSION

Table 1 shows the results obtained for the physicochemical analysis and statistical evaluations of bore hole water samples collected from Esako west local Government area of Edo State.

Phosphate: The presence and concentration of phosphorus in waters could be an important pollutant. Phosphates are retained by some complex system of biological uptake, absorption, and mineralization³⁴. Phosphates are not toxic to people or animals unless present in very high concentrations in which case they interfere with digestion. In this study, the phosphate level in the range was 0.08-0.24 mg/L. These levels of phosphates were not within the permissible limit of NAFDAC and WHO for all the samples. Hence, the bore hole water samples were contaminated with about five times the WHO standard. Sources of phosphates could be adjoining farmlands in which phosphate fertilizers have been used.

Nitrate: Values of nitrate range was in the range of 7.0-59.0 mg/L while mean nitrate levels (30 mg/l) obtained in this study were less than WHO standard of 50 mg/l. Nitrate concentrations in samples 3 (59.0 mg/L) and 5 (55.0 mg/L), were slightly higher than threshold values for both NAFDAC and WHO standards. Usually, high nitrate levels are attributed to anthropogenic activities such as the indiscriminate dumping of waste³¹. Nitrates are known to interfere with the oxygen transport ability of the red blood cells with this effect felt more in the popular blue baby illness. Infants who drink water high in nitrates may turn “bluish” and appear to have difficulty in breathing since their bodies are not receiving enough oxygen³².

Chloride (Cl): Chloride is invariably present in small amounts in almost all natural waters and its contents go up appreciably with increasing salinity³³. High concentration of chlorides is considered an indicator of pollution due to organic wastes of animal or industrial origin. Chlorides are troublesome in water and also harmful to aquatic life. The chlorine content of the bore hole water samples range is within 2.0-10.0 mg/L and were generally lower than the standard limit of WHO and NAFDAC except for sample 3, 4 and 5 that showed very high concentration with values of 9.0 mg/L, 10.0 mg/L and 5.0 mg/L. It has no adverse health impact, but excess of it impacts bad taste to the drinking water³⁴.

Heavy metals studied here include iron, zinc, manganese and copper. These were selected because of their link to cancer and organ damage^{35, 36}. These metals react with oxygen to form reactive oxygen species that damage cells³⁷.

Iron (Fe): Iron is an important mineral element required by human in trace quantity. It is one of the most abundant elements in the soil, hence its concentration in ground water abounds. High concentration of iron in drinking water is deleterious due to its health hazard³⁷, because excessive ingestion can cause anemia and even death in chronic exposure. In this study, the concentration of iron the water samples are below permissible limit of NAFDAC and WHO and the range obtained in this study were 0.09-0.41 mg/L although sample 3 gave a higher value of 0.4 mg/L.

Zinc (Zn): The Zn concentrations in the studied bore hole water samples had mean values range of 0.57-3.5mg/L. The observed values were reportedly higher than the common world range for WHO standard. Environmental contamination of Zn is mainly related to anthropogenic activities. The anthropogenic sources of Zn are related to industries and the use of liquid manure, composted materials and agrochemicals such as fertilizers and pesticides in agriculture³⁸.

Manganese (Mn): The sources of manganese in water may include metal alloys, batteries, glass and ceramic materials. This is not unusual for disposal of waste on water channels that finds their way into streams, especially wasted from auto mechanic workshops. The result obtained for the bore hole water samples showed that manganese was within the range of 0.01-0.03 mg/L. This range of values were within the permissible limit of the WHO and NAFDAC.

Copper (Cu): Ingestion of metals such as copper (Cu) may pose great risks to human health. It may also cause small increases in blood pressure, particularly in middle-aged and older people and can cause anemia. Contribution of Cu may be envisaged from dumping of solid wastes, application of fungicides, live stock manures, sludges and atmospheric deposition³⁷. The presence of Copper in the water samples were within the range of 0.001-0.80 mg/L and below the WHO permissible limit of 3 mg/L.

pH: The pH of water is an important parameter that determines the quality of drinking water. It is the measure of the acidity and alkalinity of the water^{37, 38}. The pH of quality water according to national agency for food and drug, administration and control (NAFDAC) and world health organization (WHO) standard should be within the range of 6.8-9.0. Hence below or above this limit, the water is termed unfit for human consumption. In this study, it was revealed that the pH of the majority of the water samples were below NAFDAC and WHO limit with. The results obtained were in the range of 4.36-6.88.

Electrical Conductivity (EC): Mean value of electrical conductivity of borehole water was 47.83, much lower than 100 the standard value for NAFDAC and WHO. It is known that acidic pH encourages high EC, but this was not so for borehole water at Esako west. EC is a measure of dissolved salts and can affect taste of water^{40, 41}.

Total dissolved solids: Total dissolved solids of the water samples range from 13.87-276mg/L in which sample 1,2,4,6-8 were within the acceptable range of WHO and NAFDAC while samples 3 and 5 were above the limit with values of 256 and 276 mg/L respectively. However, sample 3 has the highest TDS with 276 mg/L and sample 1 have the lowest with 13.87 mg/L. TDS may cause water to be corrosive and the consumption of water with high dissolved solids could lead to gastrointestinal diseases. These solids which can be suspended can cause damage to human if the concentration is too high³⁹.

Carbon dioxide content (CO₂): Carbon dioxide content in water is a very important parameter for defining water quality. The source of carbon dioxide in water is usually from the atmosphere. High concentration of carbon dioxide in water will increase the CO₂ content of the human when

ingested constantly. In this study, the carbon dioxide content of the water sample exceeded the recommended limit for quality water set by NAFDAC having a range of 75-280mg/L

Contamination and Pollution Indices

Table 2 shows the results obtained for the contamination factor. The results obtained for the contamination factors shows moderate contamination of metals. The results also showed that there is a very high contamination of phosphate in the water samples collected from the area. This pollution could be attributed to the continues uses of phosphate containing materials like NPK fertilizers which increases the level of phosphate in the soil which could percolate into the soil or be washed away by runoff during rainfall into water bodies.

CONCLUSION

It can be concluded that high phosphate and CO₂ content in the bore hole water made the water not fit for consumption. The overall results showed that the bore hole water samples collected from Esako west LGA of Edo State are considerably contaminated and unhealthy for human consumption without any form of treatment. The quality of available water has implication on the health status of a community. Polluted water is the major reason for the spread of many endemic diseases.

REFERENCES

1. W.H.O (2012) Guidelines for drinking-water quality. Geneva, Switzerland; 2012.
2. Mgbemena, N.M and Okwunodulu ,F.U.(2015). Physicochemical and microbiological assessment of borehole waters in Umudike, Ikwuano LGA, Abia State, Nigeria. *AdvApplSci Res.* 6(4):210–214.
3. Moyo, N.A.G.(2013). An analysis of the chemical and microbiological quality of ground water from boreholes and shallow wells in Zimbabwe. *PhysChem Earth, Parts A/B/C.* 66:27–32.
4. Iyasele, J.U. and Idiata, D.J. (2012). Determining the borehole water quality in Edo south and Edo north areas of Edo state. *Res J EngAppl Sci.* 2012;1(4):209–213.
5. Akpoveta, O.V. Okoh B.E. Osakwe, S.A.(2011). Quality assessment of borehole water used in the vicinities of Benin, Edo State and Agbor, Delta State of Nigeria. *Curr Res Chem.*;3(1):62–69.
6. Tamungang, N.E.B. Alakeh, M.N. Niba F.L.M. and Sunjo J. (2016). Physicochemical and bacteriological quality assessment of the Bambui community drinking water in the North West Region of Cameroon. *Afr J Environ Sci Technol.* 10(6):181–91.
7. Ugbaja V. and Otokunefor T. (2015). Bacteriological and Physicochemical Analysis of Groundwater in Selected Communities in Obio Akpor, Rivers State, Nigeria. *Br Microbiol Res J.* 7(5):235–42.

8. Collins, N.A. and Merapelo, D.M. (2011). Detection of Enterococcus species in groundwater from some rural communities in the Mmabatho area, South Africa: A risk analysis. *Afr J Microbiol Res.* 2011;**5**(23):3930–3935.
9. Isa, A. (2013). Physicochemical and bacteriological analyses of drinking water from wash boreholes in Maiduguri Metropolis, Borno State, Nigeria. *Afr J Food Sci.* **7**(1):9–13.
10. Ikeme, C.H. Dioha, I.J. Olasusi K.A. and Chukwu, P.U.(2014). Physico-Chemical Analysis of Selected Borehole Water In Umuhi, Town Imo State, Nigeria. *IntJ SciEng Res.* **5**:680–9.
11. John, A.A. Fehintolu, F.A. and Asabe, C.L. (2010). Assessment Of water quality from hand dug wells In Samaru – Zaria. *Nigeria J. Chem. Society, Nigeria, Zaria Chapter*, 3:42-44. DOI 10.4409.
12. Langegger, O. (1994). Groundwater quality and hand pump corrosion in West Africa. UNDP-World Bank water and sanitation program Report series. No. 8 Washington D.C., U.S.A. 158pp.
13. Mustapha, S, and Adamu, E.A. (1991). Discussion on Water Problems in Nigeria: Focus on Bauchi State. *National Res. Inst.*
14. American Public Health Association (2005). *Standard Methods for the Examination of Water and Wastewater.* Washington, DC, USA; 42.
15. WHO (2011). *Guidelines for Drinking Water Quality.* World Health Organization, Geneva.
16. Ademorati, C.M.A. (1996). *Environmental chemistry and toxicology,* Foludex press Ltd Ibadan, Nigeria p. 218.
17. American Public Health (1985). American Water Works Association (AWWA) and Water Pollution Control Federation (WPCF): *Standard Methods for the Examination of Water and Waste Waters* 16th edition, American Public Health Association Inc. New York, p.4.
18. American Public Health Association, APHA (1999). *Standard Methods for Examination of Water and waste water* 20th Ed. P. 5.
20. American Society for Testing and Material, ASTM (2004) *Annual Book of ASTM Standard, water and environmental technology, volume 11.01, Water (i)* ASTM International, West Conshohocken, PA, pp. 352-371.
21. Borne, R.A. (1978). *The Chemistry of our Environment,* J Wiley & Sons, Inc., NY . 2nd Ed. Pp. 187-189.
22. Chandra, S, Singh, A. and Tomar, P. K.(2012) Assessment of water quality values in Porur Lake Chennai, HussainSagar Hyderabad and Vihar Lake Mumbai, India. *Chem. Sci. Trans. 1:* 508-515.
23. Edward, E.J (1980) *Practical Information on surface Water Problems. The Chemistry of Water* Bulletin NO. 1237, Johnson Water screens Overseas Ltd, Northern Island. P.16

24. Erah P.O, Akujieze C.N, and Oteze G.E (2002). A quality of ground water in Benin City: A baseline study on inorganic chemicals and microbial contaminants of health importance in boreholes and open wells. *Trop. J. Pharm. Res.* 1(2): 75-82.
25. Golterman, H.I. (1978); *Methods for Physical and Chemical Analysis of Fresh Waters*, Billing and Sons Ltd. U.S.A. 2nd Ed. P.25
26. Hammer, M. J. and Kenneth M. A. (1981); *Hydrology and Quality of Water Resources*, John Wiley and Sons Inc. New York, U.S.A 2nd Ed. P.92
27. Hutton, M. (1987); *Human Health Concerns of Lead, Mercury and Arsenic*, Hutchinson, E.C. and Meama, K. M., Eds. *Lead, Mercury, Cadmium and Arsenic in the Environment*; Wiley scope, 2nd Ed. Pp. 85-94.
28. Kirk-Othmer (1964); *Encyclopedia of Chemical Technology*, John Wiley and Sons Inc. New York, U.S.A. p. 26.
29. Lind, J.E. (1959); *Water Pollution*, Clarendon Press, Oxford, p. 9.
30. Obi C.N, Okocha C.O (2007) Microbiological and Physico-Chemical of Selected Bore-hole Waters in World Bank Housing Estate, Umuahia, Abia State, Nigeria. *J. Eng. Applied Sci.* 2(5):920-929.
31. Okonkwo I.O, Ogunjobi A.A, Kolawale O.O, Babatunde S, Oluwole I, Ogunnusi T.A, Adejuyi O.D, and Fajobi E.A (2009). Comparative Studies and Microbial Risk Assessment of a Water Samples Used for Processing Frozen Sea foods in Ijora- Olopa, Lagos State, Nigeria. *EJEAFCh.* 8(6): 408-415.
32. Olowe O.A, Ojuronbe O, Opaleye O.O, Adedosu O.T, Oluwe R.A, and Eniola K.I.T (2005) Bacteriological Quality of Water Samples in Osogbo Metropolis. *Afr. J. Clin. Exper. Microbiol.* 6(3): 219-222.
33. World Health Organization., (2004), *Guidelines for drinking water recommendation*, Geneva, Switzerland Vol.1
<http://www.who.int/publications/2004/9241546387pdf>. Assessed 6/09/2021.
34. World Health Organization (2009). *Copper in drinking water. Background document for development of WHO guidelines for drinking water quality*. www.who.int/water/health/.../copper.pdf assessed 19/09/2018.
35. Magda M, Usalam A.B.D, Gaber I and Ahu Z, (2015) Impacts of landfill leachates on ground water quality: A case study of Egypt. *Journal of advanced research* 6(4) 579-586.
36. FAO/WHO (1988). *Requirement of Vitamin A, Iron, Folate and Vitamin B12. Report of a joint FAO/WHO Expert Consultation Rome*, Food and Agricultural Organization of the United Nations (FAO Food and Nutrition series No. 23).

37. Yusuf, Y.O., Jimoh, A.I., Onaolapo, E.O. and Dabo, Y(2015). An assessment of sachet water quality in Zaria Area of Kaduna State, Nigeria. *Journal of Geography and Regional Planning*, Vol. 8(7), pp. 174-180.
38. Osibanjo O, Ajayi S, Adebyi F, and Akinyanju P (2000) Public analysis reporting System as Applied to Environmental issues IPAN News, a Publication of the Institute of Public Analysts 1(3):10.
39. Mustapha S, and Adamu E.A, (1991). Discussion on Water Problems in Nigeria: Focus on Bauchi State. *National Res. Inst. P.* 8.
40. NAFDAC (1993) Standard operating procedure for water analyses. 1-30
41. Verla Andrew Wirnkor, Dikeocha Sandra Uzoamaka, Verla Evelyn Ngozi, Muna Stella Ebele, Njoku-Obi Treasure Njideka (2022) Determining water quality for drinking and irrigation of surface water in Oguta Lake, *International Journal of Advanced Chemistry Research* Volume 4 (1): 20-27. www.chemistryjournals.net.