

Original Research Article
**Analysis of Genetic Variability, Correlation and
Path for Yield and its Attributing Traits in Wheat
(*Triticum aestivum* L.)**

UNDER PEER REVIEW

ABSTRACT

The current study was undertaken to analyze the genetic variability, correlation and path for yield and its attributing traits at agriculture research farm, Department of Genetics and Plant Breeding, School of Agriculture, Lovely Professional University, Punjab during during *Rabi* 2021-22. Augmented design with F_2 seeds planted in ear to row fashion in 4 rows were used to study variability, correlation and path including 2 parent and its 6 F_{2s} for the 9 traits *viz.* Plant height, Spike length, Number of tillers per plant, Number of spikes per plant, Number of spikelets per ear, Number of Grains per ear, Biological yield, Harvest index and Grain yield. Finding revealed that Grain yield revealed positive and highly significant correlation with number of tillers per plant ($r_p= 0.8387$, $r_g= 0.8709$) and biological yield ($r_p= 0.9671$, $r_g= 0.9965$) while positive and significant correlation with Number of spikes per plant ($r_p= 0.7168$, $r_g= 0.7253$) at both phenotypic and genotypic level. Biological yield per plant revealed highest positive direct effect (1.0642, 0.8700) with significant association with grain yield (0.9965, 0.9671) whereas Number of tillers reveals least positive direct effect on grain yield (0.0183, 0.0834) at both genotypic and phenotypic level. PCV was higher than GCV for all the traits indicated an influence of environment over the traits. Number of spikelets per ear recorded highest PCV and GCV followed by Grain yield. Heritability (h^2_b) found higher for grain yield (84.2) followed by Number of spikelets per ear (83.4), Number of Grains per ear (82.7) and Plant height (80.1). Genetic advance (GA) recorded higher Number of spikelets per ear (35.07) followed by Grain yield (11.93). Moderate GA recorded for Plant Height (7.67) and Biological yield (6.88) whereas Spike length (0.62) recorded lowest GA. Hence, selection for these characters could bring improvement in yield and yield components.

Keywords: Correlation, Path, Phenotypic, Genotypic, Direct and indirect effect, grain yield.

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) having chromosome number $2n=6x=42$, [AABBDD] (hexaploid) is a widely cultivated, self-pollinating annual plant belonging to the true grass (Gramineae) family. According to [1], wheat holds great significance as one of the primary cereal crops as a staple food source worldwide. In terms of global food crops, wheat ranks second after rice [2]. Wheat plays a crucial role in providing approximately 20% of the calories and protein consumed by people worldwide, contributing to the sustenance of around 40% of the global population.

According to the Food and Agriculture Organization FAO [3], global wheat production reached approximately 776.5 mt in 2020. The production was estimated to slightly increase to 778.3 mt in 2021 and then decrease to 770.3 mt in 2022. Wheat holds a crucial position not only as a key crop for food security but also as a valuable cash crop on a global scale [4]. The importance of wheat is reflected in the significant global trade volume. In the 2020-21 season, world wheat trade reached a record high of 186.6 mt, surpassing the previous season by 1.2 percent or 2.3 mt FAO [5]. This highlights the significant role of wheat in meeting global food demands and economic activities.

In order to establish a successful breeding program, the presence of genetic diversity within the population is crucial. [6] confirm that plants with higher genetic variability are more likely to benefit from favourable hybrid crosses and generate productive recombinants, thus leading to a broader heritability within the population

Comment [IP1]: Mention traits

Comment [IP2]: Mention proper key words that represent your manuscript

Comment [IP3]: Use , delete having

Comment [IP4]: delete

Comment [IP5]: belongs

Comment [IP6]: delete

Comment [IP7]: rearrange the sentence and correct it

Comment [IP8]: rewrite in a scientific way

Comment [IP9]: give appropriate number

during the process of genetic improvement. Estimation of heritability in combination with genetic advance are usually more useful than heritability alone for forecasting genetic gain under selection. However, a trait with a high heritability does not always have a high genetic advance [7]. Correlation reveals the degree and direction of association at phenotypic and genotypic levels btw the yield and its contributing traits. However, it should be noted that the correlation could sometimes fail to give accurate insights into the individual impact of each character on the dependent character. It is understandable that a path analysis would be necessary to determine which characters actually affect seed yield. So path analysis is used to measure the indirect and direct effects of traits [8]. It is impossible to begin an effective breeding programme without first establishing genetic variability using appropriate metrics including GCV and PCV, Genetic advance (GA) and heritability (h^2_b) [9].

Comment [IP10]: rewrite it a proper scientific way

Comment [IP11]: Rewrite in a proper way

2. MATERIALS AND METHODS

The present investigation was carried out at the research farm, Department of Genetics and Plant Breeding, Lovely Professional University, Phagwara, Punjab during *Rabi* 2021-22. The F_2 seeds were space planted in ear to row fashion to study variability, correlation and path including 2 parent and its 6 F_{2s} for the 9 traits viz. Plant height, Spike length, Number of tillers per plant, Number of spikes per plant, Number of spikelets per ear, Number of Grains per ear, Biological yield, Harvest index and Grain yield. The experiment material consists of crosses obtained from matting of female line GS/2020-21/8049 (provided by Borlaug Institute for South Asia, BISA-CIMMYT) which are heat tolerant and susceptible to rust, with the PBW 766 male parent, which are newly released local cultivars. The list of wheat cultivars (along with Pedigree) used for study tabulated in (Table 1).

Comment [IP12]: Mention properly how and where did you prepare F1s, F2s and also give the number of years of study.

Table 1. Parents and crosses of wheat used in the experiment

S.No	Genotype	Parentage
1	GS/2020-21/8049 (Female)	WBLL1*2/BRAMBLING//SAAR/2*WAXWING/4/PBW343*2/KUKUNA//KRO NSTAD F2004/3/PBW343*2/KUKUNA/5/2*MUCUY
2	PBW 766 (Male)	PBW 766 NAC/TH.AC//3*PVN/3/MIRLO/BUC/4/2*PASTOR/5/KACHU/6/KACHU
S.No	Crosses	F_2 Population of GS/2020-21/8049X PBW 766
1	C1	(GS/2020-21/8049 X PBW 766) 1
2	C2	(GS/2020-21/8049 X PBW 766) 2
3	C3	(GS/2020-21/8049 X PBW 766) 3
4	C4	(GS/2020-21/8049 X PBW 766) 4
5	C5	(GS/2020-21/8049 X PBW 766) 5
6	C6	(GS/2020-21/8049 X PBW 766)6

Using a Spacing 22.5 x 15 cm (row to row 22.5 and plant-to-plant 15 cm) in augmented Design for F2 generation. The mean value was then determined for analysis over three replications Figure in (Table 2). The formulas provided by [7] were used to compute the different genetic factors viz. GCV, PCV, Heritability (h^2_b) and Genetic advance (GA). Using the formula provided by [10] for correlation co-efficient and [11] for path co-efficient analysis were obtained.

2.1 Estimating of correlation

Now, genotypic and phenotypic correlation coefficients were calculated using formula

$$\text{Phenotypic correlation (r}_p\text{)} = \frac{PCov_{xy}}{\sqrt{PV_x \cdot PV_y}}$$

$$\text{Genotypic correlation (r}_g\text{)} = \frac{GCov_{xy}}{\sqrt{GV_x \cdot GV_y}}$$

$$r_{xy} = \frac{Cov(x, y)}{\sqrt{V(x)} \times \sqrt{V(y)}}$$

Where,

r_{xy} = Correlation coefficient between character x and y

$Cov_{x,y}$ = Co-variance of character x and y

V_x = Variance of character x, and

V_y = Variance of character y

r_p = Phenotypic correlation

r_g = Genotypic correlation.

2.2 Path Analysis

Path analysis splits the correlation coefficient into the measures of direct and indirect effects and measures contribution of each independent variable on the dependent variable and estimates residual effects. It helps in determining the yield and yield contributing characters.

To estimate various direct and indirect effects, the following equations were used

$$r_{1y} = P_{1y} + r_{12}P_{2y} + r_{13}P_{3y} + \dots + r_{1l}P_{ly}$$

$$r_{2y} = r_{2y}P_{1y} + P_{2y} + r_{23}P_{3y} + \dots + r_{2l}P_{ly}$$

$$r_{ly} = r_{l1}P_{1y} + r_{l2}P_{2y} + r_{l3}P_{3y} + \dots + P_{ly}$$

Where,

r_{1y} to r_{ly} = Coefficient of correlation between factor 1 to l and dependent character y

r_{12} to $r_{l-1,l}$ = Coefficient of correlation among causal factors themselves

P_{1y} to P_{ly} = Direct effects of characters 1 to l on character y.

2.3 Residual effect

Residual effect, which measures the contribution of the characters was obtained by

$$(PRV) = \sqrt{1 - R^2}$$

Where,

$$R^2 = \frac{\sum_{ij} R_i^2 Y + 2 \sum_{i < j} R_i R_j R_{ij}}{\sum_{ij} R_i^2 Y + 2 \sum_{i < j} R_i R_j R_{ij}}$$

2.4 Estimation of GA and GAM

The genetic advance (GA) for selection intensity (K) at 5% was calculated by the formula suggested by [12].

$$GA = (K) (\sigma_p) (H^2)$$

Where,

GA = Expected genetic advance at 5% selection intensity, σ_p = Phenotypic standard deviation

H^2 = Heritability and K = selection differential (K= 2.063 at 5% selection intensity).

The genetic advance as percent of the mean (GAM) was calculated by formula given by [7]

$$GAM = \frac{GA}{\bar{X}} * 100$$

Where,

GAM = Genetic advance as percent of mean

GA = Genetic advance at 5% selection intensity

\bar{X} = Population Mean

Heritability (H^2) was computed by formula developed by [12]

$$H^2 = \frac{\sigma_g^2}{\sigma_p^2} * 100$$

Where,

σ_p^2 = Phenotypic Variance σ_g^2 = Genotypic Variance H^2 = Heritability in broad sense

3. RESULTS AND DISCUSSION

Significant differences ($P < 0.01$) were observed among the tested genotypes for all the traits investigated viz. plant height, spike length, number of tillers per plant, number of spikes per plant, number of spikelets per ear, number of grains per ear, biological yield, harvest index, and grain yield. These findings indicate the presence of substantial variability among the genotypes, which can be harnessed through selection and hybridization. The significant differences observed among the genotypes for the studied characteristics justify conducting further genetic analysis [13]. The substantial genetic variation among the genotypes suggests that they exhibit genetic diversity, providing an excellent opportunity for breeders to select suitable genotypes for specific traits of interest in variety development. A similar results reported by [14-16] that the analysis of variance revealed significant genetic variability among all traits for the treatments. Table 3 presents the estimated range,

Comment [IP13]: Not mentioned in results

Comment [IP14]: Not mentioned in results

Comment [IP15]: H2b

Comment [IP16]: Author name with date

minimum and maximum values, along with their corresponding genotypes, mean values, and corresponding standard errors for the nine traits examined in wheat genotypes.

Table 2. ANOVA for nine traits in wheat

Characters	Replication (d.f =4)	Treatment (d.f =7)	Error (d.f =28)
Plant Height (cm)	13.22	108.10	21.51
Spike length (cm)	0.53	1.511	0.68
Number of tillers per plant	3.10	23.377	6.08
Number of spikes per plant	6.25	23.02	7.35
No. of spikelets per ear	1.60	9.585	1.657
No. of Grains per ear	33.06	171.75	73.81
Biological yield (gm)	17.43	2089.39	345.20
Harvest index (%)	22.52	56.52	37.57
Grain yield (gm)	12.00	236.55	37.28

4.1 Variability, Heritability and Genetic advance

Genetic variability parameters viz. GCV, PCV, Heritability (broad sense), Genetic advance and genetic advance as 5 % of mean tabulated in Table 4. PCV (Phenotypic coefficient of variance) is higher than GCV (Genotypic coefficient of variance) for all the characters that indicates an influence of environment on traits. Data pertaining to Number of spikelets per ear recorded highest PCV and GCV followed by Grain yield. The current findings are consistent with the results reported by [17], who also observed high PCV and GCV for grain yield. Similarly, moderate PCV and GCV was recorded for Number of tillers per plant followed by Number of spikes per plant. [18] also reported moderate PCV and GCV number of tillers per plant and low PCV and GCV were recorded for Plant Height, Spike length and Harvest index. The present result in agreement with [19].

Heritability serves as a predictor of the traits that parents will pass on to their offspring. [20] found that selection processes become easier as heritability estimates increased. The heritability (H^2) revealed the extent to which quantitative traits inherited, but it fails to disclose the extent to which genetic gain may be attained by the selection of the ideal plant from the elite populations. Therefore, the combination of heritability and genetic advance is beneficial than either factor alone [21]. Heritability (h^2b) found higher for grain yield (84.2) followed by Number of spikelets per ear (83.4), Number of Grains per ear (82.7) and Plant height (80.1). [17-18], documented similar result that high estimates of heritability. These traits exhibit a high degree of heritability, indicating that genetic factors have a strong influence on their expression. Consequently, environmental factors are likely to have a lesser impact on the development of these traits. Medium (h^2b) recorded for Number of tillers per plant (74) and Number of spikes per plant (68.1) and lower for Harvest index (32.9). Genetic advance (GA) recorded higher Number of spikelets per ear (35.07) followed by Grain yield (11.93). Moderate GA recorded for Plant Height (7.67) and Biological yield (6.88) whereas Spike length (0.62) recorded lowest GA. [22] also found similar results.

Comment [IP17]: Mention author findings

Comment [IP18]: Write the author name here

Comment [IP19]: Again mention author

Comment [IP20]: Add for

Comment [IP21]: Delete

Comment [IP22]: And lower for

Comment [IP23]: Delete, mention author and his findings

4.2 Estimation of Correlation

Assessment of the genotypic and phenotypic correlation between various traits tabulated in Table 5. Data pertaining to plant height it was found that it revealed negative and highly significant correlation with spike length ($r_g = -0.8613$) at genotypic level only. [23] also observed a significant positive correlation between plant height and spike length in their study. Spike length revealed positive and highly significant correlation with number of spikelets per ear ($r_g = 0.9883$) whereas negative and highly significant for plant height ($r_g = -0.8613$) at genotypic level only. Whereas negative and significant correlation was found for number of tiller per plant ($r_p = -0.7068$) and number of spikelets per plant ($r_p = -0.7847$) at phenotypic level only. Number of tillers per plant revealed positive and highly significant correlation with number of spikes per plant ($r_p = 0.9409$, $r_g = 0.9958$) and Grain yield ($r_p = 0.8387$, $r_g = 0.8709$) while positive and significant correlation with Biological yield ($r_p = 0.7789$, $r_g = 0.7683$) at both phenotypic and genotypic level. Whereas negative and significant correlation was found for spike length ($r_p = -0.7068$) at phenotypic level only. [15] found similar results. Number of spikes per plant revealed positive and highly significant correlation with number of tillers per plant ($r_p = 0.9409$, $r_g = 0.9958$) while positive and significant correlation with grain yield ($r_p = 0.7168$, $r_g = 0.7253$) at both phenotypic and genotypic level. Whereas negative and significant correlation was found for spike length ($r_p = -0.7847$) at phenotypic level only. [24] identified noteworthy associations between the plant height and number of spikelets. Biological yield revealed positive and highly significant correlation with grain yield ($r_p = 0.9671$, $r_g = 0.9965$) while positive and significant correlation with Number of tillers per plant ($r_p = 0.7789$, $r_g = 0.7683$) at both phenotypic and genotypic level. Whereas negative and significant correlation was found for harvest index ($r_p = -0.7519$) at phenotypic level only. Number of spikelets per ear revealed positive and highly significant correlation with Number of Grains per ear ($r_p = 0.8573$) at phenotypic level only while negative and significant correlation was found for harvest index ($r_g = -0.8024$) at genotypic level only. [22] also found similar results. Number of Grains per ear revealed positive and highly significant correlation with Number of spikelets per ear ($r_p = 0.8573$) at phenotypic level only while negative and significant correlation was found for harvest index ($r_g = -0.7091$) at genotypic level only. Harvest index revealed negative and significant correlation for grain yield ($r_g = -0.9970$), number of Grains per ear ($r_g = -0.7091$) and Number of spikelets per ear ($r_g = -0.8024$) at genotypic level only while, biological yield ($r_p = -0.7519$) for phenotypic level only. Contrary to the present study, [25] reported a positive correlation between the number of spikelets and tillers per plant with grain yield at both genotypic and phenotypic levels. Grain yield revealed positive and highly significant correlation with number of tillers per plant ($r_p = 0.8387$, $r_g = 0.8709$) and biological yield ($r_p = 0.9671$, $r_g = 0.9965$) while positive and significant correlation with Number of spikes per plant ($r_p = 0.7168$, $r_g = 0.7253$) at both phenotypic and genotypic level. [18] reported yield per head significantly correlated with harvest index. Whereas negative and highly significant correlation was found for harvest index ($r_g = -0.9970$) at genotypic level only. [25] and [23] reported similar results, where the genotypic correlation coefficients (r_g) were found to be higher than the corresponding phenotypic correlation coefficients (r_p) for most of the character combinations. This suggests that the observed correlations between traits are predominantly influenced by genetic factors rather than environmental factors. Genotypic correlations provide valuable insights into the underlying genetic relationships between traits, which can be useful for plant breeders in selecting desirable combinations of traits for crop improvement programs.

Comment [IP24]: Mention author

Comment [IP25]: Mention author and his findings

Comment [IP26]: Mention author

Comment [IP27]: Mention author

Comment [IP28]: Mention author

Comment [IP29]: Mention author

Comment [IP30]: Mention author

4.3 Path analysis

Merely examining correlations does not offer a comprehensive understanding of how each specific trait contributes to the overall picture. Path analysis, on the other hand, proves valuable in evaluating variables by quantifying their respective contributions and distinguishing between partially indirect and direct sources of association. This analytical approach enables a comparative assessment of variables based on the magnitude of their influences. Table 6 and Figure 1 reveals the path analysis through direct and indirect effect of each trait on selected variable viz. grain yield. Biological yield per plant revealed highest positive direct effect (1.0642, 0.8700) with significant association with grain yield (0.9965, 0.9671) at both genotypic and phenotypic level. Similarly harvest index also showed highest negative indirect effect (0.2067, 0.2676) associated with biological yield (-0.2126, -0.2012), Number of spikelets per ear (-0.1658, -0.1337) and Number of grains per ear (-0.1466, -0.1237). This result in agreement with [26-28]. Number of spikelets per ear (0.1115, 0.5200) which exhibits close association with grain yield (0.5324, 0.4622) and Number of tillers reveals least positive direct effect on grain yield (0.0183, 0.0834) at both the levels and significant association with grain yield (0.8709, 0.8387). Number of spikelets per plant and plant height showed positive direct effect on grain yield at genotypic level (0.1018 and 0.0108) and showed negative

direct effect on grain yield at phenotypic level (-0.1834 and -0.1271) respectively. Number of grains per ear (-0.0074, -0.0578) and spike length (-0.0060, -0.5770) exhibited highest negative indirect effect on grain yield at both genotypic and phenotypic levels. Spike length showed highest positive indirect effect through number of spikes per plant (0.4528), number of tillers per plant (0.4078) and plant height (0.3654) at phenotypic level. Similar result in agreement with [14], [18] and [22].

Comment [IP31]: Mention author and his findings

REFERENCES

1. Wani SH, Sheikh F, Najeeb S, Iqbal AM, Kordrostami M, Parray G, Jeberson MS. Genetic variability study in bread wheat (*Triticum aestivum* L.) under temperate conditions. *Current Agriculture Research Journal*. 2018;6(3):268.
2. Bhanu AN, Arun B, Mishra VK. Genetic variability, heritability and correlation study of physiological and yield traits in relation to heat tolerance in wheat (*Triticum aestivum* L.). *Biomedical Journal of Scientific & Technical Research*. 2018;2(1):2112-6.
3. FAO 2022. Crop Prospects and Food Situation – Quarterly Global Report No. 2, July 2022. Rome. <https://doi.org/10.4060/cc0868en>.
4. Crespo-Herrera LA, Crossa J, Huerta-Espino J, Vargas M, Mondal S, Velu G, Payne TS, Braun H, Singh RP. Genetic gains for grain yield in CIMMYT's semi-arid wheat yield trials grown in suboptimal environments. *Crop Science*. 2018;58(5):1890-98.
5. FAO 2021. Crop Prospects and Food Situation - Quarterly Global Report No. 1, March 2021. Rome. <https://doi.org/10.4060/cb3672en>.
6. Verma PN, Singh BN, Yadav RK. Genetic variability and divergence analysis of yield and its contributing traits under sodic soil condition in wheat (*T. aestivum* L.). *Int. J. Agric. Sci*. 2013;3(2):395-9.
7. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybeans. *Agronomy journal*. 1955;47(7):314-8.
8. Dhunde BB, Devmore JP, Mahadik SG, Palshetkar MG, Vanve PB, Bhavne SG, Thorat BS. Correlation and path analysis studies on yield and its components in green gram [*Vigna radiata* L. Wilczek]. *The Pharma Innovation Journal*. 2021;10(1):727-30.
9. Salman MA, Anuradha C, Sridhar V, Babu ER, Pushpavalli SN. Genetic variability for yield and its related traits in green gram [*Vigna radiata* (L.) Wilczek]. *Legume Research-An International Journal*. 2021;1:5.
10. Al-Jibouri H, Miller PA, Robinson HF. Genotypic and environmental variances and covariances in an upland Cotton cross of interspecific origin 1. *Agronomy journal*. 1958;50(10):633-6.
11. Dewey DR, Lu K. A correlation and path-coefficient analysis of components of crested wheatgrass seed production 1. *Agronomy journal*. 1959;51(9):515-8.
12. Allard RW. *Principals of Plant Breeding* John Wiley and Sons. Inc., USA Wiley International Edition. 1960:85.
13. Chaudhary PL, Kumar B, Kumar R. Analysis of Heterosis and Heterobeltiosis for Earliness, Yield and Its Contributing Traits in Okra (*Abelmoschus esculentus* L. Moench). *International Journal of Plant & Soil Science*. 2023;35(11):84-98. <https://doi.org/10.9734/ijpss/2023/v35i112949>
14. Getachew A, Worede F, Alamerew S. Phenotypic variation and traits interrelationships in bread wheat (*Triticum aestivum* L.) genotypes in Northern Ethiopia. *Acta agriculturae Slovenica*. 2021;117(3):1-9.
15. Ullah MI, Mahpara S, Bibi R, Shah RU, Ullah R, Abbas S, Ullah MI, Hassan AM, El-Shehawi AM, Brestic M, Zivcak M. Grain yield and correlated traits of bread wheat lines: Implications for yield improvement. *Saudi Journal of Biological Sciences*. 2021;28(10):5714-9.
16. Yared S, Hussein S, Mark L, Isack M. Genetic variability and association of yield and yield components among bread wheat genotypes under drought-stressed condition. *AJCS*. 2021;15(06):863-870.
17. Upadhyay K, Adhikari NR, Sharma S. Genetic variability and cluster analysis of wheat (*Triticum aestivum* L.) genotypes in foot hill of Nepal. *Archives of Agriculture and Environmental Science*. 2019;4(3):350-5.
18. Rajput RS. Path analysis and genetic parameters for grain yield in bread wheat (*Triticum aestivum* L.). *Annual Research & Review in Biology*. 2019;28:1-8.
19. Bayisa T, Tefera H, Letta T. Genetic variability, heritability and genetic advance among bread wheat genotypes at Southeastern Ethiopia. *Agriculture, Forestry and Fisheries*. 2020;9(4):128-134.
20. Khan AA, Alam MA, Alam MK, Alam MJ, Sarker ZI. Correlation and path analysis of durum wheat (*Triticum turgidum* L. var. Durum). *Bangladesh Journal of Agricultural Research*. 2013;38(3):515-21.

21. Kumar J, Kumar A, Singh SK, Singh L, Kumar A, Chaudhary M, Kumar S, Singh SK. Principal component analysis for yield and its contributing traits in bread wheat (*Triticum aestivum*) genotypes under late sown condition.
22. Kumar P, Solanki YP, Singh V. Genetic Variability and Association of Morpho-physiological Traits in Bread Wheat (*Triticum aestivum* L.). *Current Journal of Applied Science and Technology*. 2020;39(35):95-105.
23. Baye A, Berihun B, Bantayehu M, Derebe B. Genotypic and phenotypic correlation and path coefficient analysis for yield and yield-related traits in advanced bread wheat (*Triticum aestivum* L.) lines. *Cogent Food & Agriculture*. 2020;6(1):1752603.
24. Ul-Allah S, Azeem A, Sher A, Ijaz M, Sattar A, Saleem MA, Bibi M, Abbas N, Hussain M. Assessment of genetic variability and direct-indirect contribution of post-anthesis traits to the grain yield in bread wheat (*Triticum aestivum*) at different sowing dates. *Int. J. Agric. Biol.* 2021;26:193-200.
25. Verma P, Saini P, Singh V, Yashveer S. Genetic variability of wheat (*Triticum aestivum* L.) genotypes for agro-morphological traits and their correlation and path analysis. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(4):2290-4.
26. Ayer DK, Sharma A, Ojha BR, Paudel A, Dhakal K. Correlation and path coefficient analysis in advanced wheat genotypes. *SAARC Journal of Agriculture*. 2017;15(1):1-2.
27. Meles B, Mohammed W, Tsehaye Y. Genetic variability, correlation and path analysis of yield and grain quality traits in bread wheat (*Triticum aestivum* L.) genotypes at Axum, Northern Ethiopia. *Journal of plant breeding and crop science*. 2017;9(10):175-85.
28. Chimdesa O, Mohammed W, Eticha F. Analysis of genetic variability among bread wheat (*Triticum aestivum* L.) genotypes for growth, yield and yield components in Bore district, Oromia regional state. *Agriculture, Forestry and Fisheries*. 2017;6(6):188-99., 188-199.

GENOTYPES	PH	SL	NTP	NSP	BY	NSPE	NGPE	HI	GY
(GS/2020-21/8049 X PBW 766) 1	80.00	12.60	14.00	14.00	86.40	21.00	70.20	38.58	32.86
(GS/2020-21/8049 X PBW 766) 2	81.40	12.40	11.40	11.40	60.40	19.40	66.80	42.10	25.28
(GS/2020-21/8049 X PBW 766) 3	79.20	13.40	7.40	7.40	46.60	20.20	59.80	43.05	18.78
(GS/2020-21/8049 X PBW 766) 4	87.20	12.00	11.20	13.20	55.60	18.60	56.80	43.53	23.96
(GS/2020-21/8049 X PBW 766) 5	89.00	12.80	12.20	12.20	89.40	22.60	73.00	41.72	36.85
(GS/2020-21/8049 X PBW 766) 6	87.40	12.60	11.00	11.00	56.00	19.80	64.00	44.55	24.12
PBW 766	92.00	11.80	14.20	13.80	103.80	19.80	61.60	37.24	38.40
GS/2020-21/8049	87.4	11.8	13	13	60.8	18.2	57.8	47.91	28.8
Mean	85.45	12.42	11.80	12.00	69.87	19.95	63.75	42.34	28.63
min	79.2	11.8	7.4	7.4	46.6	18.2	56.8	37.24	18.78
max	91	13.4	14.2	14	103.8	22.6	73	47.91	38.4
C.V.	5.42	6.63	20.90	22.59	26.59	6.45	13.47	14.47	21.32
C.D. 5%	6.00	-	3.19	3.51	24.07	1.66	11.13	-	7.91
C.D. 1%	8.10	-	4.31	4.73	32.47	2.24	15.01	-	10.67

PH- Plant height, SL- Spike length, NTP- Number of tillers per plant, NSP- Number of spikes per plant, NSPE- Number of spikelets per ear, NGPE- Number of Grains per ear, BY- Biological yield, HI- Harvest index and GY- Grain yield

Table 3. Mean of parent and crosses for nine traits with CV, CD at 5 and 1 % respectively

Characters	Min	Max	Mean	GCV	PCV	h ² b	GA (5%)	GA Mean (5%)
Plant Height (cm)	79.2	91	85.45	4.870	5.441	80.1	7.672	8.979
Spike length (cm)	11.8	13.4	12.42	3.280	4.424	55	0.622	5.009
Number of tillers per plant	7.4	14.2	11.80	15.757	18.322	74	3.294	27.915
Number of spikes per plant	7.4	14	12.00	14.757	17.884	68.1	3.010	25.083
Number of spikelets per ear	46.6	103.8	69.87	26.676	29.206	83.4	35.071	50.191
Number of Grains per ear	18.2	22.6	19.95	6.312	6.940	82.7	2.359	11.826
Biological yield (gm)	56.8	73	63.75	6.942	9.194	57	6.885	10.799
Harvest index (%)	37.24	47.91	42.34	4.537	7.906	32.9	2.271	5.363
Grain yield (gm)	18.78	38.4	28.63	22.046	24.020	84.2	11.936	41.683

Table 4. Variability (GCV & PCV), Heritability (broad sense), Genetic advance (GA) and GAM at 5% in wheat

		Phenotypical and Genotypic Correlation Matrix								
		PH	SL	NTP	NSP	BY	NSPE	NGPE	HI	GY
PH	r _p	1.0000	-0.6333	0.4832	0.4993	0.4751	-0.0599	-0.1433	-0.0115	0.5745
	r _g	1.0000	-0.8613 **	0.6021	0.6664	0.5501	-0.0811	-0.2107	0.0332	0.6796
SL	r _p		1.0000	-0.7068*	-0.7847*	-0.3074	0.5950	0.3871	-0.0477	-0.4048
	r _g		1.0000	-1.2255	-1.4055	-0.5855	0.9883**	0.6233	0.4326	-0.6089
NTP	r _p			1.0000	0.9409**	0.7789*	0.0458	0.2900	-0.3861	0.8387**
	r _g			1.0000	0.9958**	0.7683*	0.0717	0.3880	-0.4136	0.8709**
NSP	r _p				1.0000	0.6473	-0.0808	0.1440	-0.3008	0.7168*
	r _g				1.0000	0.6023	-0.1826	0.1556	-0.1443	0.7253*
BY	r _p					1.0000	0.5088	0.5100	-0.7519*	0.9671**
	r _g					1.0000	0.5817	0.6473	-1.0288	0.9965**
NSPE	r _p						1.0000	0.8573 **	-0.4997	0.4622
	r _g						1.0000	1.1176	-0.8024 *	0.5324

NGPE	r_p	1.0000	-0.4623	0.5007
	r_g	1.0000	-0.7091 *	0.6765
HI	r_p		1.0000	-0.5678
	r_g		1.0000	-0.9970**
GY	r_p			1.0000
	r_g			1.0000

PH- Plant height, SL- Spike length, NTP- Number of tillers per plant, NSP- Number of spikes per plant, NSPE- Number of spikelets per ear, NGPE- Number of Grains per ear, BY- Biological yield, HI- Harvest index and GY- Grain yield

Table 5. Phenotypal and Genotypic Correlation for nine traits in wheat

Figure 1. Genotypic and Phenotypic Path matrix of nine traits in wheat for Grain Yield

Figure 1. Genotypic and Phenotypic Path matrix for Grain Yield




Table 6. Phenotypic and Genotypic Path matrix of nine traits in wheat for Grain Yield

Phenotypic and Genotypic Path Matrix for GY

		PH	SL	NTP	NSP	BY	NSPE	NGPE	HI	GY
PH	P	-0.1271	0.0805	-0.0614	-0.0635	-0.0604	0.0076	0.0182	0.0015	0.5745
	G	0.0108	-0.0093	0.0065	0.0072	0.006	-0.0009	-0.0023	0.0004	0.6796
SL	P	0.3654	-0.5770	0.4078	0.4528	0.1773	-0.3433	-0.2234	0.0275	-0.4048
	G	0.0052	-0.0060	0.0073	0.0084	0.0035	-0.0059	-0.0037	-0.0026	-0.6089
NTP	P	0.0403	-0.0590	0.0834	0.0785	0.065	0.0038	0.0242	-0.0322	0.8387
	G	0.0110	-0.0225	0.0183	0.0182	0.0141	0.0013	0.0071	-0.0076	0.8709
NSP	P	-0.0916	0.1439	-0.1725	-0.1834	-0.1187	0.0148	-0.0264	0.0552	0.7168
	G	0.0678	-0.1430	0.1013	0.1018	0.0613	-0.0186	0.0158	-0.0147	0.7253
BY	P	0.4133	-0.2674	0.6776	0.5631	0.8700	0.4426	0.4437	-0.6541	0.9671
	G	0.5854	-0.6231	0.8177	0.641	1.0642	0.6191	0.6889	-1.0949	0.9965
NSPE	P	-0.0312	0.3094	0.0238	-0.042	0.2645	0.5200	0.4458	-0.2598	0.4622
	G	-0.0090	0.1102	0.008	-0.0204	0.0649	0.1115	0.1246	-0.0895	0.5324
NGPE	P	0.0083	-0.0224	-0.0167	-0.0083	-0.0295	-0.0495	-0.0578	0.0267	0.5007
	G	0.0016	-0.0046	-0.0029	-0.0012	-0.0048	-0.0083	-0.0074	0.0053	0.6765
HI	P	-0.0031	-0.0128	-0.1033	-0.0805	-0.2012	-0.1337	-0.1237	0.2676	-0.5678
	G	0.0069	0.0894	-0.0855	-0.0298	-0.2126	-0.1658	-0.1466	0.2067	-0.9970

PH- Plant height, SL- Spike length, NTP- Number of tillers per plant, NSP- Number of spikes per plant, NSPE- Number of spikelets per ear, NGPE- Number of Grains per ear, BY- Biological yield, HI- Harvest index and GY- Grain yield

UNDER PEER REVIEW

