

Unraveling the Genetic Architecture of Rice (*Oryza sativa* L.) Using Variability, Correlation, Path analysis and Diversity Analysis: Insights for Crop Improvement Strategies

ABSTRACT

The present study aimed to assess the genetic architecture of rice through various biometrical techniques and this was achieved through a field experiment conducted during the *Kharif* 2022 season utilizing 23 distinct genotypes of rice. The trial was performed at the field experimentation center, Department of genetics and plant breeding, School of agriculture, Lovely Professional University. The research concluded that the highest genotypic and phenotypic coefficients of variation were for the number of productive tillers per plant, grain L/B ratio and grain yield per plant. Additionally, high heritability was observed for all traits except for kernel breadth. The investigation also explains high genetic advance as a percent of the mean for all traits except spikelet fertility and kernel breadth. Grain yield per plant had a significant positive genotypic correlation with the number of productive tillers per plant, filled grains per panicle, and total number of grains. Path analysis revealed that kernel L/B ratio, kernel length, filled grains per panicle, total number of grains, and number of productive tillers per plant had the maximum positive direct effects. The study suggests that selection based on these traits could be more effective in improving rice yield. The germplasms falling in different clusters with high mean for grain yield and other component characters can be utilized for hybridization programme to obtain elite segregants.

Keywords: Variability, Correlation, Path analysis, Diversity, Rice

1. INTRODUCTION

Rice is one of the chief cereals consumed across the world. Rice proteins are known for having amino corrosive substances and are uniquely wealthy in lysine content [1]. India ranks first in the area of rice production and exports. The interest in food is consistently extending with the increment of populace, making a yearly increment of 3% for each year of popular rice [2]. The production of rice in India, for example, was estimated to be around 118.87 million tonnes in the year 2020-21, according to the Ministry of Agriculture and Farmers' Welfare. Variation among the varieties for the economical yield and associated traits is the foundation of any selection program in breeding.

The heritable variability within the population or intermating resource can be concluded by genetic diversity. This helps to succeed in the subject of plant breeding by projecting genetic variation, and keen information about the concerned traits in the given resource. The information about the gene exchange to offsprings can be evaluated using heritability. Heritability provides information about the capability of a trait that can be transmitted to possible generations. Heritability estimates thus helps to predict the expression of traits in further generations. A combination of genetic advance and heritability is helpful for proper selection as said by [3]. High heritability coupled with high genetic advance results helps breeder to select preferable estimations in breeding programme. Correlation studies can provide information on the improvement of multiple characteristics simultaneously. Therefore, knowledge of the relationship between yield and its contributing characters is

essential in rice breeding programs. Path coefficient analysis is a partial regression coefficient and well-established technique practiced in breeding to conclude the relation between grain yield and its contributing characters. The path coefficient analysis predicts the direct and indirect effects of source variables on the expressing variable. The nature of effect like direct or indirect effect of yield and its characteristics can be explained through path coefficient analysis [4]. Thereafter this method can be used in selection techniques which helps to read about specific traits resulting in correlation between grain yield and other traits.

2. MATERIAL AND METHODS

The study was conducted at the Field Experimentation Centre, Department of Genetics and Plant Breeding, School of Agriculture, Lovely Professional University in Punjab.

The experimental material for this study consisted of 23 genotypes sourced from various places, such as Punjab Agricultural University (PAU) and Agricultural Research Station (ARS) in Nellore, Andhra Pradesh. The genotypes were used for a field experiment conducted during the *Kharif* 2022 season. The experimental design used was a randomized block design with three replications, and the spacing between rows and plants was 30 cm and 15cm, respectively. The cultural practices followed during the experiment were based on recommended guidelines of PAU to ensure proper growth and development of the plants.

2.1 Recorded observations

Five randomly selected plants from each entry and replication were evaluated for traits like days to 50% flowering (DF), plant height (PH) (cm), panicle length (PL) (cm), number of tillers per plant (NTL), number of productive tillers per plant (NPL), number of Spikelets per panicle (NSP), spikelet fertility (SF), flag leaf length (FL) (cm), number of filled grains per panicle (NFG), days to maturity (DM), single plant yield (SYP) (g), 1000 grain weight (TW) (g), biological yield (BY) (g) and harvest index (HI).

2.2 Statistical analysis

The techniques suggested by Singh and Chaudhary [5] are used to evaluate the genotypic coefficient of variation (GCV) and the phenotypic coefficient of variation (PCV). Heritability in a broad sense is concluded using the formula established by [6]. Genetic advance is estimated using the method proposed by [3]. The path analysis is calculated by the method of Dewey and Lu [7]. The concept of D^2 statistics was originally developed by P.C. Mahalanobis [8].

3. RESULTS AND DISCUSSION

The analysis of variance conducted in this study indicated that the genotypes used in the experiment showed significant differences for each trait studied, as shown in Table 1. This indicates the presence of genetic variation among the genotypes used in the study [27]. The mean performance of the genotypes also showed a wide range, indicating the existence of significant variation among the genotypes. Similar results highlighting the presence of variation in different quantitative traits of rice have been reported in other studies [9-11].

Table 1: Analysis of variance for grain yield and yield components in rice

Source of variation	d.f	DF	PH	PL	NTL	NPL	NSP	SF
Replication	2	13.43	62.26	9.33	12.28	6.44	78.98	6.04
Treatment	23	40.57	588.82	21.75	6.66	8.16	2317.76	42.33

Error	44	3.27	8.19	1.48	2.99	1.89	20.50	2.55
C.V.		2.17	2.35	4.14	13.95	12.07	1.85	1.84
Source of variation	d.f	FL	DM	BY	SPY	TW	NFG	HI
Replication	2	11.75	54.91	87.62	24.78	9.49	6.64	0.03
Treatment	23	19.52	236.23	168.78	17.77	9.34	1565.38	0.006
Error	44	2.92	13.16	26.73	7.18	2.48	9.60	0.001
C.V.		5.71	2.88	5.92	8.08	6.80	1.46	9.70

3.1 GCV and PCV of variation

The phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) disguise the environmental effect over several traits. Estimates of minute difference between the PCV and GCV define the environmental effect on the appearance of traits. Low estimates of PCV and GCV were recorded for days to 50% flowering, panicle length, spikelet fertility, flag leaf length, days to maturity, biological yield, grain yield per plant and 1000 seed weight. Similar findings were recorded for days to 50% flowering, panicle length, spikelet fertility by [9], days to maturity by [12] that say direct selection for these traits is not recommended. Moderate PCV and GCV were recorded for plant height, number of productive tillers, number of spikelets per panicle, and number of filled grains per panicle. Similar findings were recorded for plant height, harvest index, this is the indication of an appreciable amount of genetic variability and negligible influence of environment over the expression of the trait and suggests the scope of improvement of these genotypes for the traits, relevant discovery by [13-15]. Number of total tillers per plant was recorded as low PCV and Medium GCV.

3.2 Heritability and Genetic advance

As mentioned in (Table 2), the estimated heritability varied between (55.17-99.39) percent between days to 50% flowering to harvest index. High heritability recorded characteristics for days to 50% flowering, plant height, panicle length, number of spikelets per panicle, number of filled grains per panicle, spikelet fertility, flag leaf length, days to maturity, biological length, 1000 grain weight, harvest index and, harvest index shows the presence of additive gene action. Similar results were recorded for days to 50 % flowering by [16], Plant height by [17], Moderate heritability recorded characteristics for number of total tillers per plant, single plant yield, Similar results were recorded for number of total tillers per plant by [18]. Low heritability was not recorded, likewise results in [19]. Low or moderate estimates of heritability coupled with genetic advance indicated the presence of non-additive gene action, this criterion was observed in traits like, single plant yield and no. of tillers per plant.

The genetic advancement percentage mean observed ranges from (1.69-56.75) days to 50% flowering to harvest index the presence of high GA percentage is observed in plant height, number spikelet's per panicle, number of filled grains per panicle, likewise results observed for plant height, spikelet's per panicle by [18]. Moderate GA percentage are observed in days to maturity, biological yield, and similar results for days to maturity by [20]. Low GA percentage observed in days to 50% flowering, panicle length, number of total tillers per plant, number of productive tillers, spikelet fertility, flag leaf length, single plant yield, 1000 grain weight, harvest index similar results recorded for days to 50 % flowering by [16] and spikelet fertility for [21].

Table 2: Coefficient of variation, heritability, and genetic advance of rice genotypes.

Characters	Mean	Range		Variance		Coefficient of variation		Heritability (%)	Genetic advance (5%)	G.A. as percentage of mean
		Min.	Max.	Genotypic	Phenotypic	Genotypic	Phenotypic			
DF	83.26	74.33	91.33	12.43	13.52	4.23	4.41	91.9	6.96	8.36
PH	121.52	91.47	136.13	193.5	196.20	11.44	11.52	98.6	28.45	23.41
PL	29.36	24.87	33.47	6.755	7.24	8.85	9.16	93.1	5.16	17.60
NTP	12.39	10.33	15.70	1.225	2.22	8.93	12.02	55.17	1.69	13.66
NPL	11.39	9.33	15.03	2.0894	2.71	12.68	14.47	76.82	2.60	22.90
NSF	245.07	170.13	286.93	765.75	772.58	11.29	11.34	99.1	56.75	23.15
SF	86.69	81.20	94.86	13.261	14.11	4.20	4.33	93.9	7.27	8.38
FL	29.95	25.83	33.67	5.532	6.50	7.85	8.51	85.03	4.46	14.91
DM	125.78	118.67	142.67	74.3584	78.74	6.85	7.05	94.43	17.26	13.72
BY	87.28	77.20	105.67	47.3509	56.20	7.88	8.59	84.16	13.04	14.90
SPY	33.18	28.73	36.40	3.5304	5.923	5.66	7.33	59.6	2.98	9.04
TW	23.17	20.20	29.40	2.2849	3.11	6.52	7.61	73.41	2.66	11.51
NFP	211.88	142.53	245.53	518.5	521.79	10.74	10.78	99.39	46.76	22.07
HI	0.38	0.28	0.47	0.01	0.02	11.10	12.44	79.72	0.07	20.43

3.3 Correlation coefficient analysis

The genotypic and phenotypic correlation values are showing minute differences, which destinate the effect of environment is low on these traits. The grain yield per plant revealed positively correlation with panicle length, no. of tillers per plant, no. of productive tillers per plant, no. of spikelet's per panicle, spikelet fertility, days to maturity, no. of filled grains per panicle and harvest index which designates that these traits are positively arbitrary at both phenotypic and genotypic levels. Similar results were recorded for panicle length, number of productive tillers per plant, total tillers, days to maturity, total tillers per plant, and number of filled grains per panicle [22]. Hence the selection of these traits for high yield is recommended. Negatively correlated traits for both phenotypic and genotypic correlations viz. days 50% flowering, plant height, flag leaf length, biological yield, and thousand seed weight. Similar results recorded for plant height by [23], flag leaf length by [14] that are not suggested for trait selection to improve the grain yield.

3.4 Path Coefficient analysis

The outcome of path coefficient analysis says that the fourteen traits viz, panicle length, number of productive tillers per panicle, flag leaf length, biological yield, number of filled grains per panicle, and harvest index have a direct positive effect and accordingly these traits are useful for selection to improve the grain yield. Similar results for flag leaf length and panicle length by [24]. Characters showed highly positive effects for grain yield also revealed high direct effects. Negative direct effects on grain yield are days to 50 % flowering, plant height, number of total tillers per panicle, number of spikelets per panicle, spikelet fertility, days to maturity, and test weight. Similar results for days to maturity by [15] plant height by [25-16].

Fig.1 Phenotypic (r_p) (above diagonal) and Genotypic (r_g) (below diagonal) correlation coefficients among 14 characters in 23 genotypes



TABLE 3: Genotypic path coefficient for grain yield and its component traits in rice

Traits	DF	PH	PL	NTL	NPL	NSP	SF	FL	DM	BY	TW	NFG	HI	SYP
DF	-0.2134	0.0135	0.0344	-0.1163	-0.0862	0.029	0.0952	-0.0647	-0.109	0.0219	0.0841	0.0706	0.0055	-0.178
PH	0.0033	-0.0515	-0.0336	0.0068	0.0153	-0.0292	-0.0048	0.0007	-0.0241	-0.0071	0.0221	-0.0323	0.0087	-0.2093
PL	-0.0248	0.1003	0.1536	0.0186	-0.0184	0.0666	0.0403	0.0276	0.0282	0.0117	-0.0411	0.0846	-0.0064	0.0251
NTL	-0.1494	0.0362	-0.0333	-0.2741	-0.2828	0.1398	-0.0405	0.0864	-0.0917	0.1244	-0.0053	0.1333	-0.1388	0.4318
NPL	0.186	-0.1364	-0.055	0.4748	0.4603	-0.2451	0.0836	-0.1589	0.0632	-0.178	0.0943	-0.2232	0.1823	0.2782
NSP	0.634	-2.6435	-2.021	2.3774	2.4815	-4.661	1.5028	-0.0444	-2.4492	-1.123	0.1759	-4.3014	0.4019	0.0904
SF	0.9127	-0.1896	-0.5363	-0.302	-0.3714	0.6597	-2.046	-0.2117	1.0279	0.5288	0.0634	-0.1376	-0.5582	0.2514
FL	0.0457	-0.0019	0.0271	-0.0475	-0.0521	0.0014	0.0156	0.1508	-0.0245	-0.0145	-0.0568	0.0044	-0.0067	-0.1545
DM	-0.0534	-0.0488	-0.0192	-0.035	-0.0144	-0.0549	0.0525	0.017	-0.1046	0.0034	0.0236	-0.0372	-0.0209	0.2989
BY	-0.1084	0.1454	0.0803	-0.4792	-0.4083	0.2544	-0.2729	-0.1014	-0.0348	1.0559	-0.4383	0.1633	-0.9479	-0.4935
TW	0.1035	0.1125	0.0703	-0.0051	-0.0538	0.0099	0.0081	0.0989	0.0594	0.109	-0.2626	0.0136	-0.0553	-0.1687
NFG	-1.4649	2.7753	2.4369	-2.1514	-2.146	4.084	0.2976	0.1299	1.5765	0.6843	-0.2295	4.4254	0.053	0.1863
HI	-0.0489	-0.3207	-0.0792	0.9648	0.7545	-0.1643	0.5199	-0.0847	0.3816	-1.7104	0.4014	0.0228	1.9053	0.8223

RESIDUAL EFFECT = 0.1178

3.5 Diversity analysis

A range of variation was observed among twenty-three rice germplasms that were evaluated. These germplasms were then grouped into different clusters based on their similarity, as determined by the Mahalanobis D^2 statistical method. To select parents for a hybridization program that can lead to maximum heterosis, it is recommended to consider the magnitude of genetic distance, the contribution of different characters to total divergence, and the magnitude of cluster means for different characters' performance. The

germplasms in clusters I & III, II & I, III & I, IV & III, and V & I exhibit significant variability, and it is expected that subsequent generations will accumulate favorable genes, leading to high heterosis.

Fig. 2 Graphical depiction of a dendrogram that shows the clustering of 23 rice genotypes based on 14 observable traits.




Table 4 Average inter and intra-cluster distance values in rice

	Cluster I	cluster II	cluster III	cluster IV	Cluster V
cluster 1	0	64.074	101.963	46.8275	99.7712
cluster 2	64.075	21.91	45.863	35.3407	38.1032
cluster 3	101.963	45.8629	14.34	79.1143	23.5643
cluster 4	46.828	35.3407	79.114	20.71	71.0568
cluster 5	99.771	38.1032	23.564	71.0568	16.81

4. CONCLUSION

It is concluded that the high heritability coupled with high genetic advance possesses the additive gene action which prioritizes selection of traits for breeding program and our experiment also concludes that the same for traits like plant height, no. of productive tillers per plant, 1000 seed weight and harvest index shows high heritability and high genetic advance. The genotypic and phenotypic correlations result the positive and direct effect of harvest index and number of tillers per plant over grain yield, which indicates the true

association between the traits and environmental effect is not evident. Hence concluded the selection among the traits is preferred to increase the yield of rice per plant.

REFERENCES

1. Rathna Priya TS, Eliazer Nelson AR, Ravichandran K, Antony U. Nutritional and functional properties of coloured rice varieties of South India: a review. *Journal of Ethnic Foods*. 2019;6(1):1-1.
2. Dong J, Xiao X, Kou W, Qin Y, Zhang G, Li L, Jin C, Zhou Y, Wang J, Biradar C, Liu J. Tracking the dynamics of paddy rice planting area in 1986–2010 through time series Landsat images and phenology-based algorithms. *Remote Sensing of Environment*. 2015; 160:99-113.
3. Johnson HW, Robinson HF, Comstock RE. Genotypic and phenotypic correlations in soybeans and their implications in selection. *Agronomy journal*. 1955;47(10):477-83.
4. Prasad KR, Radha-Krishna KV, Bhave MH, Subba-Rao LV. Genetic variability, heritability and genetic advance in boro rice (*Oryza sativa* L.) germplasm. *International Journal of Current Microbiology and Applied Sciences*. 2017;6(4):1261-6.
5. Singh RK, Chaudhary BD, 1985. *Biometrical Methods in Quantitative Genetic Analysis*, pp. 318. Kalayani Publishers, New Delhi, India
6. Hanson CH, Robinson HF, Comstock RE. Biometrical studies of yield in segregating populations of Korean lespedeza 1. *Agronomy journal*. 1956;48(6):268-72.
7. Dewey DR, Lu K. A correlation and path-coefficient analysis of components of crested wheatgrass seed production 1. *Agronomy journal*. 1959;51(9):515-8.
8. Mahalanobis PC. Historical note on the D^2 -statistics, appendix I. anthropological survey of United Provinces, 1941: a statistical study.
9. Maurya RK, Dwivedi DK, Khan NA, Giri SP, Dixit S. Genetic variability studies for qualitative and quantitative traits in rice (*Oryza sativa* L.). *The Pharma Innovation Journal*. 2022;11(5):1140-3.
10. Deepthi KP, Mohan YC, Hemalatha V, Yamini KN, Singh TV. Genetic variability and character association studies for yield and yield related, floral and quality traits in maintainer lines of rice (*Oryza sativa* L.). *The Pharma Innovation Journal*. 2022;11(2):191-7.
11. Akshay M, Chandra BS, Devi KR, Hari Y. Genetic variability studies for yield and its attributes, quality and nutritional traits in rice (*Oryza sativa* L.). *The Pharma Innovation Journal*. 2022;11(5):167-72.
12. THUY NP, NHU TT, TRAI NN, NHU NK, THAO NH, PHONG VT, KHOA BD, LUAN NT. Genetic variability for micronutrients under an aerobic condition in local landraces of rice from different agro-ecological regions of Karnataka, India. *Biodiversitas Journal of Biological Diversity*. 2023;24(1).
13. Naik MVK, Pillai MA, Saravanan S. Assessment of Correlation and Path Coefficient Analysis for Yield Attributing and Quality Traits in promising rice varieties cultivated in Tamil Nadu. *The Journal of Phytopharmacology*. 2021;10(2):139-143.
14. Yadav EK, Pitha CC. Studies on Genetic Variability, Divergence and Characters Associated with Yield Components in Rice. *International Journal of Environment and Climate Change*. 2022;12(11):2105-15.
15. Beena R, Veena V, Jaslam MP, Nithya N, Adarsh VS. Germplasm innovation for high-temperature tolerance from traditional rice accessions of Kerala using genetic variability, genetic advance, path coefficient analysis and principal component analysis. *Journal of Crop Science and Biotechnology*. 2021;24(5):555-66.

16. Basavaraj PS, Gireesh C, Bharamappanavara M, Manoj CA, Ishwaryalakshmi LV, Senguttuvel P, Sundaram RM, Subbarao LV, Anantha MS. Genetic analysis of introgression lines of *Oryza rufipogon* for improvement of low phosphorous tolerance in indica rice. *Indian Journal of Genetics and Plant Breeding*. 2022;82(02):135-42.
17. Awad-Allah MM, Elekhtyar NM, El-Abd MA, Abdelkader MF, Mahmoud MH, Mohamed AH, El-Diasty MZ, Said MM, Shamseldin SA, Abdein MA. Development of new restorer lines carrying some restoring fertility genes with flowering, yield and grains quality characteristics in rice (*Oryza sativa* L.). *Genes*. 2022;13(3):458.
18. Surjaye N, Singh Y, Singh SK, Rahangdale S, Mehta AK. Genetic Variability, Correlation and Path Coefficient Study for Various Yield and Quality Traits in NPT Lines of Rice (*Oryza sativa* L.). *Environment and Ecology*. 2022;40(1):115-22.
19. Keerthiraj B, Biju S. Genetic variability, heritability and genetic advance of yield and lodging-related traits in rice (*Oryza sativa* L.). *Electronic Journal of Plant Breeding*. 2020;11(04):1093-8.
20. Chavan BR, Dalvi VV, Kunkerkar RL, Mane AV, Gokhale NB. Studies on genetic variability for yield and yield contributing traits in aromatic rice (*Oryza sativa* L.). *The Pharma Innovation Journal* 2022;11(2):1732-1735.
21. Pallavi M, Shanthi P, Kumar VR. Genetic variability, heritability and genetic advance for yield and related traits in rice (*Oryza sativa* L.). *The Pharma Innovation Journal* 2022; 11(9): 429-433.
22. Lakshmi MS, Suneetha Y, Srinivas T. Genetic variability, correlation and path analysis for grain yield and yield components in rice genotypes. *Journal of Pharmacognosy and Phytochemistry*. 2021;10(1):1369-72.
23. Jasmine C, Shivani D, Senguttuvel P, Naik DS. Genetic variability and association studies in maintainer and restorer lines of rice [*Oryza sativa* (L.)]. *The Pharma Innovation Journal*. 2022;11(1):569-76.
24. Faysal AS, Ali L, Azam MG, Sarker U, Ercisli S, Golokhvast KS, Marc RA. Genetic variability, character association, and path coefficient analysis in transplant Aman rice genotypes. *Plants*. 2022;11(21):2952.
25. Shahani M, Kumar R, Nilanjaya AK, Mishra A, Bijarania S. Association study on yielding attributes and brown spot disease components in rice germplasm under aerobic condition. *The Pharma Innovation Journal* 2022;11(2):2163-2166.
26. Kondi RK, Kar S, Mandawi NC. Study of genetic parameters, correlation and path analysis for yield and quality characters in fine scented rice genotypes. *ORYZA-An International Journal of Rice*. 2022;59(1):20-30.
27. Chaudhary PL, Kumar B, Kumar R. Analysis of Heterosis and Heterobeltiosis for Earliness, Yield and Its Contributing Traits in Okra (*Abelmoschus esculentus* L. Moench). *International Journal of Plant & Soil Science*. 2023;35(11):84-98.