

# Pokkali rice cultivation: An Review on the indigenous traditional rice cultivation method in Kerala

## Abstract

The Pokkali style of rice farming in Kerala's acidic, saline soil is a distinctive way of rice cultivation. A native historic salt-tolerant type of rice called Pokkali, with a 120-day growing season, can withstand flooding by growing up to a height of 1.5 meters.- Chettivirippu, Vyttila 1, Vyttila 2, Vyttila 3, Vyttila 4 and Vyttila 5 are the other rice varieties cultivated using the Pokkali rice cultivation method. With this technique, a single crop of rice crop is harvested on mounds during the low salinity part of the production cycle (June to mid-October), with prawn aquaculture taking place during the high salinity phase (November to April). Pokkali farmers only produce 2000 kg of rice on average per hectare, making it slightly unprofitable. The residue of rice plant residue after harvesting are left to rot in the water, which subsequently acts as prawn feed. Prawn farming takes place from November to April. Sluice gates help in prawn culture filtering by maximizing the amount of saline water that enters the field and preventing the prawns from escaping. The two main species of cultivated prawn are *Penaeus monodon* (Tiger prawn) and *Penaeus indicus* (White prawn). Prawns are typically collected in mid-April, especially two to three days before new full moon days, as tidal activity affects their movement. Prawn excreta serve as a natural fertilizer for rice cultivation,; prawns feeding on rice cultivation stubbles helps to minimize the occurrence of illness in rice to a minimal minimum. It is an organic rice production approach with lower costs than commercial rice farming because chemical fertilizers and pesticides are not used.

**Keywords:** Ecosystem, Pokkali, Prawn farming, Relay cropping and Vyttila

## Introduction

Kerala has both upland and low-lying locations where rice is grown (Vijayan, 2016). The state's diverse rice ecology is significantly influenced by the local climate and topography (Kumar & Kunhamu, 2021). In Kerala, there are several different methods of growing rice, including intercropping it with other crops, relay planting, sequential cropping, and the Integrated Farming System (IFS) (Kerala State Biodiversity Board, 2009). Moreover, the rice ecosystems in Kerala varies from below sea level, as is the case with Pokkali, Kuttanad, and Kole lands, to the rice paddies in the High Mountains of Idukki and Wayanad (800 to 1500 m above mean sea level) (Kumar & Kunhamu, 2021). Saline-resistant indigenous rice cultivars helped farmers in Kerala's coastal areas create a distinctive method of farming (Gopi & Manjula, 2018). In Kerala, over 25% of the entire paddy lands are submerged in water; as a result, integrated rice-shrimp/prawn farming is done here (Jayan & Sathyanathan, 2010). The Pokkali system of rice cultivation is a traditional technique for growing rice that systematically combines prawn farming with the saline-tolerant wild rice type known as Pokkali. Considering the state's rice consumption is predicted to be between 3.5 and 4 million tonnes per year The state's rice consumption is predicted to be between 3.5 and 4 million tonnes per year, but production is just a

sixth of this (Krishnankutty *et al.*, 2021). The overall area under Pokkali has decreased from 25,000 ha a few decades ago to about 8,500 ha today, of which only 5,500 ha are really used for rice cultivation. The remainder is either kept fallow or is primarily used for prawn farming (Suchithra & Venugopal, 2005). Just 967 acres of paddy is periodically grown using the Pokkali method when the climate is suitable, with the area under Pokkali agriculture decreasing yearly (Jayan & Sathyanathan, 2010). Due to the shrinking area under rice cultivation and the widespread conversion of paddy lands for other crops or for residential uses, there is a severe problem with the traditional method of rice farming in many regions of Kerala (Kumar, 2007). In order to boost the state's rice production, attention must be paid to traditional rice growing techniques, including both wild and cultivable kinds. Understanding the Pokkali rice farming method used in Kerala is the goal. In areas of the wetland environment, fishing is the main industry (Maliyekkal & Thomson, 2008).

### Geographical Location

Kerala is located in the south-west region of the Indian peninsula, in the southern portion of the Western Ghats, bordered to the east and north-east by Tamil Nadu and Karnataka, and to the west by the Arabian Sea (Krishnankutty *et al.*, 2021). Kerala state is essentially categorized into 13 agroecosystems, of which eight are relevant for rice growing, based on topography, soil, abiotic variables, ~~and resource endowments as well as resource endowments, and~~ the seasonality of cultivation (Kumari, 2012). The Laterite Midlands, Malayorum (hill slopes), the Palakkad Plains, the Chittoor Black Soil, the Kuttanad, the Pokkali, Kole and Kaipad, the Onattukara, and the High Ranges are among of them (Kumar & Kunhamu, 2021). Kerala has a variety of geographical features, including wetlands, plains, and coastland to the west, as well as foothills of the Western Ghats to the east (Krishnankutty *et al.*, 2021). ~~Both wild and cultivated rice in the state have a wide variety of germplasm as a result of~~ Wild and cultivated rice in the state have a wide variety of germplasm due to the ecological circumstances there (Sharma, 1991). Kerala, Kuttanad and Palakkad are two of Kerala's principal ~~rice-rice~~-growing regions (Vijayan, 2016). The overall Pokkali lands were thought to measure 25,000 acres, but vast tracts have been transformed for coconut farming and other uses (Jayan & Sathyanathan, 2010).

The Kuttanad region, which produces roughly 20% of Kerala's rice, is referred to as the "Rice Bowl of Kerala" (Thomas, 2002). Located between the districts of Ernakulam and Alappuzha, Kuttanad is a lowlying 875 km<sup>2</sup> area with coordinates of 90171 N to 90401 N and 75019 E to 76033 E. (Padmanabhan *et al.*, 2001). ~~The majority~~ Most of the Kuttanad region, which is made up of reclaimed land from Vembanad lake, is located between 0.6 and 2.1 metres below sea level (Vijayan, 2016). ~~In Kerala's Kuttanad region, there is~~ Kerala's Kuttanad region has an ecological scenario known as the tidal wet land ecosystem coupled with deep water status (Vijayan, 2016). The entire wetland is split into three parts: the Kari lands (a marshy area with black peat soil that is highly acidic), the shallow reclaimed lands known as Karappadams, and the Kayal land (land in backwaters) (Vijayan, 2016 and Padmaja *et al.*, 1994). The Kuttanad region's paddy field is organized into contiguous blocks known as "Padasekharams," which are delimited by canals and natural partitions and range in size from 1 to 1000 ha (Thomas, 2002).

### Climate

Due to the presence of four rivers, primarily Manimala, Meenachil, Achencovil, and Pampa, the cultivable region of Kuttanad endures numerous natural calamities, such as heavy flooding during the South West monsoon period from July to August (Vijayan, 2016). Saline water incursion at high tide is also ~~common-typical~~ (Balchand, 1983). Floods transport alluvial materials from the Western Ghats that are rich in nutrients and deposit them along rivers in low-lying areas. Rice agriculture, especially deep water rice, is very suitable in the tidal wetlands ~~that form as a result of~~ resulting from floods and tidal intrusions (Richharia and Govindaswamy, 1990). Growing rice in locations where there is 50 cm of flooding for more than a month during the growing season is known as deep water rice or floating rice (Catling, 1992).

## Soil

Throughout the soil of Kerala's Kuttanad region, sand and clay are mixed in various amounts (Vijayan, 2016). The extremely fertile pokkali fields are also periodically submerged in water (Jayan & Sathyanathan, 2010; Pervez et al. 2017). The pokkali fields are exceptionally fruitful due to the daily tidal inflows and outflows ~~as well as~~ and the intense microbiological activity brought on by the abundance of organic matter (decomposed aquatic weed mass and paddy stubbles) (Shylaraj & Sasidharan, 2005).

## Pokkali rice cultivation Peculiarities

The Pokkali style of rice farming in Kerala's acidic, saline soil is a distinctive way to grow rice (Vijayan, 2016). It is an integrated farming approach that doesn't interfere with the environment's normal ecological processes (Kerala State Biodiversity Board, 2009). A historic salt-tolerant type of rice called Pokkali is cultivated from June to November along Kerala's coast (Ramaiah *et al.*, 2020). ~~In general, paddy cultivation is done during the low salinity phase, and~~ Paddy cultivation is generally done during the low salinity phase, while prawn cultivation is done during the high salinity phase (Shylaraj and Sasidharan, 2005). It is an organic rice production approach with lower costs than commercial rice farming because chemical fertilizers and pesticides are not used (Vijayan, 2016). This traditional way of growing rice doesn't use artificial fertilizers or pesticides on the crop (Shylaraj & Sasidharan, 2005 ~~and~~ Sathiadhas, 1989). With this technique, a single ~~crop of rice~~ rice crop is harvested on mounds during the low salinity part of the production cycle (June to mid-October), with prawn aquaculture taking place during the high salinity phase (November to April). The soggy, waterlogged fields don't require ~~the utilization of~~ labour-saving heavy machinery or manure addition (Nambiar & Raveendran, 2009). According to the custom, the variety has been given a Geographical Indicator (GI) tag (Pavithran, 2014). Pokkali rice production is an environmentally benign method of growing rice because it doesn't result in biodiversity loss, overuse of natural resources, or coastline deterioration (Paimpillil, 2007). Despite this, pokkali farmers only produce 2000 kg of rice ~~on average~~ per hectare, making it slightly unprofitable to grow rice here (Vijayan, 2016). Consequently, to maintain rice farming in the pokkali areas of Kerala, it is crucial to develop suitable varieties with high yield potential and the capacity to operate well under ~~the~~ acidic saline conditions with little to no external inputs.

## Rice varieties in Pokkali system

Both wild and cultivated rice in the state have a wide variety of germplasm as a result of the ecological circumstances there (Sharma, 1991). Pokkali, a native rice type with a 120-day growing season, is used by farmers (Kumari, 2007). This species, which can withstand salt and flooding, can reach a height of 1.5 meters (Vijayan, 2016). Chettivirippu, Vyttila 1, Vyttila 2, Vyttila 3, and Vyttila 4 are more types (KAU, 2002). ~~The~~ However, the export potential of the four pokkali rice cultivars (VTL 1 to VTL 4) that were made available by the Vyttila Rice Research Station is minimal. In particular, the red bold grain type, preferred by the local consumers, is not considered ~~a desirable trait~~ desirable by customers elsewhere (Vijayan, 2016). A potential "Mahsuri" mutant named VTL 5 has numerous abiotic stress tolerances, including salt, acidity, and submersion. It is a tall, medium-duration, high yielding kind of rice with white kernels that is also excellent for cooking (Vijayan, 2016).

### **Mound preparation**

In the pokkali system of rice growing, land preparation begins in May. (Vijayan, 2016). Bunds are raised first, then mounds with a 1 m<sup>2</sup> foundation and 50 cm height are prepared (KAU, 2002). By restricting ~~the flow of water~~ water flow into the fields, the mounds are allowed to dry up (Vijayan, 2016).

### **Sowing**

Sprouting seeds are sown on Pokkali field's mounds (Paimpillil, 2007). The seeds are sprouted ~~by placing them in fresh water for 12 to 15 hours while being in fresh water for 12 to 15 hours~~ while securely wrapped in coconut leaves with a layer of banana or teak leaves inside (Thomas, 2002 and Paimpillil, 2007; Pervez et al. 2021).

### **Dismantling**

The mounds are levelled a month after seeding. In the field, the ~~seedlings on the mound~~ mound seedlings are dispersed evenly (Vijayan, 2016). The ~~rapid growth and field establishment of the seedlings~~ seedlings' rapid growth and field establishment makes them more resilient to the ensuing floods. Because the prevalence of pests and diseases is below the threshold level, manuring and plant protection practices are not required ~~for the growth of~~ to grow Pokkali rice, resulting in the production of natural, organic rice (Jayan & Sathyanathan, 2010).

### **Harvesting**

Except for the erect panicles, which are harvested, the 1.5 to 2 m tall rice plant bends and collapses when it reaches maturity (Nambiar & Raveendran, 2009). The remaining stalks are left to rot in the water, where they subsequently provide prawn feed. Just 2000 kg ha<sup>-1</sup> of rice is often harvested by pokkali farmers, making it slightly unprofitable to grow rice here (Shylaraj & Sasidharan, 2005). Traditional rice varieties are declining for several reasons, including their low production, very ~~lengthy~~ long growing seasons, lack of price premiums for some types, and relatively lengthy cooking times (NICRA, 2019 and Xie, 2011). According to Jayan and Sathyanathan (2010), lodging and associated damages brought on by fish, tortoises, and rodents result in a loss of between 40 and 50 per cent of the potential yield. In addition to these losses

and the challenges associated with harvesting paddy, clearing fields for the subsequent selective stocking of prawns is ~~a challenge~~ challenging in these circumstances (Shylaraj *et al.*, 2006).

### **Prawn cultivation**

Typically, prawn farming ~~takes place~~ occurs from November to April (Vijayan, 2016). The backwaters and canals in this region become salted after the monsoon season, and a lot of young prawns enter them (Paimpillil, 2007). Sluice gates, or water canals with a head at their gate, are used to direct young prawns into pokkali fields. They are unable to leave due to these gates. In Pokkali field, this is known as prawn culture filtering (Thomas, 2002). In order to maintain the entry and outflow of water during the farming season, sluice gates are crucial in the prawn farming industry. They are kept up in a way that maximizes the amount of saline water that enters the field and prevents prawns from escaping (Vijayan, 2016). Prawn culture uses the leftover residue from rice cultivation as natural feed. While prawn excreta serve as a natural fertilizer for rice cultivation, prawns feeding on rice cultivation stubbles helps to minimize the occurrence of illness in rice to a ~~minimal~~ minimum (Vijayan, 2016).

### **Harvesting of Prawn cultivation**

The two main ~~species of cultivated prawn~~ cultivated prawn species are *Penaeus monodon* (Tiger prawn) and *Penaeus indicus* (White prawn). Prawns are typically collected in mid-April, especially two to three days before new full moon days, as tidal activity affects their movement (Vijayan, 2016).

### **~~Prospective~~ The prospective aspect of Pokkali rice cultivation**

Prawns raised in the field ~~give the farmers a boost in~~ boost the farmers' income (Vijayan, 2016). In Northern Kerala, it's usual practice to use the fertile bottom muck of the land as manure for coconut plantations (Nambiar & Raveendran, 2009). ~~Saline~~ In addition, saline-resistant cultivars and methods can help farmers prepare for the effects of climate change. Climate change is expected to result in sea level rise, which can increase sea-water intrusion in more coastal locations and cause agricultural setbacks that can be avoided by using these types (Gopi & Manjula, 2018). According to estimates, this state needs between 3.5 and 4.0 million tonnes of rice annually ~~-~~.

Nevertheless, Kerala only generates a quarter of this quantity (Krishnankutty *et al.*, 2021). A decrease in the area under rice cultivation has resulted in a growing annual shortfall in rice production. In Kerala, where rice farming has been ~~practiced for centuries, a severe issue has arisen due to the widespread conversion of paddy lands for other crops~~ for centuries, a severe issue has arisen due to the widespread conversion of paddy lands for other crops or for residential purposes (Kumari, 2007; Pervez *et al.* 2022). By erecting barriers to the entry of brackish water from the lakes and oceans and directing the flow of fresh water from the river system to these fields, the reclamation schemes were primarily designed to increase the amount of area under the rice crop and increase the output of the current fields (Korakandy, 2005). The primary reason for the decline in the adoption of Pokkali rice growing practices is the lack of farm labour, particularly for harvesting (Jayan & Sathyanathan, 2010).

## Conclusion

The shrinking area under rice cultivation and the widespread conversion of paddy lands for other crops or for residential uses, ~~there is~~ has a severe problem with meeting the demand and supply of rice within the state. ~~In order to~~ To boost the state's rice production, attention must be paid to traditional rice growing techniques, including both wild and cultivable kinds. The Pokkali system of rice cultivation is a traditional ~~technique for growing rice~~ rice-growing technique that systematically combines prawn farming with the saline-tolerant wild rice type known as Pokkali. ~~In general, paddy cultivation is done during the low salinity phase, and~~ Paddy cultivation is generally done during the low salinity phase, while prawn cultivation is done during the high salinity phase. Harvesting ~~is carried out when the rice crop reaches to~~ occurs when the rice crop reaches a height of 1.5 to 2 m with the help of boats. The remaining stalks are left to rot in the water where they subsequently acts as prawn feed whereas prawn excreta as a natural fertilizer for rice cultivation. - Pokkali rice production is an environmentally benign method of growing rice because it doesn't result in biodiversity loss, overuse of natural resources, or coastline deterioration Despite this, the primary reason for the decline in the adoption of Pokkali rice growing practices is the lack of farm labor, particularly for harvesting.

## Reference

- Balchand, A. N. 1983. Kuttand: A case study on environmental consequences of water resources mismanagement. *Water Int.* 8(1): 35-41p.
- Catling, H.D. 1992. Rice in deep water. IRRI [International Rice Research Institute]. Macmillan Press, London, 213-243p.
- Gopi, G. and Manjula, M. 2018. Speciality rice biodiversity of Kerala: Need for incentivizing conservation in the era of changing climate. *Current Science.* 114(5):997- 1006p.
- Jayan, P. R., and Sathyanathan, N. 2010. Overview of farming practices in the water-logged areas of Kerala, India. *Int. J. Agric. & Biol. Eng.*, 3(4): 25-30p.
- KAU [Kerala Agricultural University]. 2002. Package of Practices Recommendations, Thrissur. 278p.
- Kerala State Biodiversity Board. 2009. Available: <http://www.keralabiodiversity.org/pdf/policy.pdf>.
- Korakandy, R. 2005. Coastal Zone Management. Gyan Publishing House, 585 p.
- Krishnankutty, J. Blakeney, M. Raju, R.K. Siddique, K.H.M. 2021. Sustainability of Traditional Rice Cultivation in Kerala, India—A Socio-Economic Analysis. *Sustainability*, 13, 980p.
- Kumar, S. L. 2007. Rice varieties in Kerala. *J. of tropical Agriculture.* 27(12);16-18p.
- Kumar, B. and Kunham, T. K. 2021. Ecological and historical perspectives of rice cultivation in Kerala: a synthesis. *Oryza.* 58(2): 241-261p.
- Kumari, L.S. 2012. Status paper on rice in Kerala. Available at <http://www.rkmp.co.in/>

Kumari, S.L. Status. 2007. Paper on Rice. Rice Knowledge Management Portal (RKMP) Directorate of Rice Research, Rajendranagar, Hyderabad. Available online: <http://www.rkmp.co.in/sites/default/files/ris/rice-state-wise/Status%20Paper%20Kerala>

Maliyakkal, G. and Thomson, K. 2008. Water sharing institution in complex coastal commons: A study of Cochin Estuary in Kerala in India, 12<sup>th</sup> Biennial Conference of the International Association for the Study of Commons, England

Nambiar, G. R. and Raveendran, K. 2009. Exploration of Untapped Potentiality of Coastal Paddy Fields of Kerala (India) - A Case Study. *Middle-East J. of Scientific Research*, 4(1): 44-47p.

NICRA [National Innovations on Climate Resilient Agriculture]. 2019. Available online: <http://www.nicra-icar.in/nicrarevised>

Padmaja, P. Kumari, V. L. G. Nair, H.K. Chinnamma, N. P. Sasidharan, N. K. Rajan, K.C. 1994. A glimpse to problems soils of Kerala. Kerala Agricultural University, Vellanikkara, Thrissur. 116p.

Padmanabhan, P. G., Narayanan, N.C. and Padmakumar, K. G. 2001. Economic viability of an integrated and Sustainable Resource Use Model for Kuttanad. Discussion paper on Kerala Research Programme on Local Level Development (KRPLLD)

Paimpilli, J. S. 2007. Ecofriendly prawn farming with Pokkali cultivation: Role of women and children in coastal resource management and poverty reduction. Proceedings of Pacem in Mauritius XXXIII Conference, Malta, 274 – 280p.

Pavithran, K. 2014. Pokkali Rice Production under Geographical Indication Protection: The Attitude of Farmers. *Journal of Intellectual Property Rights*, 19 (1): 24-29p.

Pervez, A. K. M. K., Gao, Q., Zeng, Y., & Uddin, M. E. (2017). Hybrid rice: Bangladesh's failure and China's success. *Journal of Agriculture and Ecology Research International*, 10(1), 1-10.

Pervez, A. K. M., Riar, T. S., & Sheikh, M. (2021). Integrated farming systems: A review of farmers friendly approaches. *Asian Journal of Agricultural Extension, Economics & Sociology*, 39(4), 88-99.

Pervez, A. K., Shah, A. A., Uddin, M. E., Sarker, M. N. I., & Islam, M. M. (2022). Risks affecting hybrid rice yield in Bangladesh: Two-tier approach of appraisal and management. *Journal of Agricultural Sciences–Sri Lanka*, 17(1), 211-227.

Ramaiah, M. H. S. Jagadheesh, T. Dey, P. Jambagi, S. M. Kumari, M. V. Karaba, N. Ravishankar, K. V. N. Ravikanth, G., and Shanker, R. U. 2020. An endophyte from salt-adapted Pokkali rice confers salt-tolerance to a salt-sensitive rice variety and targets a unique pattern of genes in its new host. *Nature*. 10: 3237p.

Richharia, R.; Govindasamy, S. 1990.- Rices of India; Academy of Development Science: Karjat, India. *Science and Research*, 7(4): 719–730p.

Sathiadhas, R., Panikkar, K.K.P., Satyavan U.K. & Jacob, T. (1989). Economic valuation of Paddy - Prawn Integrated Farming in Kerala. *Sea Food Export Journal*, 9-21p.

Sharma, R.D. 1991. *The Story of Rice*; National Book Trust: New Delhi, India, 68p.

Shylaraj, K. S. and Sasidharan, N. K. 2005. VTL5: High yielding salinity tolerant rice variety for the coastal saline ecosystems of Kerala. *J. of Tropical Agriculture*. 43 (1): 25 -28p.

Shylaraj K S, Sasidharan N K, Sreekumaran V. 2006. VTL 6: A semi-tall, non-lodging, and high yielding rice (*Oryza sativa* L.) variety for the coastal saline zones of Kerala. *Journal of Tropical Agriculture*, 44 (2): 48–51p.

Suchitra, M. and Venugopal. 2005. In Troubled waters. Agriculture. Available: [www.questfeatures.org/articles/pokkali.html](http://www.questfeatures.org/articles/pokkali.html).

Thomas, P. M. 2002. Problems and prospects of paddy cultivation in Kuttanad region. Discussion paper on Kerala Research Programme on Local Level Development (KRPLLD)

Vijayan, R. 2016. Pokkali Rice Cultivation in Kerala. *Agriculture Update*.11(3):329-333p.

Xie, J.; Wu, X.; Tang, J.-J.; Zhang, J.-E.; Luo, S.-M.; Chen, X. 2011. Conservation of Traditional Rice Varieties in a Globally Important Agricultural Heritage System (GIAHS): Rice-Fish Co-Culture. *Agric. Sci. China*, 10, 754–761p.