

***Original Research Article***

**Response of Biofertilizers and Foliar application of Zinc on Growth and Yield of Lentil (*Lens culinaris* L.)**

Formatted: Font: Italic

**ABSTRACT**

A field experiment was conducted during *Rabi* 2022 at Crop Research Farm, Department of Agronomy, Sam Higginbottom Institute of Agricultural Sciences, Prayagraj, UP, during the *Rabi* season of 2022. The experiment consisted of two variables with three levels i.e, biofertilizer (*Rhizobium spp*, *Pseudomonas fluorescens*, *Rhizobium spp* + *Pseudomonas fluorescens*) and foliar application of zinc (0.25%, 0.50%, 0.75%) at pre flowering and pod development stage. The soil had a sandy loam texture, with a pH of 7.2, organic carbon (0.72%), available N (178.48 kg/ha), available P (27.80kg/ha) and available K (233.24 kg/ha). The experiment conducted in randomized block design with ten treatments replicated thrice. Application of biofertilizers and zinc significantly influenced the growth parameters, yield attributing characters and yield. The results revealed that treatment eight (*Rhizobium spp* + *Pseudomonas fluorescens* + Zinc 0.50%) recorded significantly highest plant height (43.29 cm), dry weight (35.63 g), number of pods/plant (111.33), number of seeds/pod (2.27), seed yield (1530.94 kg/ha) and stover yield (3020.00 kg/ha) in Lentil.

Formatted: Font: Italic

Formatted: Font: Italic

**Key words:** *Biofertilizer, Rhizobium spp, Pseudomonas fluorescens, Growth, Yield, Rabi, Lentil.*

**Comment [AC1]:** The abstract must include the problem and the justification of the work. In the methodology, include the variables and statistical analyzes performed. Results are presented but not their impact for the study area.

**Comment [AC2]:** Avoid repeating words from the title.

## Introduction

Lentil (*Lens culinaris*) is an important annual leguminous crop which is locally called Masoor belongs to the family Fabaceae. It is one of the most nutritious and also known as poor man's meat because of its rich protein content and high concentration of essential amino acid as isoleucine and lysine (Nasar and Shah, 2017). Lentils contain important components of human nutrition, such as 25% protein, 1.1% fat, 59% carbohydrate, and is also rich in soluble and insoluble dietary fiber. It is also rich in calcium, iron, phosphorus, potassium, zinc, magnesium and vitamins like niacin, riboflavin, thymine and ascorbic acid (Fatima et al., 2018). Lentils are naturally gluten-free. Their exceptionally low glycemic index values and resistant starch content make them suitable for a diabetic diet. Uttar Pradesh is the second largest growing state of lentil after Madhya Pradesh in the country. Lentil production of U.P was 0.47 million tonnes from an area of 0.47 million hectares and productivity of 988 kg/ha in 2020-2021 (Anonymous, 2021). It is one of the principal pulse crops cultivated in semi-arid region of the world, particularly in India sub-continent and the dry areas of Middle East (Malik, 2005).

Bio-fertilizers are living microorganisms of bacterial, fungal and algal origin. Biofertilizers fix atmospheric nitrogen in the soil and root nodules of legume crops and make it available to the plant. They solubilize the insoluble forms of phosphates into available forms. They produce hormones and antimetabolites which promote root growth. When applied to seed or soil, biofertilizers increase the availability of nutrients and improve the yield by 10 to 25% without adversely affecting the soil and environment (Kumar and Chandra, 2008). Production of lentil enhanced by optimum use of nutrients. <sup>34</sup>freewwes Inoculation with *Rhizobium* influence growth and productivity of lentil. The efficiency of *Rhizobium* improved by co-inoculation with plant growth promoting rhizobacteria (PGPR). It shows an important role in the sustainable agriculture industry CITE-

The use of PGPR has been proven to be an environmentally sound way of increasing crop yields by facilitating plant growth through either a direct or indirect mechanism. The mechanisms of PGPR include regulating hormonal and nutritional balance inducing resistance against plant pathogens, and solubilizing nutrients for easy uptake by plants. It shows synergistic and antagonistic interactions with microorganisms with in the rhizosphere, which indirectly boosts

Comment [AC3]: Include economic importance internationally, nationally and regionally. Updated information

Comment [AC4]: CITE.

Comment [AC5]: Most current references.

plant growth rate. There are many bacteria species that act as PGPR (Singh *et al.*, 2016). Use of plant growth promoting rhizobacteria is often associated with increased rates of plant growth, development and yield. Co-inoculation with *Rhizobium* and PGPR is even more effective for improving nodulation and growth of legumes (Biswas *et al.*, 2012).

Foliar application of micronutrients helps in the rapid translocation when compared to soil application which is very pertinent in mitigating stress in plants. Zinc role in a plant is either as a metal constituent in enzymes or as a functional co-factor of number of enzymatic reactions. Zinc is required for synthesis of tryptophan, which in turn is precursor for synthesis of IAA. After flowering high levels of zinc in plant will enhance cell differentiation (Anil *et al.*, 2013). It plays vital role in the biosynthesis of plant hormone like IAA, protein and nucleic acid which helps in utilization of nitrogen and phosphorus in plants (Saha *et al.*, 2018). At flowering, Zinc plays a greater role during reproductive phase especially during fertilization. Most of the zinc is diverted to seed only (Singh *et al.*, 2013).

Keeping the above points in view, the present investigation was carried to find out **“Response of Biofertilizers and Foliar application of Zinc on Growth and Yield of Lentil (*Lens culinaris*)”**.

**Comment [AC6]:** No antecedent studies are presented that justify the performance of the work.

**Comment [AC7]:** At the end of the introduction, it is not only to include the objective but also the impact at the biological and agronomic level that the research has.

## Materials and Methods

The experiment was conducted during *Rabi* season of 2022, at Crop research farm of department of Agronomy at Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj which is located 98 m altitude above the mean sea level. The soil in the experimental area was sandy loam texture with pH (7.2), organic carbon (0.72%), available N (178.48 kg/ha), available P (27.80 kg/ha) and available K (233.24 kg/ha) were determined by Jackson's method, Subbaiah, and Asija's method, Olsen's method, Flame photometer method, respectively.

The experiment was laid out in randomized block design (RBD) with ten treatments each replicated thrice. The experiment comprised of three biofertilizer treatments and three levels of zinc when applied in combinations as follows, T<sub>1</sub>: *Rhizobium spp*+ Zinc (0.25%), T<sub>2</sub>: *Rhizobium spp*+ Zinc (0.50%), T<sub>3</sub>: *Rhizobium spp* + Zinc (0.75%), T<sub>4</sub>: *Pseudomonas fluorescens* + Zinc (0.25%), T<sub>5</sub>: *Pseudomonas fluorescens* + Zinc (0.50%), T<sub>6</sub>: *Pseudomonas fluorescens*+ Zinc (0.75%), T<sub>7</sub>: *Rhizobium spp* + *Pseudomonas fluorescens* + Zinc (0.25%), T<sub>8</sub>: *Rhizobium spp* + *Pseudomonas fluorescens* + Zinc (0.50%), T<sub>9</sub>: *Rhizobium spp* + *Pseudomonas fluorescens* + Zinc (0.75%), T<sub>10</sub>: Control (NPK 20-40-20 kg/ha).

The variety of Lentil is "PL 406". The crop was sown with pre-sowing irrigation. The line-to-line spacing was kept at 30 cm spacing with a seed rate of 40 kg/ha. A distance of 10 cm was maintained between plant to plant. Hand weeding was done twice at 30 and 50 days after sowing. Nutrients particularly, nitrogen, phosphorus, and potassium were applied as basal dose. Foliar application of zinc was done at pre flowering stage and pod development stage. The observations on five randomly selected plants from each treatment recorded on different growth parameters at 20,40,60,80 and 100 days after sowing viz. plant height (cm), dry weight (g/plant), number of pods/plant, number of seeds/pod, seed yield and stover yield. The data collected was computed and statistical analysis by analysis of variance method (Gomez and Gomez, 1984).

**Comment [AC8]:** It must describe in detail how the variables were measured.

## Result and Discussion

### Growth attributes

At 100 DAS, significantly highest plant height (43.29 cm), dry weight (35.63 g) was recorded in treatment-8 with the combined application of *Rhizobium spp.*, *Pseudomonas fluorescens* along with zinc (0.50%). Dual seed inoculation with *rhizobium* and PGPR gave higher plant height as it has the ability to fix atmospheric nitrogen in rhizosphere of leguminous plants through the process of N fixation. The results are in conformity with those reported by (Singh *et al.*, 2016). Zinc is an important element for the synthesis of tryptophan, which is the pioneer for the synthesis of IAA (Indole acetic acid), a growth hormone, involved in stem elongation (Swargiary *et al.*, 2021). Increase in plant height might be attributed to the fact that the better nourishment causes accelerated rate of photosynthesis, assimilation, cell division and vegetative growth (Tiwar *et al.*, 2018). The favorable effect of plant growth regulator and zinc might influence the metabolism, photosynthetic pigments and activity of enzymes which in turn helps to increase in the vegetative growth. An enhanced growth and higher dry matter accumulation due to application of zinc, produce growth hormones and precursor of auxins (Mounika *et al.*, 2022).

### Yield attributes

According to yield attributes data that was collected and analyzed at harvest, significantly higher number of pods/plant (111.33), seeds/pod (2.27) were recorded in treatment-8 with the combined application of *Rhizobium spp.*, *Pseudomonas fluorescens* along with zinc (0.50%). The combination of *Rhizobium* and Zinc plays an important role in increasing the crop production. It increases microorganisms that accelerate microbial process which in turn augment the extent availability of nutrient in the form which is easily assimilated by the plant. Zinc helps in production of auxin which resulted in more number of pods/plant (Mounika *et al.*, 2022). Micro nutrients help in utilization of phosphorus and nitrogen in flower retention, pollen tube development, seed formation and seed setting through the translocation of metabolites from source to sink (Shinde *et al.*, 2017). Zinc improved translocation of photosynthates towards reproductive system and thereby enhancing the yield of the crop. Better photosynthetic activity

**Comment [AC9]:** It is important to discuss to what extent it is better to have tall plants, as this would make agronomic work difficult. Comparative studies must present context. The tables must be related in the text.

also may have resulted in better translocation of photosynthates from source to sink due to less crop competition between the plants which might have led to higher yield attributes (Swargiary *et al.*, 2021). Seeds per pod were also increased with zinc application due to the role of zinc in seed setting (Praveena *et al.*, 2018).

**Comment [AC10]:** Take into account previous observations, and that comparative studies are current.

### Yield (kg/ha)

After evaluated the data recorded post harvesting of crop show that significantly maximum seed yield (1530.94 kg/ha) and stover yield (3020.00 kg/ha) were recorded with the combined application of *Rhizobium spp*, *Pseudomonas fluorescens* along with zinc (0.50%). The beneficial effect of PGPR and *Rhizobium* had probably induced the synthesis of growth promoting substances which could stimulate growth and elongation, nitrogen fixation and crop yield. The increase in seed yield have been attributed to interaction with various soil borne pathogens, production and release of secondary metabolites for plant growth, or increased uptake of nutrients (Biswas *et al.*, 2012). *Pseudomonas fluorescens* apart from possessing phosphorus solubilizing ability improving symbiotic efficiency, growth and yield of lentil (Khanna *et al.*, 2011). Foliar spray of zinc leads to optimum availability of nutrients for luxurious crop growth and efficient partitioning of assimilates from source to sink (Lakshmi *et al.*, 2017). Zinc application improve stover yield due to its direct influence on auxin production which in turn enhanced the elongation processes of plant development (Rajana *et al.*, 2017).

**Comment [AC11]:** The results are discussed only from the biological point of view, it is recommended to do it from the agronomic point of view.

### CONCLUSION

It can be concluded that combined application of *rhizobium spp* and *Pseudomonas fluorescens* at 20 g/kg seed along with foliar application of zinc at 0.05% (treatment T<sub>8</sub>) recorded higher growth parameters, improved yield attributes, maximum seed yield, and stover yield in lentil crop.

**Comment [AC12]:** The conclusions should not be a abstract of the main results found, but show their impact.

**Ethical paper:** This article does not contain any studies with human participants or animals performed by any of the authors.

## REFERENCES

Anonymous. (2021). Directorate of Economics and Statistics. Ministry of Agriculture and Farmers Welfare, Department of Agriculture, Cooperation & Farmers Welfare, Government of India. *Agricultural Statistics at a glance data 2021*.

Biswas, P.K., Bhowmick, M.K., Malik, G.C, Banerjee, M., Ghosh, G.K. and Mitra, S.R. (2012). Synergistic Effect of Rhizobium Inoculation with Co-inoculants on Growth and Yield of Lentil (*Lens culinaris Medikus*). *International Journal of Bio-resource and Stress Management*. 3(1): 000-000.

Formatted: Highlight

Fatima, K., Ganie, S.A., Kumar, Y., Masoodi, T.H. and Shah, A. (2018). Effect of Organic and Inorganic Fertilizer Does on Growth and Yield Lentil under Cold Arid Conditions of Ladakh. *International Journal of Current Microbiology and Applied Sciences*. 7(11): 1449-1455.

Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedures for Agricultural Research*. 2<sup>nd</sup> Edition, John Wiley and Sons, New York, 680 p.

Khanna, V. and Sharma, P. (2011). Potential for enhancing lentil (*Lens culinaris*) productivity by co-inoculation with PSB, plant growth promoting rhizobacteria and Rhizobium. *Indian Journal of Agricultural Sciences*. 81(10): 932-4.

Formatted: Highlight

Kumar, R. and Chandra, R. (2008). Influence of PGPR and PSB on *Rhizobium leguminosarum* Bv. Viciae Strain Competition and Symbiotic Performance in Lentil. *World Journal of Agricultural Sciences*. 4(3): 297-301.

Formatted: Highlight

Lakshmi, E.J., Babu, P.V.R., Reddy, G.P., Maheswari, P.U. and Reddy, A.P.K. (2017). Effect of foliar application of secondary nutrients and zinc on growth and yield of Blackgram. *International Journal of Chemical Studies*. 5(6): 944-947.

Malik, R. (2005). Genetic divergence analysis in lentil (*Lens culinaris Medik*). M.Sc. Thesis, Department of Agricultural Botany, Chaudhary Charan Singh University, Meerut (U.P), India. 1pp.

Mounika, S., Mehra, B., Sathvik, D. and Chandan, N. (2022). Influence of Bio Fertilizers and Zinc on Growth and Yield of Lentil (*Lens culinaris L.*). *International Journal of Environment and Climate Change*. 12(10): 788-793.

Nasar, J. and Shah, Z. (2017). Effect of Iron and Molybdenum on Yield and Nodulation of Lentil. *Journal of Agricultural and Biological Science*. 12(11): 1190-6145.

Formatted: Highlight

- Praveena, R., Ghosh, G. and Singh, V. (2018). Effect of Foliar Spray of Boron and Different Zinc Levels on Growth and Yield of Green gram (*Vigna radiata*). *International Journal of Current Microbiology and Applied Sciences*. **7**(8): 1422-1428.
- Saha, G., Ghosh, M., Nath, R., Gunri, S.K., Roy, K. and Saha, B. (2018). Effect of Boron and Zinc on growth, yield and economics of lentil (*Lens culinaris*) in New Alluvial Zone of West Bengal. *Indian Journal of Agronomy*. **63**(3): 391-393.
- Shinde, P., Doddagoudar, S.R. and Vasudevan, S. N. (2017). Influence of seed polymer coating with micronutrients and foliar spray on seed yield of chickpea (*Cicer arietinum* L.). *Legume Research*. **40**(4): 704-709.
- Singh, A.K. and Bhatt, B.P. (2013). Effect of foliar application of zinc on growth and seed yield of late-sown lentil (*Lens culinaris*). *Indian Journal of Agricultural Sciences*. **83**(6): 622-6.
- Singh, N., Singh, G. and Khanna, V. (2016). Growth of Lentil (*Lens culinaris* Medikus) as influenced by phosphorus, rhizobium and plant growth promoting rhizobacteria. *Indian Journal of Agriculture Research*. **50**(6): 567-572.
- Swargiary, S., Umesha, C. and Dwivedi, N. (2021). Influence of Spacing and Zinc Levels on Growth and Yield of Lentil (*Lens culinaris*). *Biological forum*. **13**(3): 114-117.
- Tiwari, A.K., Prakash, V., Ahmad, A. and Singh, R.P. (2018). Effect of Biofertilizers and Micronutrients on Nutrient Uptake, Growth, Yield and Yield Attributes of Lentil (*Lens culinaris* L.). *International Journal of Current Microbiology and Applied Sciences*. **7**(2): 3269-327.

Formatted: Highlight

Formatted: Highlight

**Table 1. Response of Biofertilizers and Foliar application of Zinc on Growth attributes of Lentil.**

Comment [AC13]: Complement with the place where the work was carried out.

S.No	Treatment combinations	100 DAS	100 DAS	40-60 DAS
		Plant height (cm)	Dry weight (g/plant)	CGR (g/m <sup>2</sup> /day)
1.	<i>Rhizobium</i> spp +Zinc -0.25%	32.08	30.27	10.30
2.	<i>Rhizobium</i> spp +Zinc -0.50%	33.14	29.87	9.68
3.	<i>Rhizobium</i> spp +Zinc -0.75%	34.37	30.43	10.45
4.	<i>Pseudomonasfluorescens</i> +Zinc -0.25%	35.27	30.57	11.69
5.	<i>Pseudomonasfluorescens</i> +Zinc-0.50%	36.62	32.75	12.65
6.	<i>Pseudomonasfluorescens</i> +Zinc-0.75%	37.05	31.00	12.47
7.	<i>Rhizobium</i> spp + <i>Pseudomonasfluorescens</i> +Zinc-0.25%	39.32	33.24	15.18
8.	<i>Rhizobium</i> spp + <i>Pseudomonasfluorescens</i> +Zinc -0.50%	43.29	35.63	17.53
9.	<i>Rhizobium</i> spp + <i>Pseudomonasfluorescens</i> +Zinc -0.75%	42.69	34.95	15.51
10.	Control(20-40-20NPKkg/ha)	30.94	28.20	9.38
	F-test	S	S	S
	SEm(±)	0.22	1.08	1.03
	CD(p=0.05)	0.66	3.21	3.08

**Table 2. Response of Biofertilizers and Foliar application of Zinc on Growth attributes of Lentil**

S.No	Treatment combinations	Number of pods/ plant	Number of seeds/pod	Seed yield (kg/ha)	Stover yield (kg/ha)
1.	<i>Rhizobium</i> spp + Zinc -0.25%	103.00	1.73	1025.37	2076.67
2.	<i>Rhizobium</i> spp + Zinc -0.50%	103.67	1.73	1038.44	2216.67
3.	<i>Rhizobium</i> spp + Zinc -0.75%	104.00	1.80	1082.46	2346.67
4.	<i>Pseudomonas fluorescens</i> + Zinc -0.25%	105.33	1.93	1192.33	2463.33
5.	<i>Pseudomonas fluorescens</i> + Zinc -0.50%	105.67	2.00	1257.02	2570.00
6.	<i>Pseudomonas fluorescens</i> + Zinc -0.75%	107.00	2.07	1309.93	2676.67
7.	<i>Rhizobium</i> spp + <i>Pseudomonas fluorescens</i> + Zinc -0.25%	107.33	2.13	1378.94	2843.33
8.	<i>Rhizobium</i> spp + <i>Pseudomonas fluorescens</i> + Zinc -0.50%	111.33	2.27	1530.94	3020.00
9.	<i>Rhizobium</i> spp + <i>Pseudomonas fluorescens</i> + Zinc -0.75%	109.33	2.20	1451.81	2920.00
10.	Control (20-40-20 NPK kg/ha)	101.00	1.73	997.02	2036.67
	F-test	S	S	S	S
	SEm(±)	0.912	0.06	39.13	45.98
	CD(p=0.05)	2.710	0.17	116.27	136.61

UNDER PEER REVIEW

