

Economic factors and different growth phases of sweet corn (*Zea mays* L. var. *Saccharata*) in the South Gujarat area, India were affected by intra-row spacing and potassium levels

ABSTRACT

To determine how intra-row spacing and potassium levels affected sweet corn, a field experiment was conducted in Junagadh (Gujarat) during the Rabi season of 2016–17. (*Zea mays* L. var. *Saccharata*). Four levels of intra-row spacing (5, 10, 15 and 20 cm) and four potassium levels (0, 20, 40, and 60 kg K₂O/ha) were combined into sixteen treatment combinations. Three replications of a factorial randomised block design were used to set up the experiment. Based on the data of growth attributes significant and maximum recorded dry matter accumulation at (102.46 and 144.54 g plant⁻¹ at 60 DAS and harvest), absolute growth rate at (3.07 and 1.41 g day⁻¹ at 45-60 DAS and 60 DAS and harvest) and crop growth rate at (0.00275 and 0.00126 g m⁻²day⁻¹) under the treatment intra-row spacing (20 cm), respectively. According to data on growth attributes, the treatment (K₄) 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha resulted in significant and maximum dry matter accumulation at (96.93 and 142.00 g plant⁻¹ at 60 DAS and harvest), absolute growth rate at (3.03 and 1.39 g day⁻¹ at 45-60 DAS, and 60 DAS-harvest), and crop growth rate at (0.00272 and 0.00124 g m⁻²day⁻¹, respectively. Economic analysis showed that higher net returns and B: C ratio from sweet corn (Sweet-16) can be secured by sowing the crop at 20 cm intra-row spacing + application of 60 kg K₂O/ha.

Key words: *Economics, spacing, Gujarat, growth and potassium*

1. INTRODUCTION

Due to its classification as a C₄ type crop, maize (*Zea mays* L.) is often grown effectively throughout the year. Among the numerous forms of maize, sweet corn is particularly famous for the use of its green cobs all throughout the world. Popular vegetable sweet corn is in second in farm value and fourth in terms of commercial crops. The sweet corn is possible to boost agricultural revenue because of the rise in demand. The major consideration is to maintain stand density in order to increase cob yields. The form and size of plant-1's leaf area are determined by its spatial layout, which in turn affects how well it can absorb solar energy and how quickly its roots can develop and function. Only when plant population permits each plant to reach its

full natural potential can maximum production be anticipated. In order to get the best population density, inter- and intra-row spacing must be modified in connection to other agronomic parameters [7].

For maize, potassium (K) is a macronutrient because the plant absorbs a lot of it during the growth season. K serves as an activator for several enzymes and metabolic processes, including those involved in photosynthesis, protein synthesis, and starch production in grains, even though the plant does not use it as a building block for organic molecules. Potassium has a function in the flow of water, minerals, and carbohydrates inside the plant. It controls how stomata close and open, which affects how much water and gas are exchanged. Moreover, K is crucial for cell wall strength and cellulose formation. Strong cell walls that increase disease resistance and the capacity of the crop to keep firm, robust stalks are linked to high K fertility. For regulating disease incidence and stalk strength as corn output levels rise, it's critical to maintain a balance between nitrogen (N) and potassium (K) levels. The plant's ability to take N from the soil is constrained when K is a limiting factor, which has an effect on stalk strength, disease resistance, and grain output [11].

2. MATERIALS AND METHODS

The outcomes of a field experiment titled "Study of intra-row spacing and potassium levels on growth, yield, and quality of sweet corn (*Zea mays* L. var. *Saccharata*) under South Saurashtra conditions" carried out at the Farming System Research Centre, Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat during *Rabi* season of 2016-17.

2.1 Crop husbandry

Dry matter accumulation: Five plants were randomly selected from border lines of each experimental plot at 30, 45, 60 DAS and harvest. After chopping, plant samples were placed separately in perforated paper bags and oven dried at 65°C till a constant weight is obtained. Later, these were weighted and dry matter was expressed as g plant⁻¹. *Relative growth rate (RGR):* The values for relative growth rate were calculated for the stage between 30 DAS and 45 DAS, 45 DAS and 60 DAS and then between 60 DAS and at harvest with the help of following formula. $RGR \text{ g g}^{-1} \text{ day}^{-1} = (\log_e w_2 - \log_e w_1) / (t_2 - t_1)$, Where, $\log_e w_1 = \log_e$ of dry weight of plant at time interval t_1 . $\log_e w_2 = \log_e$ of dry weight of plant at time interval t_2 . *Crop growth rate*

(CGR): The values for crop growth rate were calculated for the stage between 30 DAS and 45 DAS, 45 DAS and 60 DAS and then between 60 DAS and harvest with the help of following formula: $CGR = 1/p (w_2-w_1)/(t_2-t_1) \text{ g m}^{-2} \text{ day}^{-1}$. Where, w_1 = weight of dry matter of plant at time t_1 . w_2 = weight of dry matter of plant at time t_2 . p = ground area (m^2)

Absolute growth rate: The values for absolute growth rate were calculated for the stage between 30 DAS and 45 DAS, 45 DAS and 60 DAS and then between 60 DAS and harvest with the help of following formula $AGR = (w_2-w_1)/(t_2-t_1) \text{ g day}^{-1}$. Where, w_1 = dry weight of plant at time t_1 . w_2 = dry weight of plant at time t_2 .

2.2 Crop economics

Cost of cultivation: The expenses incurred for all the routine operations from preparatory tillage to harvesting including threshing, cleaning as well as the cost of inputs viz. seeds, fertilizers, pesticides, irrigation etc. applied to each treatment were calculated on the basis of prevailing local charges and then cost of cultivation was calculated. *Gross returns:* The gross realization in terms of rupees per hectare was worked out separately for each treatment considering the green cob and green fodder yields from each treatment and local market prices. *Net returns:* The total cost of cultivation was deducted from the gross realization to work out the net income for each treatment combinations and was recorded accordingly. *Benefit: cost ratio (B: C):* The Benefit: Cost ratio (B: C) ratio was calculated with the help of following formula. $B: C = \text{Gross returns } (\text{₹/ha}) / \text{Total cost of cultivation } (\text{₹/ha})$

2.3 Crop statistical analysis

By using the appropriate analysis of variance as suggested by [2] the data was subjected to statistical analysis. The critical difference (CD) values were generated for each instance when the F values were determined to be significant at the 5% level of probability in order to compare the treatment means.

3. RESULTS AND DISCUSSION

3.1 Growth attributes

Different intra-row spacing levels had no discernible impact on the accumulation of dry matter at 30 and 45 DAS, according to an analysis of the data (Table 1). Nevertheless, intra-row spacing of 20 cm (S_4) recorded considerably greater dry matter accumulation at 60 DAS and harvest (102.46 g and 144.54 g, respectively), which was deemed statistically comparable to treatment S_3 (intra-row spacing of 15 cm).

Due to the consideration of various intra-row spacing values, absolute growth rate between 30-45 DAS was not significantly impacted. Nevertheless, during 45-60 DAS, intra-row spacing of 20 cm (S_4) recorded considerably the higher absolute growth rate (3.07 g/day) which was remained statistically at par with treatment S_3 (15 cm). Nevertheless, between 60 DAS-harvest and treatment S_4 (20 cm), which was discovered statistically at par with treatment S_3 , absolute growth rate continued to drop and was recognized to be greater (1.41 g/day) (15 cm). Crop growth is a well-established consequence of environmental interaction. In the current study, it was revealed that intra-row spacing had a significant impact on crop production for each plant, indirectly dictating the amount of rivalry among plants for different growth inputs as well as the availability of different growth nutrients to individual plants in the commodity.

Also, between 45 and 60 DAS, the treatment with a 20 cm intra-row spacing (S_4) recorded statistically equivalent results to treatment S_3 but with a much greater crop growth rate ($0.00275 \text{ g m}^{-2} \text{ day}^{-1}$) (15 cm). Nevertheless, between 60 DAS-harvest and treatment S_4 (20 cm), which was found statistically at par with treatment S_3 , the crop growth rate started to drop and was recorded higher ($0.00126 \text{ g m}^{-2} \text{ day}^{-1}$) (15 cm).

It is common knowledge that N, P, and K are key nutrients for crop growth and development. The greatest levels of N, P, and K in the crop's plant portion at the recommended intra-row spacing (S_4) of 20 cm may have aided in the promotion of plant development through active cell division and elongation. Under 20 cm of spacing, there appears to be a larger accumulation of photosynthates and ultimately a higher accumulation of dry matter by individual plants as a result of the enhanced nutritional state. In comparison to limited intra-spacing, higher values of growth characteristics were seen with broader intra-row spacing. The increase seen with greater intra-row separation may be related to a reduction in plant competition for nutrients and light under equidistant spatial arrangement [10]. The result is in close accordance with findings of [4, 8, 3 & 1].

Application of treatment K_4 ($60 \text{ K}_2\text{O} + 120 \text{ N}_2\text{O} + 60 \text{ P}_2\text{O}_5 \text{ kg/ha}$) resulted in considerably larger dry matter accumulation at 60 DAS and harvest than treatment K_3 ($40 \text{ K}_2\text{O} + 120 \text{ N}_2\text{O} + 60 \text{ P}_2\text{O}_5 \text{ kg/ha}$) (96.93 g and 142.00 g respectively). Significantly reduced levels of dry matter accumulation (73.11 g and 108.14 g) were observed at 60 DAS and harvest, respectively, with treatment K_1 ($0 \text{ K}_2\text{O} + 120 \text{ N}_2\text{O} + 60 \text{ P}_2\text{O}_5 \text{ kg/ha}$). The application of $60 \text{ K}_2\text{O} + 120 \text{ N}_2\text{O} + 60 \text{ P}_2\text{O}_5$ (K_4) between 45 and 60 DAS also recorded a considerably greater absolute growth rate

(3.03 g/day), which was statistically comparable to treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha). However, between 60 DAS-harvest and treatment K₄ (60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), which was found statistically at par with treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), the absolute growth rate continued to drop and was recorded higher (1.39 g/day). When treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), which was statistically comparable to treatment K₄, applied 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha, it considerably increased crop growth rate (0.00272 g m⁻² day⁻¹) between 45 and 60 DAS. Nevertheless, between 60 DAS-harvest and treatment K₄ (60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), which was found statistically at par with treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), the crop growth rate started to fall and was recorded higher (0.00124 g m⁻² day⁻¹).

Since potassium has a favourable influence on growth and enhances cell division and cell expansion, it has a beneficial effect on growth. Potassium's impact on the production of phytohormones plays a significant part in meristematic growth. Cytokinin, one of several plant hormones, is crucial for the development of tillers and buds. High spikelet fertility is a result of improved pollen germination in the florets due to potassium feeding. Such an increase may also be attributable to the root systems of plants receiving enough potassium from the soil, which increases photosynthesis and the production of metabolites and enzymes in plants [5]. According to [6] potassium boosts the plant's potential resistance to illnesses and insect pests. [12 & 9] both indicated that K had positive impacts on growth.

3.2 Economics

According to a review of the data (Table 2), treatment S₄ (intra-row spacing of 20 cm) considerably produced greater gross returns, which were ₹ 92734/ha, whereas treatment S₁ (intra-row spacing of ₹ 65575/ha) significantly produced lower gross returns (5 cm). Net return considerably was secured with an intra-row spacing of 20 cm (S₄), which stayed on the same bar as treatment S₃ (15 cm), and the lower net returns of ₹ 39719/ha were accumulated under S₁ (5 cm). Due to greater availability of nutrients, moisture, solar radiation, and room for growth and development, population maintenance at intra-row spacing of 20 cm (S₄) provided greatest net returns of ₹ 67408/ha and BCR 3.7.

Also, S₄ with an intra-row spacing of 20 cm acquired the greatest benefit cost ratio of 3.7, while S₁ with an intra-row spacing of 25 cm accumulated the lowest benefit cost ratio of 2.5. (5 cm). It

is clear from the data (Table 2) that the application of 60 K₂O + 120 N₂O + 60 P₂O₅ (K₄), which remained equivalent to treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅), was responsible for the higher gross returns of ₹ 93513/ha. Treatment K₁ (0 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) had lower gross yields, with a total of ₹ 66242/ha. When potassium was applied at a rate of 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha (K₄), the cost of culture was found to be greatest (₹ 26193/ha), whereas treatment K₁ (0 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) recorded the lowest cost of cultivation (₹ 24841/ha).

It is clear from the data (Table 2) that treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅), which came in second, and application of 60 K₂O + 120 N₂O + 60 P₂O₅ (K₄), which recorded much higher net returns of (₹ 67319/ha), respectively. The lower net yields of (₹ 41401/ha) were seen with no potassium treatment. The highest observed 3.6 was in the benefit cost ratio application for 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha (K₄). Use of K₁ (0 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) resulted in the lowest benefit cost ratio of 2.7.

CONCLUSIONS

Sweet corn (Sweet-16) may be planted with an intra-row spacing of 20 cm and an inter-row spacing of 45 cm, and it can be fertilised with 60 kg/ha of K₂O, 120 kg/ha of N₂O, and 60 kg/ha of P₂O₅ in addition to the recommended amounts of N and P₂O₅.

Conference disclaimer:

Some part of this manuscript was previously presented in the conference: 3rd International Conference IAAHAS-2023 "Innovative Approaches in Agriculture, Horticulture & Allied Sciences" on March 29-31, 2023 in SGT University , Gurugram, India. Web Link of the proceeding: <https://wikifarmer.com/event/iaahas-2023-innovative-approaches-in-agriculture-horticulture-allied-sciences/>

ACKNOWLEDGEMENT

Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat- 362001, India for providing me the necessary resources and support for carrying out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bozorgi, H. R., Dousta, H. Z., Sadeghi, S. M., Keshavarz, A., Faraji, A. Study effect of plant density and nitrogen fertilizer on yield and yield components of maize. *World Applied Sciences Journal*. 2011; 13: 147-151.
2. Gomez, K. A. and Gomez, A. A. Statistical procedures for agricultural research. An International Rice Research Institute Book. John Wiley and Sons. New York.. 1984; 680.
3. Gozubenli, H and Konuskan, O. Nitrogen dose and plant density effects on popcorn grain yield. *African Journal of Biotechnology*. 2010; 9(25): 3828-3832.
4. Hamni. H. and Dadari, S. A. Growth and yield analysis of irrigated popcorn (*Zea mays everta* L.) grown in Kadawa as affected by sowing date and intra-row spacing, using correlation coefficient. *ASSET Series*. 2003; 4:63-70.
5. Jat, G., Majumdar, S. P., Jat, N. K. and Majumdar, S. P. Potassium and zinc fertilization of wheat in western arid zone of India. *Indian Journal of Agronomy*. 2013. 58 (1): 67-71.
6. Kumar, Pradeep., Kumar, Rajeev., Singh, S. K. and Kumar, Anil. Effect of fertility on growth yield and yield attributes of pearl millet (*Pennisetum Glaucum* L.) under rainfed condition. *Agriways*. 2014; 2(2): 89-93.
7. Manishaben, Rathod., Bavalgave, V. G., Patil, V. A. and Deshmukh, S. P. Growth, yield and quality of sweet corn (*Zea mays* L. var. *Saccharata*) as influenced by spacing and INM practices under South Gujarat condition. *International Journal of Economic Plants*. 2018; 5(4):170-173.
8. Paradkar, V. K. Effect of N levels and plant spacing on performance of baby corn at Chindwara. Annual Progress Report, Directorate of Maize Research, IARI, New Delhi, 2004. A-77.
9. Patel, P. C. and Kotecha, A. C. Effect of phosphorus and potassium on growth characters, forage yield, nutrient uptake and quality of lucerne (*Medicago sativa* L.). *Indian Journal of Agronomy*. 2006; 51: 242-244.
10. Shekhawat, P. S and Gautam, R. C. Effect of row spacing and weed control methods on the growth attributes and grain yield of maize under tilled and untilled conditions. *Annals of Agricultural Research*. 2002; 23(4): 626-629.

11. Silva, G. Nutrient removal rates by grain crops. Michigan State University Extension. 2017. https://www.canr.msu.edu/news/nutrient_removal_rates_by_grain_crops.
12. Singh, V., Ali, J., Seema, A. K. and Chauhan, T. M. Productivity, nutrient uptake and economics of wheat (*Triticum aestivum* L.) under potassium and zinc nutrition. *The Indian Society of Soil Science*. 2015; 60 (3): 426-430.
13. Yadav, S. S. Tikkoo. A. and Singh, J. P. Effect of potassium on pearl millet-wheat cropping system in coarse textured soils of southern Haryana. *Journal of the Indian Society of Soil Science*. 2012. 60(2): 145-149.

Table 1: Effect of intra-row spacing and potassium levels on dry matter accumulation at relative growth rate, absolute growth rate and crop growth rate of sweet corn

Treatments	Dry matter accumulation at (g plant ⁻¹)				Relative growth rate at (g g ⁻¹ day ⁻¹)			Absolute growth rate at (g/day)			Crop growth rate at (g m ⁻² day ⁻¹)		
	30 DAS	45 DAS	60 DAS	Harvest	30-45 DAS	45-60 DAS	60 DAS-Harvest	30-45 DAS	45-60 DAS	60 DAS-Harvest	30-45 DAS	45-60 DAS	60 DAS-Harvest
Intra-row spacing (cm)													
S ₁ : 5 Intra-row + 45 Inter-rows	24.37	51.02	70.72	106.33	0.0535	0.0497	0.0135	1.81	2.58	1.11	0.00163	0.00231	0.00100
S ₂ : 10 Intra-row + 45 Inter-rows	25.84	52.75	82.90	122.73	0.0561	0.0505	0.0139	1.84	2.64	1.18	0.00165	0.00237	0.00106
S ₃ : 15 Intra-row + 45 Inter-rows	25.85	55.75	92.96	133.62	0.0562	0.0525	0.0140	1.85	2.89	1.38	0.00166	0.00259	0.00124
S ₄ : 20 Intra-row + 45 Inter-rows	27.48	57.94	102.46	144.54	0.0609	0.0571	0.0141	1.88	3.07	1.41	0.00168	0.00275	0.00126
S.Em.±	0.75	2.14	3.42	4.27	0.0020	0.0023	0.0003	0.06	0.11	0.05	0.00005	0.00010	0.00004
C.D. at 5%	NS	NS	9.87	12.32	NS	NS	NS	NS	0.32	0.14	NS	0.00029	0.00013
Potassium levels (kg/ha)													
K ₁ : 0 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	24.95	51.01	73.11	108.14	0.0535	0.0478	0.0136	1.82	2.55	1.14	0.00163	0.00229	0.00102
K ₂ : 20 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	25.31	53.48	85.10	122.34	0.0561	0.0517	0.0139	1.83	2.69	1.20	0.00165	0.00241	0.00107
K ₃ : 40 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	26.55	56.25	93.89	134.74	0.0576	0.0534	0.0140	1.85	2.90	1.36	0.00166	0.00260	0.00122
K ₄ : 60 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	26.72	56.73	96.93	142.00	0.0595	0.0569	0.0141	1.87	3.03	1.39	0.00168	0.00272	0.00124
S.Em.±	0.75	2.14	3.42	4.27	0.0020	0.0023	0.0003	0.06	0.11	0.05	0.00005	0.00010	0.00004
C.D. at 5%	NS	NS	9.87	12.32	NS	NS	NS	NS	0.32	0.14	NS	0.00029	0.00013
Interaction (S × K)													
S.Em.±	1.51	4.27	6.84	8.53	0.0039	0.0045	0.0006	0.12	0.22	0.10	0.00010	0.00020	0.00009
C.D. at 5%	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
C.V. %	10.08	13.62	13.57	11.65	11.99	14.99	7.80	10.85	13.91	13.36	10.76	13.91	13.36

Table 2: Effect of intra-row spacing and potassium levels on economics of sweet corn

Treatments	Gross returns (₹ /ha)	Cost of cultivation (₹ /ha)	Net returns (₹ /ha)	BCR
Intra-row spacing (cm)				
S ₁ : 5 Intra-row + 45 Inter-rows	51.02	70.72	106.33	0.0535
S ₂ : 10 Intra-row + 45 Inter-rows	52.75	82.90	122.73	0.0561
S ₃ : 15 Intra-row + 45 Inter-rows	55.75	92.96	133.62	0.0562
S ₄ : 20 Intra-row + 45 Inter-rows	57.94	102.46	144.54	0.0609
S.Em.±	2.14	3.42	4.27	0.0020
C.D. at 5%	NS	9.87	12.32	NS
Potassium levels (kg /ha)				
K ₁ : 0 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	51.01	73.11	108.14	0.0535
K ₂ : 20 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	53.48	85.10	122.34	0.0561
K ₃ : 40 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	56.25	93.89	134.74	0.0576
K ₄ : 60 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	56.73	96.93	142.00	0.0595
S.Em.±	2.14	3.42	4.27	0.0020
C.D. at 5%	NS	9.87	12.32	NS
Interaction (S × K)				
S.Em.±	4.27	6.84	8.53	0.0039
C.D. at 5%	NS	NS	NS	NS
C.V. %	13.62	13.57	11.65	11.99