

# Antifungal activity of different *Trichoderma* spp. and plant extracts against *Sclerotinia sclerotiorum* (Lib.) de Bary causing stem rot of mustard

## Abstract

The present study was conducted to evaluate the bio-efficacy of different *Trichoderma* spp. and plant extracts against *Sclerotinia sclerotiorum* under laboratory conditions. All the tested *Trichoderma* spp. and plant extracts showed a significant effect on growth inhibition of test fungus over control. Among different *Trichoderma* spp., maximum growth inhibition (73.33%) was observed with *T. viride* followed by *T. harzianum* (67.77%). While as the least efficacy was found in *T. atroviride* (38.37%) against the pathogen. Of all, *Allium sativum* was proved to be most potent at all three tested concentrations (5, 10, and 15%), thereby registering 59.26, 67.41, and 100.0% growth inhibition followed by *Azadirachta indica* and *Lantana camara*, respectively. However, *Argemone maxicana* was least effective at all concentrations in this study.

**Keywords:** *Trichoderma* spp., plant extracts, rapeseed-mustard, *Sclerotinia* stem rot, *in vitro*

## Introduction

India is one of the leading edible oil economy after the USA, China, and Brazil, which contributes about 10 per cent to the global oilseed oil. Rapeseed-mustard is the third major oilseed crop after soybean and palm in the world (Kumar *et al.*, 2014), and about 25 per cent of total oilseed produced by this crop (Yadav *et al.*, 2019). This crop used as an important source of edible oil, industrial oil, livestock feed, vegetable, and soil amendment. The mustard has nutritional value, *viz.*, sugar 1.41g, carbohydrates 4.51g, fat 0.47g, protein 2.56g per 100 g (3.5 oz), and dietary fiber 2g. India holds second place in the area (20.50%) and fourth rank in yield production (10.72%) of rapeseed and mustard, contributing 6.23 m ha and 9.34 mt, respectively, with an average productivity of 1499 kg ha<sup>-1</sup> (Agricultural Statistics at a Glance, 2021). A wide gap occurs between the probable yield and the actual yield at the farmer's field, which is mainly because of biotic and abiotic factors to which this crop is exposed. Amongst, diseases caused by fungi are posing a severe menace. These fungal diseases cause deterioration in oilseed quality and also reduce production. About thirty diseases are known to occur on crops in India (Saharan,

1992). Some of the economically critical diseases of rapeseed-mustard reported in India or abroad are white rust, downy mildew, *Alternaria* blight, stem rot, powdery mildew, *Rhizoctonia* leaf blight, black leg/canker, grey mold, *Fusarium* wilt, damping off, black rot and mosaic (Saha *et al.* 1989).

Amongst, *Sclerotinia* stem rot (SSR) has become one of the most ubiquitous and devastating soil-borne disease distributed worldwide. The disease occurs commonly in cool and wet regions on a number of plants that are predominantly grown in temperate and subtropical regions all over the world (Purdy, 1979; Saharan and Mehta, 2008; Sharma *et al.*, 2014). Earlier, this disease was considered to be of lesser importance, but due to newer crop practices and large cultivation of susceptible cultivars by the farmers, it has now achieved the next place to *Alternaria* blight in their economic importance (Kolte, 1985; Parveen *et al.*, 2007). In India, this disease has been reported in serious proportion in states like Uttar Pradesh, Rajasthan, Madhya Pradesh, Punjab, Haryana, and Bihar (Saharan and Mehta, 2002). In Rajasthan, about 60% yield loss has been observed in heavily infected plants (Ghasolia *et al.*, 2004). According to Shukla (2005), plants infected before initiation of flower can result in 100% crop losses, whereas the plants attacked after flowering suffer 50% yield loss.

The pathogen appears to be non-specific and attacks different plant parts, *viz.*, leaf, stem, and pods. The characteristic symptoms appeared as elongated water-soaked lesions at stem base, which usually enlarge quickly, become necrotic, and subsequently develop patches of white fluffy mycelium (Bolton *et al.*, 2006). Later, the stem is girdled completely and covered by a cottony mycelium growth; it breaks from where it shows rotting and drying. The pathogen survives through latent sclerotia which can remain viable for relatively longer time as they are resistant to adverse conditions (physically and chemically), as well as to degradation by another beneficial microorganism (Wu *et al.* 2008).

The management of *Sclerotinia* rot is absolutely difficult because of the broad host range of the pathogen, variability, and soil-inhabitant nature (Mondal *et al.*, 2015; Smolinska and Kowalska, 2018). The management of stem rot needs a proper understanding of pathogen etiology and epidemiology. Keeping in mind the significance of crop, the current experiment was

conducted *in vitro* to the effectiveness of different *Trichoderma* spp. and plant extracts in managing the disease.

## **Material and Methods**

### ***In vitro* evaluation of *Trichoderma* spp. against *S. sclerotiorum***

Four *Trichoderma* spp. namely, *T. harzianum*, *T. atroviride*, *T. viride*, and *T. virens* were tested to check their effect on the growth of this fungus by using dual culture method (Dennis and Webster, 1971). Twenty ml of sterilized medium (PDA) was transferred into a sterile Petri plate and allowed for solidification. Five mm diameter discs from fresh colony of the pathogen were cut with the help of sterile cork borer and positioned near the periphery of the plate having PDA. Similarly, *Trichoderma* spp. was also placed on another side, i.e., at 180° angle. Plates with no antagonists treated as a control for the pathogen. Each treatment was replicated thrice and incubated at 25±2°C for seven days. The antagonistic activity by *Trichoderma* spp. was recorded after an incubation period of 7 days by assessing the growth of the fungus in both treated and untreated plates. The growth inhibition of the pathogen was measured by using the method given by Vincent (1927).

$$\text{Per cent inhibition (\%)} = (C - T)/C \times 100$$

Where,

C=pathogen growth in control plate

T=pathogen growth in dual culture plate

### ***In vitro* evaluation of plant extracts against *S. sclerotiorum***

Antifungal activity of four plant extracts i.e., Neem (*Azadirachta indica*), Lantana (*Lantana camara*), Satyanashi (*Argemone maxicana*) and Garlic (*Allium sativum*) were also evaluated against *S. sclerotiorum* at three different concentrations (5%, 10% and 15%) by using poison food technique (Dubey and Patel, 2001). About 100 gm of fresh leaves/clove were used and carefully washed in distilled water. Such plant parts were cut into small pieces and then grinded in a mortar and pestle by including 100 ml of distilled water. The basic material was then filtered through muslin cloth (double layer), and then the filtrate was filtered through Whatman

no. 1 filter paper. The prepared plant extracts were autoclaved at temperature of 40°C for 5 minutes to prevent contamination.

The requisite amount of plant extracts were incorporated into sterilized potato dextrose agar medium (non-solidified) and stirred well to make it standardized. Thereafter, 20 ml of amended medium was then poured into 90 mm measuring Petri plates. The observation, thus, recorded on the growth of fungus at all tested concentrations until the growth of test fungus fully covered the un-poisoned Petri plates (check). The percent inhibition in radial growth (T) over control (check) was assessed by using following method given by Vincent, 1947.

$$\text{Per cent inhibition (\%)} = (C - T)/C \times 100$$

Where,

C= growth of pathogen in control plates


T= growth of pathogen in treated plates

## **Result and Discussion**

### ***In-vitro* efficacy of *Trichoderma* spp. against *S. sclerotiorum***

All *Trichoderma* spp. showed a significant impact on radial growth inhibition of test fungus over control. Among all tested *Trichoderma* spp., significant maximum growth inhibition (73.33%) was recorded due to *T. viride* which was followed by *T. harzianum* (67.77%), *T. virens* (44.44%). While as, least efficacy was found in *T. atroviride* (38.37%) against the pathogen (Fig 1).

It is clear from the present observations that among all *Trichoderma* spp., *T. viride* was the most efficient in radial growth inhibition of the fungus followed by *T. harzianum*, *T. virens*, while *T. atroviride* was the least effective in this study (Fig 1). The present findings are in testimony to the results of several earlier reports that also indicated the efficacy of *T. virens*, *T. viride*, and *T. harzianum* against *S. sclerotiorum* in reducing the mycelial growth and sclerotial development (Pandey *et al.*, 2011; Yadav *et al.*, 2011; Sharma *et al.*, 2017). In a study, Mehta *et al.* (2012) also tested the effect of several antagonists against *S. sclerotiorum* (stem rot of mustard) and reported that *T. harzianum* was most superior as compared to other strains of



**Fig 1. Efficacy of different *Trichoderma* spp. on growth inhibition of *S. sclerotiorum***

*Trichoderma in vitro* and it was also found to be excellent in checking length of lesions and post emergence damping-off *in vivo*.

Bio-control agents are known to produce some growth promoting substances (increase seed germination and root and shoot length) and enhance plant nutrition or production of some growth-regulating substances (Cook and Baker, 1983).

#### ***In vitro* efficacy of different plant extracts against *S. sclerotiorum***

All plant extracts significantly reduced the growth of fungus at all concentrations over control. Of all, *Allium sativum* was found to be most effective at all three concentrations of 5, 10, and 15 per cent, thereby registering 59.26, 67.41, and 100% radial growth inhibition, followed by *Azadirachta indica* (51.11, 65.55, and 82.6 %) and *Lantana camara* (43.33, 64.44 and 77.77%), respectively. However, *Argemone maxicana* was least effective at all concentrations in

**Table 1. Effect of different plant extracts on growth inhibition of *S. sclerotiorum***

Plant extracts	Concentrations					
	5%		10%		15%	
	Radial growth (mm)	% Inhibition	Radial growth (mm)	% Inhibition	Radial growth (mm)	% Inhibition
<i>Azadirachta indica</i>	51.00	51.11 <sup>b</sup>	31.00	65.55 <sup>d</sup>	15.66	82.6 <sup>b</sup>
<i>Lantana camara</i>	36.66	43.33 <sup>c</sup>	32.00	64.44 <sup>c</sup>	20.00	77.77 <sup>c</sup>
<i>Argemone maxicana</i>	60.00	33.33 <sup>d</sup>	43.66	51.48 <sup>b</sup>	33.00	63.33 <sup>d</sup>
<i>Allium sativum</i>	44.00	59.26 <sup>a</sup>	29.33	67.41 <sup>a</sup>	00.00	100.0 <sup>a</sup>
<b>Control</b>	90.00	-	90.00	-	90.00	-
<b>L.S.D. (P≤0.05)</b>	4.28	4.76	3.54	3.93	2.83	3.14
<b>S.E(m)±</b>	0.94	1.04	0.78	0.86	0.62	0.69

\*Each value is an average of three replicates. Values within a column followed by different alphabets are significant and some alphabets are non-significant according to Tukey's Test at P≤0.05

comparison to other plant extracts and resulted in 33.33, 51.48, and 63.33 per cent inhibition of growth at 5, 10, and 15% concentrations, respectively (Table 1).

It is, thus, clear from this study *Allium sativum* was the most effective plant extracts, followed by *Azadirachta indica* and *Lantana camara*. However, *Argemone maxicana* was the least effective in this study (Table 1). The results, thus, obtained in the present study are in agreement with the findings of other researchers, who have also noted the efficacy of garlic and neem extracts for inhibition of mycelial growth and sclerotial formation of *S. sclerotiorum* (Kapil and Kapoor, 2005; Tripathi and Tripathi, 2009; Mehta *et al.*, 2011; Yadav *et al.*, 2011; Meena *et al.*, 2013; Fatehpuria *et al.*, 2017). In a study, Fagodia *et al.* (2017) also investigated the efficacy of different botanicals on the growth of *S. sclerotiorum* causing stem rot of coriander and found that *Allium sativum* extract was most potent in inhibiting mycelial growth of fungus, followed by Eucalyptus leaf extract.

## Conclusion

All *Trichoderma* spp. significantly inhibited the growth of *S. sclerotiorum*. However, *T. viride* and *T. harzianum* was noted to be most effective antagonist in inhibiting the maximum radial growth of the pathogen. While as, *T. atroviride* proved to be least efficacious in this study. Of all plant extracts, *Allium sativum* was found superior over all others at all concentrations followed by *Azadirachta indica* and *Lantana camara*. However, *Argemone maxicana* was found least effective in this study. The biopesticides should be used by the farmers alone or in combinations with other cultural practices because it is economical and eco-friendly than chemicals.

## References

**Agricultural Statistics at a Glance (2021)**. Directorate of Economics and Statistics Department of Agriculture and Farmers Welfare Ministry of Agriculture and Farmers Welfare Government of India.

- Bolton, D. M., Thomma, P. H. J. B. and Nelson, D. B. (2006).** *Sclerotinia sclerotiorum* (Lib.) de Bary: biology and molecular traits of a cosmopolitan pathogen. *Mol Plant Pathol*, 7: 1-16.
- Cook, R. J. and Baker, K. F. (1983).** Approaches to biological control. In: The natural and practice of biological control of plant pathogens. St. Paul, Minnesota, USA, *American Phytopathological Society*, pp. 84-131.
- Dennis, C. and Webster, J. (1971).** Antagonistic properties of species groups of *Trichoderma* - III. Production of volatile antibiotics. *Trans. Br. mycol. Soc*, 51: 363-369.
- Dubey S. C, Patel B. (2001).** Evaluation of fungal antagonist against *Thanatephorus cucumeris* causing web blight urd and mung bean. *Indian Phytopath*, 54: 206–209.
- Fagodia, R. K., Godika, S. and Fagodia, B. L. (2017).** Effect of physical parameters on the growth and sclerotia formation of *Sclerotinia sclerotiorum* (Lib.) de Bary, causing stem rot of coriander. *Annals of Plant and Soil Research*, 19 (1): 54 – 58.
- Fatehpuria, P. K., Sasode, R. S., Pandya, R. K., Singh, R., & Gupta, J. C. (2017).** Efficacy of different inoculation techniques for testing the pathogenicity of *Sclerotinia sclerotiorum* causing Sclerotinia blight of *Brassica juncea*. *IJCS*, 5 (5), 1937-1940.
- Ghasolia, R. P., Shivpuri, A. and Bhargava, A. K. (2004).** Sclerotinia rot of Indian mustard (*Brassica juncea*) in Rajasthan. *Indian Phytopathol*, 57:76-79.
- Kapil, R. and Kapoor, A. S. (2005).** Management of white rot of pea incited by *Sclerotinia sclerotiorum* using *Trichoderma* spp. and biopesticides. *Indian Phytopathology*, 58:10-16.
- Kolte, S. J. (1985).** Diseases of Annual Edible Oilseed Crops, Rapeseed-Mustard and Sesame Diseases. *CRC Press Inc Boca Raton*, 2, 135p.
- Kumar, A. (2014).** Challenge of edible oils: Can Brassicas deliver. *J Oilseed Brassica*, 5:83-86.
- Meena, P. D., Gour, R. B., Gupta, J. C., Singh, H. K., Awasthi, R. P., Netam, R. S., Godika, S., Sandhu, P. S., Prasad, R., Rathi, A. S., Rai, D., Thomas, L., Patel, G. A.**

**and Chattopadhyay, C. (2013).** Non-chemical agents provide tenable, eco-friendly alternatives for the management of the major diseases devastating Indian mustard (*Brassica juncea*) in India. *Crop protection*, 53: 169-174.

**Mehta, N., Hieu, N. and Sangwan, M. (2012).** Efficacy of various antagonistic isolates and species of against causing white stem rot of mustard. *Journal of Mycology and Plant Pathology*, 42(2): 244-250.

**Mehta, N., Hieu, N. T. and Sangwan, M. S. (2011).** Efficacy of some botanicals against *Sclerotinia sclerotiorum* inciting white stem rot of rapeseed-mustard. *Plant Dis Res*, 26: 82-86.

**Mondal, B., Khatua, D. C., Hansda, S., Sharma, R. (2015).** Addition to the host range of *Sclerotinia sclerotiorum* in West Bengal. *Scholars Academic Journal of Biosciences*, 3: 361-364.

**Pandey, P., Kumar, R. and Mishra, P. (2011).** Integrated approach for the management of *Sclerotinia sclerotiorum* (Lib.) de Bary, causing stem rot of chickpea. *Indian Phytopath*, 64 (1): 37-40.

**Parveen, K., Haseeb, A. and Shukla, P. K. (2007).** Management of *Sclerotinia sclerotiorum* on *Mentha arvensis* C.V.Gomti. *J. Mycol. Pl. Pathol*, 37 (1):33-36.

**Purdy, L. H. (1979).** *Sclerotinia sclerotiorum*: history, diseases and symptomatology, host range, geographic distribution, and impact. *Phytopathology*, 69: 875-880.

**Saha, L. R., Singh, H. B., Raychaudhuri, S. P. and Verma, J. P. (1989).** Diseases of rapeseed and mustard and their management. *Rev. Tropical Pl. Pathol*, 5: 47-77.

**Saharan, G. S. (1992).** Management of rapeseed and mustard diseases. In: Advances in oilseed research. Vol.1 Rapeseed and Mustard (Kumar, D. and Rai, M, eds.) Scientific Pub. Jodhpur, pp,155-188.

**Saharan, G. S. and Mehta, N. (2002).** Fungal Diseases of Rapeseed-Mustard. In: Diseases of Field Crops (Gupta, V.K. and Paul. Y.S. eds.). Indus Publishing Company, New Delhi, pp.193-228.

- Saharan, G. S. and Mehta, N. (2008).** *Sclerotinia* Diseases of Crop Plants: Biology, Ecology and Disease Management: Springer Press. *Sclerotinia sclerotiorum* Isolates in Northern Parts of Iran. *World Appl. Sci. J*, 8 (3):326-333.
- Sharma, P., Meena, P. D. and Singh, D. (2014).** Effect of *Sclerotinia sclerotiorum* culture filtrate on seed germination and seedling vigour of Indian mustard (*Brassica juncea* cv. Rohini). *J Oilseed Brassica*, 5: 158-161.
- Sharma, P., Meena, P. D., Singh, S. and Rai, P. K. (2017).** Efficacy of Micro-Nutrients, Fungicides and Bio-Agents against Sclerotinia Stem Rot (*Sclerotinia sclerotiorum*) of Indian Mustard. *Int. J. Curr. Microbiol. App. Sci*, 6(10), 620-626.
- Shukla, A. K. (2005).** Estimation of yield losses to Indian mustard (*Brassica juncea*) due to Sclerotinia stem rot. *J Phytol Res*, 18: 267-268.
- Singh, R., Palat, R., Singh, N. and Narayan, P. (2002).** Effects of different solid media on the growth and sclerotial production of *Sclerotinia sclerotiorum*. *Prog Agric*, 2: 180.
- Smolinska, U., B. Kowalska, (2018).** Biological control of the soil-borne fungal pathogen *Sclerotinia sclerotiorum* a review. *Journal of Plant Pathology*, 100:1- 12.
- Tripathi, A. K. and Tripathi, S. C. (2009).** Management of Sclerotinia stem rot of Indian mustard through plant extracts. *Vegetos*, 22 (1):1-3.
- Vincent, J. M. (1947).** Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*, 159: 239-241.
- Wu, B. M. and Subbarao, K. V. (2008).** Effects of soil temperature, moisture and burial depths on carpogenic germination of *Sclerotinia sclerotiorum* and *S. minor*. *Phytopathology*, 98: 1144–1152.
- Yadav, M. S., Godika, S., Yadava, D. K., Ahmad, N., Mehta, N., Bhatnagar, K., Agrawal, V. K., Kumar, A., Thomas, L. and Chattopadhyay, C. (2019).** Prioritizing components of package of integrated pest management in Indian mustard (*Brassica juncea*) in India for better economic benefit. *Crop Protection*, 120: 21-29.

**Yadav, M., Rakholiya, K. B. and Pawar, D. M. (2011).** Evaluation of bioagents for management of the onion purple blotch and bulb yield loss assessment under field conditions. *The Bioscan*, 8(4): 1295-1298.

UNDER PEER REVIEW