

Nuclear Technology in Horticulture: Boosting Productivity, Controlling Pests, Conserving Water

ABSTRACT

Horticulture faces significant challenges due to the destruction of crops by pests and diseases, both before and after harvesting, resulting in a significant reduction in crop production. Nuclear technology has emerged as a valuable tool for enhancing agricultural productivity. Despite the common association of nuclear technology with energy and weaponry, its use in Horticulture is diverse and multifaceted. Physical mutagens such as nuclear radiation, including gamma rays, X-rays, and UV light, as well as particle radiation, can be employed to induce chromosomal breakages, cross-linking of DNA strands, nucleotide deletion, and substitution. UV rays can be used to irradiate cell suspensions and pollen grains in the early or late uninucleate phases. Through the application of nuclear technology, agricultural productivity can be improved by increasing crop yields, controlling pests and diseases, and enhancing water quality. Food preservation can also be achieved through food irradiation, where ionizing radiation is applied to target pathogens to break their DNA bonds. Nuclear technology can also be utilized to induce beneficial mutations in crops through genetic modification, sterilization-based pest control, and water usage management. In conclusion, nuclear technology presents an innovative and effective solution for addressing challenges in Horticulture, thus contributing significantly to global food security.

Keywords : Mutagenesis, Sterile insect technique, Crop improvement, Food irradiation, Water use efficiency

INTRODUCTION

Although nuclear technology is mostly associated with nuclear energy and weapons, it has numerous applications in Horticulture. These technologies have significantly contributed to the development of horticultural practices by promoting crop production, disease and pest control, and improved water quality. Physical mutagens, including electromagnetic radiation such as gamma rays (emitted from radioactive cobalt-60), X-rays, UV light, and particle radiation such as fast and thermal neutrons, alpha and beta particles, have been utilized in these applications. Ionizing radiation, for example, induces chromosomal breakages that facilitate cross-linking of DNA strands, nucleotide deletion, and substitution (Oladosuet *al.*, 2016). Non-ionizing physical mutagens such as UV rays have also been used to irradiate cell suspensions and pollen grains in the late or early uninucleate phases due to their low penetration. Additionally, various radioisotopes such as P^{32} ($t_{1/2}$ 14.3 days, β 1.7 meV, γ 0), C^{14} ($t_{1/2}$ 5570 yrs., β 0.155 meV, γ 0), Ca^{45} ($t_{1/2}$ 165 days, β 0.254 meV, γ 0), H^3 ($t_{1/2}$ 12.46 yrs, β 0.018 meV, γ 0), Na^{22} ($t_{1/2}$ 2.58 days, β 0.54 meV, γ 1.28 meV), Na^{24} ($t_{1/2}$ 15.05 hrs, β 1.39 meV, γ 1.368 to 2.754 meV), Co^{60} ($t_{1/2}$ 5.24 yrs, β 0.31 meV, γ 1.17 to 1.33 meV), Cu^{64} ($t_{1/2}$ 12.9 hrs, β

0.57 meV, γ 1.34 meV), Fe^{55} ($t_{1/2}$ 2.94 yrs, β 0, γ 0), Fe^{59} ($t_{1/2}$ 44.3 days, β 0.271 meV, γ 1.30 meV), Mn^{54} ($t_{1/2}$ 314 days, β 0, γ 0.84 meV), K^{42} ($t_{1/2}$ 12.47 hrs, β 3.58 meV, γ 1.51 meV), Rb^{86} ($t_{1/2}$ 18.68 days, β 1.82 meV, γ 1.08 meV), S^{35} ($t_{1/2}$ 89 days, β 0.168 meV, γ 0), Zn^{65} ($t_{1/2}$ 246 days, β 0.325 meV, γ 1.12 meV) are readily available in sufficient quantities and have been widely employed to address several problems of interest in Horticulture and other domains. Horticulturists have utilized several radioisotopes for conducting seed germination studies. For instance, P^{32} has been applied to examine phosphorus uptake in seeds, while C^{14} has been utilized to study carbon fixation and metabolism. Ca^{45} has been employed to analyze calcium absorption and transport in plant roots, and H^3 has been used to study water uptake and movement in plants. To investigate ion uptake and transport in plant roots, Na^{22} and Na^{24} have been utilized.

Moreover, a study (Srivastava, *et al.*, 2007) investigated the effects of gamma radiation emitted by ^{60}Co on *Gladiolus* plants. The exposed plants exhibited significant changes compared to the control group. Gamma radiation influenced plant height, leaf size, and overall plant structure. Flowering patterns were also modified, resulting in alterations in flower size, color, and shape, as well as the occurrence of deformities and changes in floral symmetry. Reproductive capacity was negatively impacted, leading to reduced production of viable seeds and decreased pollen fertility. Additionally, gamma radiation caused variations in chlorophyll content, photosynthetic efficiency, and antioxidant enzyme activity. These findings underscore the importance of radiation safety measures in horticulture and highlight the necessity for further research to comprehend the underlying mechanisms and develop strategies to mitigate radiation-induced damage in ornamental plants. Mn^{54} has been applied to study manganese uptake and transport, and K^{42} has been utilized to investigate potassium uptake and transport. Apart from these, several other radioisotopes such as Rb^{86} , S^{35} , and Zn^{65} have been used for various studies related to plant growth and nutrient uptake. (Sahu and Sahu, 2019)

BREEDING NEW SEED VARIETIES

Radiation generation resulted to seed types which have better yields. One well-known case of a successful crop is the "miracle" rice which has elevated the price of rice manufacturing substantially. Crop development in trendy includes genetic variation, skewed in the direction of greater beneficial developments. Different varieties of radiation can be used to initiate those mutations for crop improvement. Traits that the researcher's intend to deliver commonly encompass better yield per crop, resistance to diseases, better nice of the crop, and quicker ripening. The advance trials of genetic mutation began in the 1930s, and now this technique of the usage of radiation to stimulate mutations in plant life has been extensively used. Radiation-prompted mutation technologies have grown to be a massive part of plant breeding methods. Here are some important cultivars and varieties which had been released from various research stations with specific quality characters (Rai *et al.*, 2018).

Fruit crops	Cultivar	Year	Mutagens	Improved fruit traits
Mango	Rosica	1966	Spontaneous mutation	Large and good quality
Apple	Golden Haldegg	1966	y-rays	Fruit size
	McIntosh 8F-2-32	1970	y-rays	Skin colour
Grapefruit	Senbatsu-Fuji-2-Kei	1965	y-rays	Fruit colour
	Rio Red	1964	tHn	Fruit colour
Indian jujube	Star Ruby	1970	tHn	Seedless
	Mahong	1966	tHn	Round, pink rose sweeter taste
Loquat	Shiro-moji	1981	y-rays	Fruit size
Orange	Xuegan 9-12-1	1963	y-rays	Seedless
	Hongju 420	1986	y-rays	Seedless
Peach	Eureka 22	1987	X-rays	Fruit set, fruit quality
	Valencia 2	1967	X-rays	Fruit quality
Sweet cherry	Magnif	1968	y-rays	Large, red skin
	Plovdiv	1981	y-rays	Large, fruit quality
Pear	Lapins	1963	X-rays	Larger size, firmer
	Compact Lambert	1964	X-rays	Compact growth
Almond	Ferrovia spur	1992	X-rays	Dwarfness
	Fuxiangyanghongdili	1983	y-rays	Eating, cooking quality
Fig	Supernova	1987	y-rays	Late maturity
	Boi	1979	y-rays	Not specified
Banana	Novaria	1993	y-rays	Earliness
	Al-beely	2007	y-rays	High yield
Japanese pear	Gold Nijisseiki	1993	y-rays	Disease resistance
Papaya	PusaNanha	1986	y-rays	Dwarfness
Plum	Spurdente-Ferco	1968	y-rays	Earliness
Pomegranate	Karabakh	1979	y-rays	--
Sea buckthorn	zyriank			

Source: [31]

Table 10. Other important varieties developed through mutation breeding

Fruit crop	Variety	Parents	Nature of mutation
Mango	Rosica	Rosa-do-delca (Peru)	Natural
	Davis Haden	Haden	Natural
Grape	Marvel Seedless	Delight	Induced
Banana	High Gate	Gros Michel	Natural
	Motta Poovan	Poovan	Natural
Orange	Krishna Vazahi	Virupakshi	Natural
	Washington Novel	Navel Orange	Natural
Grape fruit	Foster	Walter	Natural
	Red Blush	Thompson	Natural
	Thompson	Marsh Seedless	Natural
	Star Ruby	Hudson	Induced

Source: Ray, P.K. 2002 [3]

Table 1 : List of Fruit cultivars generated with the induced/natural mutation

CONTROLLING PESTS AND DISEASES

Nuclear technology has been playing a crucial role in enhancing crop production by controlling pests and diseases, which typically destroy more than a third of crops before and after harvesting. Radioisotope techniques are widely used in the place of toxic chemicals in pest control. The sterile insect technique (SIT) is the most commonly used method, which involves sterilizing male insects in a laboratory using gamma radiation and releasing them into the wild. When these sterile insects mate, no offspring are produced, which helps to reduce the insect population and eventually suppress its growth. SIT has been successfully used to eliminate the Medfly, which poses a threat to around 250 species of horticultural products in some parts of the US, Chile, and Mexico. Furthermore, radioisotopes have been used in developing disease-resistant plants and animals, making them more immune to pest attacks and diseases, similar to breeding new seed varieties (FAO, IAEA, 2018).

SIT(STERILE INSECT TECHNIQUE)

There are concerns that the continued use of pesticides has a negative impact on the environment and leads to the development of pesticide resistance in many insect species (Australian National Botanic Gardens, 2005). Furthermore, pesticides kill not only the target species but also many other beneficial pest species that help to maintain the natural ecological balance in crop fields. A very powerful programme integrating the SIT had been installed against tropical fruit flies, a few species of tsetse flies - *Glossina* spp., the red bollworm *Pectinophoragossypiella* (Saunders), and the codling moth *Cydia pomonella* (L.). Development of the SIT to be used against the boll weevil *Anthonomus grandis*, *grandis* Boheman and the gypsy moth *Lymantria dispar* (L.) has ended. However, it's miles in development forswheat potato weevil species, *Cylasformicarius* (F.) and *Euscepespostfasciatus* (Fairmaire), the fake codling moth *Cryptophlebialeucotreta* (Meyrick), the carob moth *Ectomyeloisceratoniae* (Zeller), the cactus moth *Cactoblastis cactorum* (Berg), the Old World screwworm *Chrysomya bezziana* (Villeneuve), additional

Glossina spp., different *Anastrepha spp.*, the Mexican fruit fly (*Anastrephaludens*, Loew) was eradicated from most of northern Mexico. Fruit fly species, inclusive of Mediterranean fruit fly (*Ceratitidis capitata*), Caribbean fruit fly (*Anastrephasuspensa*) and Mexican fruit fly (*Anastrephaludens*) in the Americas, Queensland fruit fly (*Bactroceratryoni*) in Australia, the oriental fruit fly (*Bactrocera dorsalis*) (Australian National Botanic Gardens, 2005).

A comprehensive review of the action of ionizing radiation on insects has been presented by Hilchey (1958). Lethal effects of X-radiation on immature stages of the oriental fruit fly and the melon fly were reported by Koidsumi (1930). Generally, dosages of 10,000 to 25,000 I are required to prevent reproduction in insects. Indications are that these dosages may be low enough to permit the rapid or large-volume treatment that may be required for practical quarantine purposes without injurious side effects on most host material (Pomerantz & Morgan, 1955; Richardson & Balock 1959; Carpenter, & Bloem, 2000).

IMPROVING WATER USE EFFICIENCY

Nuclear, and particularly isotopic technology can be used to improve water use performance in horticultural contexts. Since horticulture is the primary user of freshwater resources, assisting water control can deliver an extremely useful effect on the sustainable use of freshwater resources. In particular, isotopic techniques can: 1). optimize irrigation scheduling via way of correctly tracking soil in order to decrease water losses, 2). optimize crop's water absorption cost from rainfall or irrigation, and 3). help in deciding on crops with better tolerance to drought and better crop water productivity. Optimizing irrigation scheduling is stated to affect the performance of water utilization maximum directly. So far, major advances had been made in water tracking with the use of isotopic technology. Stable isotopes may be used to assesses the abundance of oxygen, carbon, nitrogen and hydrogen in soil, water and plant helping to figure out nutrient fluxes, in the soil.

FOOD PRESERVATION

Food irradiation entails subjecting the meals to cautiously managed quantities of ionizing radiation, including beta debris or gamma rays, to interrupt the DNA bonds of certain pathogens. This is particularly powerful in destroying the reproductive cycle of microorganisms and pathogens. Such radiation can eliminate undesirable organisms and specific, non-spore-forming pathogenic microorganisms including *salmonella*. It also can intrude with a physiological process including sprouting in potatoes or onions. Thus, the shelf life of many ingredients may be prolonged appreciably, and meals-born ailment organisms including E-coli (0157:H7) may be dramatically reduced. The beta rays are produced via way of means of accelerators while the gamma rays are typically produced through the radioactive decay of cobalt-60. X-rays also can be efficiently used. They are typically produced by means of accelerators wherein the beta rays are directed onto a goal cloth including tungsten that converts the power into x-rays. (Mead, 1991).

CONCLUSION

In conclusion, nuclear technology has proven to be a valuable tool for improving agricultural productivity by enhancing crop yields, controlling pests and diseases, and managing water usage. The use of physical mutagens, including ionizing and non-ionizing radiation, has facilitated the induction of beneficial mutations in crops, leading to the development of new seed varieties with improved traits. Additionally, food irradiation has been shown to be an

effective method for preserving food by breaking down DNA bonds of harmful pathogens. The utilization of radioisotopes in plant growth and nutrient uptake studies has also contributed to the development of horticultural practices. Overall, nuclear technology has played an important role in addressing challenges in Horticulture and has significant potential to contribute to global food security.

REFERENCES

A.E. Walter, (2003) [The Medical, Horticultural, and Industrial Applications of Nuclear Technology](#), *Pacific Northwest National Laboratory*, 16 Nov.

Application of Radioisotopes in Agriculture and Biology," in *Radioisotopes: Applications in Physical Sciences*, edited by Nirmal Singh and Ajaib Singh (*Springer*, 2019), pp. 405-425.

Australian National Botanic Gardens, 2005

B. Singh, J. Singh, and A. Kaur, (2013) "[Applications of Radioisotopes in Agriculture](#), *Int. J. Biotechnol. Bioeng. Res.* **4**, 3.

[Beneficial Uses of Radiation](#) (2008). Nuclear Energy Institute, November.

Carpenter, J. E., & Bloem, S. (2000). The need for and development of area-wide pest management programs using the sterile insect technique. In *Sterile Insect Technique: Principles and Practice in Area-Wide Integrated Pest Management* (pp. 1-36). *Springer*, Dordrecht.

[Deficit Irrigation Practices](#), (2002) Food and Agriculture Organization of the United Nations, 2002.

International Atomic Energy Agency (IAEA) or the United Nations Food and Agriculture Organization (FAO), (2018).

Koidsumi, Kiyooki. (1930). Quantitative studies on the lethal action of X-rays upon certain insects. *Jour. Soc. Trop. Agric.* **2**: 243-62.

Mead, G. C. (1991). The safety and nutritional adequacy of irradiated food. *Food Research International*, **24**(5), 511-517. [Nuclear Technology for a Sustainable Future](#) International Atomic Energy Agency, June 2012.

Oladosu Y, Rafii M, Abdullah N, Hussin G, Ramli A, Rahim H. (2016). Principle and application of plant mutagenesis in crop improvement: A review. *Biotechnology and Biotechnological Equipment*.;30(1):1-16.

Pomerantz, R., and Bruce H. Morgan. (1955). Food radiation roundup. *Food Eng.* **27**(8): 43-47,154.

Rai, M. K., & Rai, M. P. (2018). Applications of ionizing radiations in mutation breeding of vegetatively propagated crops. In *Induced Plant Mutations in the Genomics Era* (pp. 191-210). *Springer*, Cham.

Srivastava, P., Singh, R.P., & Tripathi, V.K. (2007). Response of gamma radiation ^{60}Co on vegetative and floral characters of *Gladiolus*. *Journal of Ornamental Horticulture*, **10**(2), 13-36.