

Original Research Article

Effect Ayurveda-based Comprehensive Diabetic Care Program on Glycemic Control in Type 2 Diabetic Mellitus Patients: An Observational Study

ABSTRACT

Background: The prevalence of diabetes mellitus in India is alarmingly high despite availability of several therapeutic agents. Thus, the current study was conducted to understand the role of Ayurveda based Comprehensive Diabetes Control (CDC) program in the management of type 2 diabetes mellitus.

Methods: A retrospective, observational study was conducted from January 2014 to September 2022 at Madhavbaug Clinic, Maharashtra. Patients aged 18 years and above with a diagnosis of type 2 diabetes mellitus according to the American Diabetes Association (ADA) with body mass index >24 participated in the CDC program and were included in this study. Follow-up was conducted after 90 days. Baseline data was compared with follow-up data.

Results: A total of 63 patients were assessed. Of the male patients, 29.3%, 43.9%, and 26.8% were found to have impaired, negative and positive glucose tolerance. Among the females, 22.7%, 63.6%, and 13.6% were found to be have impaired, negative and positive glucose tolerance. Weight decreased from 70.70 ± 9.82 kg to 65.15 ± 9.83 kg ($p < 0.001$) from day 1 to day 90. Body mass index decreased from 27.00 ± 3.88 to 24.91 ± 3.75 ($p < 0.001$) from day 1 to day 90. Glycated haemoglobin (HbA1c) decreased from $7.87 \pm 1.90\%$ to $5.79 \pm 0.51\%$ ($p < 0.001$) from day 1 to day 90.

Conclusion: Remission of type 2 diabetes mellitus in obese patients is possible with simple Ayurvedic procedures and diet compliance.

Keywords: Ayurveda, body mass index, diabetes mellitus, diet, exercise, glycated haemoglobin

UNDER PEER REVIEW

INTRODUCTION

Diabetes mellitus is a complex and multifactorial metabolic disorder. It is triggered by reduction in insulin or resistance to the effects of insulin leading to hyperglycaemia [1]. Symptoms include frequent urination, increased fluid uptake, blurred vision, weight loss, lethargy, and fluctuating energy levels. The risk factors of diabetes include overweight/obesity, heredity, sedentary lifestyle, poor diet, smoking, and alcohol consumption. The American Diabetes Association recommends management through exercise, diet, and lifestyle [2]. However, current management strategies focus on reducing hyperglycaemia through targeted molecular drugs which fail to address the chronicity and systemic complications of diabetes – this evokes lifetime dependency on medication. The need of the hour is a paradigm shift in the perspective of diabetes management [1].

Ayurveda is an ancient evidence-based medicinal practice that uses several herbal formulations and therapeutic measures such as *Panchakarma* including meditation and Yogic exercises to avert, alleviate and manage diabetes according to individual patients [3]. There is existing evidence indicative that Ayurveda may augment therapy and management of diabetes mellitus. It can therefore play a role as a robust tool for diabetes mellitus management. The Comprehensive Diabetic Care (CDC) Program combines *Panchakarma* and diet management to manage diabetes mellitus. Thus, the rationale of the current study was to understand the role of Ayurveda based CDC program in the management of type 2 diabetes mellitus.

MATERIALS & METHODS

Study design and patient population

A retrospective, observational study was conducted at Madhavbaug Clinic between January 2014 and September 2022. Patients aged 18 years and above with a diagnosis of type 2 diabetes mellitus according to the American Diabetes Association (ADA) [4] and with body mass index >24 who participated in the CDC program were included in this study. Patients that were underweight were excluded from the study. Patients with body mass index 25.1–28.0, 28.1–30.0, 30.1–32.0 were categorized as overweight, obese 1, obese 2, and morbid obese, respectively.

Comprehensive Diabetes Control Program

The Comprehensive Diabetes Control (CDC) program comprises of 3 steps, each session lasting for 65–75 mins. The procedures commenced after patients had a light breakfast. The first procedure was *Snehana* which is external oleation with Neem oil massaged on the hands, legs, shoulders, thorax, abdomen, and the back in a centripetal manner. The duration of this massage was 15–25 mins as 15–30 strokes were applied on each of the aforementioned body parts. *Swedana* was the next procedure and refers to heat therapy with Dashmoola decoction. The patient's body was positioned in a supine position in a wooden box. The patient's neck was permitted to protrude outside the box. The duration of this therapy was 10–15 mins followed by a relaxation period of 3–4 minutes after the therapy. The third step was *Basti* which was medicated enema of 100 ml *Gymnema sylvestre*, *Berberis aristate*, and *Glycyrrhiza glabra*. This was administered through rectum administration with the drug to remain inside the body for >15 mins to allow maximum absorption. The CDC program comprises 6 sessions in total i.e. 4 sessions within the first month, 1 in the second month and the 1 in the third month, an entire duration of 90 days. Patients were restricted to a diet plan comprising 800–1000 calories. The diet comprised low carbohydrates, moderate proteins, and low fats.

Glucose tolerance test

The glucose tolerance test, also known as oral glucose tolerance test, measures the body's response to sugar (glucose). The glucose tolerance test can be used to screen for type 2 diabetes. This test indicates glucose metabolism strength of an individual, 75 gm of oral sugar infusion was given to the patients after collection of fasting samples, and again after 1 hr and 2 hrs of post prandial blood sugar was recorded. Patients were asked to sit in one position for the duration from fasting till 2 hrs after sampling. No Orally administered antihyperglycemic agents (OHAs) or insulin was taken before or during the procedure. Normal glucose tolerance was considered as <90 mg/dL after fasting, <180 mg/dL at 1 hr, and <140 mg/dL at 2 hrs. Impaired glucose tolerance was considered as <90 mg/dL after fasting, >180 mg/dL at 1 hr, and <140 mg/dL at 2 hrs. Diabetes was considered as >90 mg/dL at fasting, >180 mg/dL at 1 hr, and >140 mg/dL at 2 hrs.

Data collection

Data for patient demographics, anthropometrics, and laboratory findings were collected and analysed from the patients' medical records. On day 1 of the CDC program, a detailed patient history was taken, anthropometric measurements were recorded and the fasting serum glycated haemoglobin (HbA1c) levels were measured. This was repeated on day 90 of the CDC program. Data of day 1 was compared with data of day 90. Data of only those patients who had completed a total of 6 sessions was collected and analysed.

Statistical analysis

Categorical data are shown as number (percentage) and continuous data are shown as mean \pm standard deviation. Paired t test was used to calculate the difference between baseline and the 90-day follow-up. P value ≤ 0.05 was considered as statistically significant.

RESULTS

Baseline demographic and anthropometric measurements

The mean age of the study population was 50.51 ± 10.62 years. Of the 61 patients, 41 (65.1%) patients were male. Weight decreased from 70.70 ± 9.82 kg to 65.15 ± 9.83 kg ($p < 0.001$) from day 1 to day 90. Body mass index decreased from 27.00 ± 3.88 to 24.91 ± 3.75 ($p < 0.001$) from day 1 to day 90. HbA1c decreased from $7.87 \pm 1.90\%$ to $5.79 \pm 0.51\%$ ($p < 0.001$) from day 1 to day 90. The anthropometric measurements at baseline and 90-day follow-up are detailed in **Table 1**. The demographic and anthropometric measurements according to age, gender, HbA1c, and body mass index at baseline and 90-day follow-up are elaborated in **Table 2**.

Glucose tolerance testing

Of the male patients, 29.3%, 43.9%, and 26.8% were found to have impaired, negative and positive glucose tolerance. Among the females, 22.7%, 63.6%, and 13.6% were found to be have impaired, negative and positive glucose tolerance. The glucose tolerance testing findings are displayed in **Figure 1**.

DISCUSSION

Overweight and obesity are significant risk factors for the development of non-communicable diseases, particularly type 2 diabetes – majority of the individuals with type 2 diabetes mellitus are overweight or obese. According to the World Health Organization, obesity and overweight account for 44% diabetes cases and predict the prevalence of obesity-related diabetes to double by 2025 [5]. The main causative factors of overweight and obesity are high-calorie or high-fat foods, inadequate physical activity, and a sedentary lifestyle. European guidelines recommend at least 150 min/week of moderate aerobic activity along with 3 weekly sessions of resistance training to increase muscle strength [6-9]. Physical activity reduces intra-abdominal fat, increase in lean mass, and causes improvement in glucose tolerance, insulin sensitivity, and physical fitness. Moreover, current evidence suggests even more favourable results of weight loss when combined with an energy-restricted diet. This refers to a low-fat and low-carbohydrate diet [5]. The Finnish Diabetes Prevention Study [10] revealed that intensive dietary and exercise regimen decreased overall risk of diabetes by 58%. The Diabetes Prevention Program [11] revealed moderate weight loss and lifestyle intervention in an obese population with impaired glucose tolerance can reduce incidence of diabetes by 58% while metformin alone reduced it by only 31%. The Look AHEAD [12] study evidenced that 5–10% body weight reduction can improve overall fitness, reduce HbA1c levels, improve cardiovascular disease risk factors, and decrease the use of anti-hyperglycaemic levels, anti-hypertensive, and lipid-lowering medications after 1 year. In the current study, weight decreased from 70.70 ± 9.82 kg to 65.15 ± 9.83 kg after 90 days. Body mass index has a strong correlation with diabetes and insulin resistance and is still used to classify obesity and overweight. Body mass index decreased from 27.00 ± 1.90 to 25.79 ± 0.51 after 90 days in our study.

Ayurveda is emerging as a complementary and alternative treatment strategy for chronic illness. The current study aimed to understand the role of Ayurveda based CDC program in the management of type 2 diabetes mellitus. The main goal of the management of diabetes mellitus is to reduce blood glucose levels within a therapeutic range through medication, diet, and lifestyle modification. HbA1c is a widely used diagnostic marker, reflective of average HbA1c levels over a 3-month period [13]. A reduction in HbA1c of at least 0.3% or 0.4% is considered to be clinically meaningful, [14]. The current study revealed changes of 0.44%, -0.97%, -2.60%, and -5.69% for HbA1c ranges >5.7, 5.8–7.0, 7.1–10.0, and 10.1–15.0, respectively. These findings are in line with other similar studies [15-17].

Yoga has found acceptance and has been practiced by individuals with all social strata. Yoga regulates eating pattern and leads to adherence of medication. A 3-arm controlled trial aimed to validate if Yoga can improve blood glucose and HbA1c in individual with pre-diabetes. Study findings revealed significant decrease in fasting blood sugar (FBS) (-21.13 ± 21.16 mg/dl) and postprandial blood sugar (PPBS) (-15.0 ± 14.89 mg/dl) in group 1 (*Rasahara* and Yoga) and FBS (20.62 ± 32.68 mg/dl) in group 2 (Yoga), while the increases in group 3 (no intervention) was only significant for the PPBS level (9.62 ± 21.83 mg/dl) [18].

CONCLUSION

Remission of type 2 diabetes mellitus with obesity is possible with simple Ayurvedic procedures and diet compliance.

NOTE:

The study highlights the efficacy of "Ayurved" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

REFERENCES

1. Thottapillil A, Kouser S, Kukkupuni SK, Vishnuprasad CN. An 'Ayurveda-Biology' platform for integrative diabetes management. *J Ethnopharmacol.* 2021 Mar 25;268:113575.
2. Gordon A, Buch Z, Baute V, Coeytaux R. Use of Ayurveda in the Treatment of Type 2 Diabetes Mellitus. *Glob Adv Health Med.* 2019 Aug 7;8:2164956119861094.
3. Banerjee S, Debnath P, Rao PN, Tripathy TB, Adhikari A, Debnath PK. Ayurveda in changing scenario of diabetes management for developing safe and effective treatment choices for the future. *J Complement Integr Med.* 2015 Jun;12(2):101-10.
4. American Diabetes Association. (2013) Diagnosis and classification of diabetes mellitus. *Diabetes Care*, 36 Suppl 1 (Suppl 1), S67-74.
5. Leitner DR, Frühbeck G, Yumuk V, Schindler K, Micic D, Woodward E, Toplak H. Obesity and Type 2 Diabetes: Two Diseases with a Need for Combined Treatment Strategies - EASO Can Lead the Way. *Obes Facts.* 2017;10(5):483-492.
6. Yumuk V, Tsigos C, Fried M, Schindler K, Busetto L, Micic D, Toplak H, Obesity Management Task Force of the European Association for the Study of Obesity: European guidelines for obesity management in adults. *Obes Facts* 2015; 8: 402–424
7. Poirier P, Despres JP: Exercise in weight management of obesity. *Cardiol Clin* 2001; 19: 459–470.

8. Willis LH, Slentz CA, Bateman LA, Shields AT, Piner LW, Bales CW, Houmard JA, Kraus WE: Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *J Appl Physiol* 2012; 113:1831–1837
9. Geliebter A, Ochner CN, Dambkowski CL, Hashim SA: Obesity-related hormones and metabolic risk factors: a randomized trial of diet plus either strength or aerobic training versus diet alone in overweight participants. *J Diabetes Obes* 2014; 1: 1–7.
10. Lindström J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, Uusitupa M, Tuomilehto J, Finnish Diabetes Prevention Study Group: The Finnish Diabetes Prevention Study (DPS): lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care* 2003; 26: 3230–3236.
11. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM, Diabetes Prevention Program Research G: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002; 346: 393–403.
12. Look ARG, Wing RR: Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med* 2010; 170: 1566–1575.
13. Saudek CD, Brick JC. The clinical use of hemoglobin A1c. *Journal of diabetes science and technology*. 2009 Jul;3(4):629-34.
14. Chattopadhyay K, Wang H, Kaur J, Nalbant G, Almaqawi A, Kundakci B, Panniyammakal J, Heinrich M, Lewis SA, Greenfield SM, Tandon N, Biswas TK, Kinra S, Leonardi-Bee J. Effectiveness and Safety of Ayurvedic Medicines in Type 2 Diabetes Mellitus Management: A Systematic Review and Meta-Analysis. *Front Pharmacol*. 2022 Jun 8;13:821810.

15. Sane, R., A, SI., S, NM, Tejaswini, S., S, MR, (2018). To study efficacy of Comprehensive Diabetes Care (CDC) management program in type II diabetic obese patients: An observational study. *Int J Ayurveda Pharma Res*, 6 (6), 6–12.
16. Sane, R., Naik, M., Khair, K., Mahajan, H., Pawar, D., and Revandkar, V., et al. (2018). Effect of comprehensive diabetes care on glycaemic control with reduction in dependency of oral hypoglycaemic medicines in pre-obese diabetic patients: A retrospective study. *Int J Curr Res Acad Rev*, 6 (11), 73–81.
17. Tejaswini Nandakumar Shingan, Swati Kamlapure, Nilesh Kulthe. Role of Ayurveda Based Comprehensive Diabetic Care Program in Restoration of Euglycemia in Known Type 2 Diabetic Mellitus, *International Journal of Diabetes and Endocrinology*. 2022 7(3), 91–95.
18. Datey P, Hankey A, Nagendra HR. Combined Ayurveda and Yoga Practices for Newly Diagnosed Type 2 Diabetes Mellitus: A Controlled Trial. *Complement Med Res*. 2018;25(1):16-23.

TABLE/FIGURE LEGENDS

Table 1: Baseline demographic and anthropometric measurements

Table 2: Demographic and anthropometric measurements according to age, gender, HbA1c, and body mass index

Figure 1: Glucose tolerance testing

UNDER PEER REVIEW

Table 1: Baseline demographic and anthropometric measurements

Variables	Day 1	Day 90	p value
Age, years	50.51 ± 10.62		
Males	41 (65.1%)		
Weight, kg	70.70 ± 13.19	65.15 ± 9.83	<0.001
Body mass index	27.00 ± 3.88	24.91 ± 3.75	<0.001
HbA1c, %	7.87 ± 1.90	5.79 ± 0.51	<0.001

All data are expressed as number (percentage) or mean ± standard deviation

HbA1c – glycated haemoglobin

UNDER PEER REVIEW

Table 2: Demographic and anthropometric measurements according to age, gender, HbA1c, and body mass index


Variables	HbA1c Day 1	HbA1c Day 90	p value	Change %	Weight Day 1	Weight Day 90	p value	Change %	BMI Day 1	BMI Day 90	p value	Change %
Age, years												
25–45	7.94 ± 1.87	6.03 ± 0.50	<0.001	-1.91	75.08 ± 9.47	70.15 ± 9.87	<0.001	-4.92	27.29 ± 4.13	25.50 ± 4.20	<0.001	-1.79
45–65	7.94 ± 2.00	5.62 ± 0.46	<0.001	-2.32	68.42 ± 9.95	62.56 ± 9.53	<0.001	-5.86	26.80 ± 3.98	24.53 ± 3.61	<0.001	-2.27
>65	7.33 ± 1.77	5.86 ± 0.60	<0.001	-1.47	68.01 ± 5.55	62.04 ± 4.25	<0.001	-5.96	27.08 ± 2.85	24.91 ± 3.06	<0.001	-2.17
Gender												
Male	8.00 ± 1.93	5.91 ± 0.50	<0.001	-2.09	72.08 ± 10.15	66.97 ± 9.54	<0.001	-5.10	25.86 ± 3.47	24.06 ± 3.33	<0.001	-1.81
Female	7.62 ± 1.90	5.56 ± 0.47	<0.001	-2.06	68.13 ± 9.07	61.76 ± 9.91	<0.001	-6.37	29.12 ± 3.87	26.50 ± 4.13	<0.001	-2.62
HbA1c, %												
<5.7	5.14 ± 0.38	5.58 ± 0.73	<0.001	0.44	75.80 ± 9.19	76.12 ± 12.33	<0.001	0.32	28.90 ± 4.68	28.98 ± 5.51	<0.001	0.09
5.8–7.0	6.62 ± 0.32	5.65 ± 0.52	<0.001	-0.97	68.83 ± 9.49	63.63 ± 9.18	<0.001	-5.21	25.85 ± 2.87	24.00 ± 3.02	<0.001	-1.85
7.1–10	8.45 ± 0.88	5.85 ± 0.47	<0.001	-2.60	71.38 ± 10.64	64.55 ± 9.63	<0.001	-6.83	27.72 ± 4.44	25.05 ± 3.73	<0.001	-2.67
10.1–15.0	11.87 ± 1.10	6.19 ± 0.25	<0.001	-5.69	70.81 ± 9.07	64.88 ± 8.84	<0.001	-5.94	26.84 ± 4.00	24.59 ± 3.85	<0.001	-2.25
Body mass index												
Normal	7.78 ± 1.95	5.92 ± 0.53	<0.001	-1.85	62.15 ± 7.01	58.40 ± 5.17	<0.001	-3.75	22.61 ± 2.12	21.32 ± 1.45	<0.001	-1.29
Overweight	7.86 ± 1.55	5.76 ± 0.50	<0.001	-2.10	70.68 ± 6.10	65.34 ± 9.03	<0.001	-5.43	26.36 ± 0.97	24.35 ± 2.14	<0.001	-2.01
Obese 1	7.98 ± 2.43	5.52 ± 0.55	<0.001	-2.47	72.50 ± 9.39	64.28 ± 8.83	<0.001	-8.22	28.84 ± 0.52	25.66 ± 1.10	<0.001	-3.18
Obese 2	8.33 ± 1.26	6.05 ± 0.54	<0.001	-2.28	80.98 ± 12.24	72.68 ± 7.76	<0.001	-8.30	30.89 ± 0.40	27.89 ± 1.74	<0.001	-3.00
Morbid	87.69 ± 2.63	5.89 ± 0.35	<0.001	-1.80	82.54 ± 7.76	78.11 ± 10.36	<0.001	-4.42	34.40 ± 2.33	32.50 ± 3.12	<0.001	-1.90

obesity												
---------	--	--	--	--	--	--	--	--	--	--	--	--

All data are expressed as mean \pm standard deviation

UNDER PEER REVIEW

Figure 1: Glucose tolerance testing



UNDER PEER REVIEW