

**Blood pressure values in apparently healthy Sudanese population**

---

**Abstract**

**Introduction:** Blood pressure is one of the most often measured clinical parameters, and assessment of blood pressure has a considerable impact on diagnostic decisions.

**Objectives:** To establish blood pressure normal reference values in Sudanese.

**Methods:** A cross-sectional study was conducted from September 2016 to November 2018. Eight hundred eighty-eight healthy adult Sudanese between the ages of 18 and 60 (203 men and 685 women) were randomly selected from the states of Khartoum, Northern, Gezira, Red Sea, and North Darfur. Clinical, anthropometric, and blood pressure measurement data were collected.

**Results:** The mean for all volunteers was  $113.93 \pm 9.917$  mmHg, systolic blood pressure (SBP) and  $75.29 \pm 6.79$  mmHg, diastolic blood pressure (DBP). SBP in men was  $118.6 \pm 7.642$  mmHg compared to  $112.53 \pm 9.121$  mmHg in women, while DBP in men was  $77.51 \pm 5.984$  mmHg compared to  $74.63 \pm 6.844$  mmHg in women. Beside the gender variations, blood pressure values also showed geographical variability. There was a positive connection between blood pressures (SBP and DBP), BMI, and age. ( $P < 0.05$ ) was used for significance.

**Conclusion:** Blood pressure of Sudanese was found to be within the normal international range with gender and geographical variability. It showed positive correlation with age and BMI.

---

Key words: Blood pressure, Sudanese, Normal reference values

**Introduction:**

Blood pressure is the pressure within the major arterial system of the body and measured in millimeters of mercury. SBP is the maximum blood pressure during systole; DBP is the minimum pressure during diastole <sup>(1)</sup>. One in every two persons  $\geq 20$  years in the USA has hypertension with only 39.64% on medications having well-controlled hypertension <sup>(2)</sup>. High blood pressure is a major risk of cardiovascular and renal diseases. <sup>(3)</sup> Hypertension requires the heart to work harder than normal to

circulate blood through the blood vessels. <sup>(4)</sup> Hypertension is defined when SBP is equal to or above 140 mm Hg and/or a DBP equal to or above 90 mm Hg. <sup>(5)</sup> Globally, the prevalence of adult hypertension was estimated to be over 40% in 2008; this number is expected to rise to above 60% by 2025. <sup>(6)</sup> The association of a high BMI with cardiovascular disease is at least partly explained by the association of BMI with hypertension and elevated serum cholesterol. <sup>(7)</sup> Numerous factors, such as advanced age, smoking, black race, low potassium and high sodium intake, inactivity, alcohol intake, stress, some chronic illnesses, overweight, and obesity, among others, have an impact on normal blood pressure. <sup>(8)</sup> Davy and Hall reported a linear relationship between adiposity and hypertension. <sup>(9)</sup> In developing countries, high blood pressure is one of the risk factors for cardiovascular diseases, and the estimated 7.1million deaths especially among middle, and old-age adults is due to high BP. <sup>(10)</sup> To establish reference BP values for Sudanese and compare them to international values, a study was conducted on 1100 individuals in Wad Medani city, Sudan. The findings indicated that while the mean BP (SBP/DBP) for women was  $133 \pm 19/73 \pm 9$  mmHg, the mean for men was  $134 \pm 18/71 \pm 10$  mmHg. <sup>(11)</sup> According to a study done in Khartoum, Sudan, by Elnagi Y. Hago et al., the mean (SBP) is greater in men ( $120 \pm 8.8$ ) compared to women ( $113 \pm 9.7$ ) and the DBP of men ( $79.3 \pm 5.9$  mmHg) is higher than that of women ( $74.8 \pm 7.1$  mmHg). In all participants, BMI was significantly positively correlated with both SBP and DBP ( $P = 0.01$ ) <sup>(12)</sup>. The average blood pressure was within the international range, and men's blood pressure was noticeably higher than that of women.

**Rationale:** From the fact that high blood pressure is a risk of for cardiovascular diseases in developing countries and the evidences of ethnic and environmental variability in blood pressure values in regional communities, plus the need to establish correlation between BMI and age with blood pressure, It is important to study normal blood pressure values in Sudanese and to derive prediction equations based on BMI and age.

**Methods:**

A cross-sectional study done between 2016 and 2018 on 888 adult healthy Sudanese (685 females and 203 males), 18–60 years old, and the study covered the states of Khartoum (392 participants), Northern (130), Gezira (82), Red Sea (160), and North Darfur (124). All participants were selected randomly from students and employees at different universities all over Sudan. The inclusion criteria were age range between 18

and 60 years old, absence of chronic diseases, and absence of acute illness at the time of data collection. Sample size was determined based on Monkey-Survey computer equation. All volunteers had completed a questionnaire covering personal, habitual, and health data. Then anthropometric blood pressure measurements were performed. The auscultation method was used for blood pressure measurements using the mercury sphygmomanometer and the stethoscope. Five minutes separated the two readings. A third reading was taken if there was a 10 mmHg or greater discrepancy between the first two values. Validity and reliability of readings were checked, and the instruments were standardized using other standard ones. All People performing the tests were qualified doctors. The final reading was the average. Weight and height were measured to determine the body mass index (BMI) as weight in kilograms divided by height in square meters. Statistical Package for the Social Sciences (SPSS) was used to analyze the collected data, which was then be displayed in tables and graphs.

### Results:

**Table 1: Mean systolic and diastolic blood pressures in different states of Sudan.**

Parameter	SBP (Mean $\pm$ SD) in mmHg		DBP (Mean $\pm$ SD) in mmHg		N
State	Males (n = 203)	Females (n = 685)	Males (n = 203)	Females (n = 685)	
Khartoum	118.88 $\pm$ 6.23	113.10 $\pm$ 8.93	79.45 $\pm$ 5.28	75.08 $\pm$ 6.97	392
Red sea	120.88 $\pm$ 7.75	109.50 $\pm$ 8.49	77.65 $\pm$ 4.37	72.78 $\pm$ 6.54	160
Gezira	114.63 $\pm$ 8.76	114.82 $\pm$ 8.92	74.89 $\pm$ 6.23	77.27 $\pm$ 6.30	82
Northern	120.00 $\pm$ 8.66	114.70 $\pm$ 10.05	79.03 $\pm$ 6.38	74.75 $\pm$ 7.16	130
North Darfur	118.63 $\pm$ 7.69	111.51 $\pm$ 8.73	75.10 $\pm$ 5.79	74.08 $\pm$ 6.40	124
Total	118.59 $\pm$ 7.61	112.52 $\pm$ 9.14	77.54 $\pm$ 5.96	74.62 $\pm$ 6.88	888

Table (1) showed variations in systolic and diastolic blood pressures between different states of Sudan. The mean SBP in male Sudanese was found to be 118.59  $\pm$  7.61 mmHg while the mean DBP was found to be 77.54  $\pm$  5.96 mmHg. The female mean SBP was found to be 112.52  $\pm$  9.14 mmHg while the mean DBP was 74.62  $\pm$  6.88 mmHg. Red sea and Northern states showed the highest mean systolic blood pressures in males (120.88  $\pm$  7.75 and 120.00  $\pm$  8.66 mmHg respectively) while the

Red Sea state showed the least mean systolic blood pressure in females ( $109.50 \pm 8.49$  mmHg). Khartoum and Northern states showed the highest mean DBP values in males ( $79.45 \pm 5.28$  and  $79.03 \pm 6.38$  mmHg respectively). The least DBP ( $72.78 \pm 6.54$ ) was obtained by Red Sea states' females.

**Table 2: Mean SBP and DBP in both sexes with the age group.**

Gender	Males		Females SBP	
Age group	SBP (Mean $\pm$ SD)	DBP (Mean $\pm$ SD)	SBP (Mean $\pm$ SD)	DBP (Mean $\pm$ SD)
20-29	$118.26 \pm 7.874$	$76.99 \pm 5.977$	$111.74 \pm 8.436$	$74.18 \pm 6.710$
30-39	$119.64 \pm 7.712$	$78.93 \pm 7.641$	$114.32 \pm 10.144$	$77.57 \pm 8.140$
40-49	$119.36 \pm 6.172$	$80.57 \pm 3.715$	$118.77 \pm 10.144$	$79.10 \pm 6.228$
50-60	$121.36 \pm 5.519$	$79.55 \pm 4.719$	$127.81 \pm 10.641$	$78.13 \pm 5.737$
Total	$118.60 \pm 7.642$	$77.51 \pm 5.984$	$112.53 \pm 9.121$	$74.63 \pm 6.844$

From table (2) it was clear that both total mean systolic ( $118.6 \pm 7.642$ ) and diastolic ( $77.51 \pm 5.984$ ) blood pressures in males were higher than in females (SBP of  $112.53 \pm 9.121$  and DBP of  $74.63 \pm 6.844$ ), but in the group age of 50-60 years old SBP was higher in females. Both SBP and DBP increased with age in both males and females. The positive correlation between age and blood pressure was more clear in females compared to males.

**Table 3: Correlation matrix between blood pressure, age, and BMI.**


Variables	SBP	DBP	Age	BMI
SBP	1.000	0.633**	0.270**	0.253**
DBP	0.633**	1.000	0.202**	0.214**
Age	0.270**	0.202**	1.000	0.307**
BMI	0.253**	0.214**	0.307**	1.000

\*\* . Correlation is significant at the 0.01 level (2-tailed).


Table (3) Showed a statistically significant positive correlation between BMI and both systolic and diastolic blood pressure. This result showed that blood pressure increases along with BMI. The positive correlation between BMI and diastolic BP was less when compared to systolic BP. A strong and positive correlation between age and

both systolic and diastolic blood pressure was also found.

**Figure 1: The correlation between SBP and BMI among Sudanese.**



**Figure 2: The correlation between DBP and BMI among Sudanese.**



From both of figures (1) and (2), it was clear that both SBP and DBP had positive

correlation with BMI, but the positive correlation of BMI was more linear with SBP compared to DBP.

(a= Age in years, b=BMI)

**Table 4: Prediction equations of blood pressure in adult Sudanese males**

Locations	Prediction for SBP males	Prediction for DBP males
Sudan	$111.322 + 0.042 a + 0.276 b$	$73.614 + 0.112 a + 0.042 b$
Khartoum state	$110.044 + 0.045 a + 0.349 b$	$74.040 + 0.030 a + 0.210 b$
Red sea state	$110.623 - 0.259 a + 0.735 b$	$70.946 + 0.181 a + 0.084 b$
Gezira state	$109.445 - 0.041 a + 0.293 b$	$70.180 + 0.070 a + 0.142 b$
North state	$118.127 + 0.056 a + 0.017 b$	$75.430 + 0.076 a + 0.110 b$
North Darfur state	$111.648 + 0.153 a + 0.138 b$	$72.276 + 0.123 a - 0.014 b$

**a = Age, b = BMI**

**Table 5: Prediction equations of blood pressure in adult Sudanese females**

Locations	Prediction for SBP females	Prediction for DBP females
Sudan	$96.695 + 0.258 a + 0.429 b$	$64.906 + 0.090 a + 0.333 b$
Khartoum state	$96.882 + 0.319 a + 0.370b$	$65.383 + 0.125 a + 0.292 b$
Red sea state	$99.718 - 0.066 a + 0.491 b$	$60.660 + 0.101 a + 0.430 b$
Gezira state	$96.150 + 0.215 a + 0.548 b$	$67.911 - 0.110 a + 0.527 b$
North state	$100.866 + 0.051 a + 0.554 b$	$67.897 - 0.036 a + 0.335 b$
North Darfur state	$97.858 + 0.198 a + 0.415 b$	$62.708 + 0.208 a + 0.301 b$

**a = Age, b = BMI**

Making use of the positive correlation between blood pressures and age, and blood pressures and BMI, predictive equations for adult males and females Sudanese from all over Sudan were derived using multiple regression.

## **Discussion:**

The previous trials to establish blood pressure normal reference values for Sudanese had limitations because of the small sample size and the lack of coverage of the whole country, but these limitations had been managed in this study.

Our sample size was designed to be statistically representative of the entire Sudanese population, considering all regions. Our results showed that, Sudanese normal blood pressure values were like the international blood pressure norms. One of the limitations in our study was that males sample size was less than females, but the overall sample was statistically representative. Also, there was mismatching of the sample size between the different states considering their population density. Great deal of the sample size was taken from Khartoum, since the researchers stayed more time there, but the positive point was that population of Khartoum were a mixture of all Sudan regions. The regional variability of blood pressure values was investigated by Kiefe CI et al. <sup>(13)</sup>.

Our result in tables (1) Shows that the mean of systolic blood pressure was differ from state to state the highest blood pressure was found in Northern state ( $115.96 \pm 9.96$  mmHg) while the lowest SBP was found in the Red sea state ( $110.71 \pm 9.10$  mmHg), also the mean of diastolic blood pressure was differs from state to state the highest diastolic blood pressure was found in Gezira state ( $76.49 \pm 6.34$  mmHg) while the lowest SBP was found in the Red sea state ( $73.29 \pm 6.51$  mmHg), this result was same as study done Kiefe CI and et al, that illustrate there was different result in different region of same country. <sup>(13)</sup> This variation in the blood pressure may be correlated with socioeconomic class and geographic location due to the emergence of new risk factors for hypertension. similar to prenatal malnutrition and psychological stress, may disproportionately impact the poor. <sup>(14,15)</sup> In our study in tables (2,3) we found there was relationship between blood pressure, sex and age. The males had high blood pressure than females, but at the age above the fifty females become had high systolic this result agree with Jane F. Reckelhoff, he proven that at similar ages, men have higher blood pressure than women. However, after menopause, women's blood pressure rises to levels that are much greater than in men. <sup>(16)</sup> Increases in androgens in people and animals may be the cause of the higher prevalence of hypertension in men than in women. Women with increased testosterone levels associated with polycystic ovarian syndrome or adrenal virilizing tumors have hypertension. <sup>(17-19)</sup> Nitric oxide (NO) generation has been demonstrated to be stimulated by estrogen.

<sup>(20,21)</sup> As a result, the loss of estrogen associated with menopause may contribute to the rise in blood pressure in women and this explain why women high blood pressure than men.

BMI and both systolic and diastolic blood pressure were positively correlated with age. Systolic and diastolic blood pressures increased steadily with age, from the youngest to the oldest age groups, demonstrating the age dependence of BP. Previous research has also linked high blood pressure to age. <sup>(22-24)</sup> In my current study, the relationship between age and systolic and diastolic blood pressures was less strong than the relationship between age and BMI. Thus, the relationship between systolic and diastolic blood pressure and age was less strong than the relationship between BMI and age.

Blood pressure was higher in the elderly strata of the sample with or without a corresponding increase in BMI, implying the presence of other contributory factors to hypertension besides fat accumulation alone. A significant correlation between BP and age was also reported in a study from North India. <sup>(25)</sup>

We found that systolic blood pressure and diastolic blood pressure were highest in overweight subjects, intermediate in normal weight subjects, and lowest in underweight subjects in both males and females and this result agree with other studies. <sup>(26,27)</sup> This result could be attributed to differences in sympathetic tone between underweight and overweight individuals. However, plus pressure was comparable. Assuming that arterial compliance did not differ between groups, this implies that stroke volume is also comparable. Heart rate was also comparable. As a result, we have indirect evidence that cardiac output does not differ significantly between the groups. Thus, variations in blood pressure may be largely due to variations in total peripheral resistance, which is heavily influenced by tonic sympathetic control of resistance vessels. <sup>(28)</sup>

### **Conclusion:**

The derived blood pressure values of Sudanese in our study were not significantly different from international values. It showed no significant variations between different parts of the country. Significant positive correlations were found between blood pressure and BMI and between blood pressure and age. So, prediction equations based on BMI and age were derived.

**Ethical Approval and Consent:**

Ethical approval was issued from the Federal Ministry of Health in Sudan (FMOH). All study participants have received an explanation of the study's goals. Each participant had signed a written consent after being informed about the study.

UNDER PEER REVIEW

## References:

1. Brzezinski WA. Blood Pressure. In: Walker HK, Hall WD, Hurst JW, editors. *Clinical Methods: The History, Physical, and Laboratory Examinations*. 3rd ed. Boston: Butterworths; 1990. Chapter 16. PMID: 21250111.
2. Muchi Ditah Chobufo et al. (2020). The prevalence and clinical significance of nocturnal hypertension in pregnancy. *Int J Cardiol Hypertens*. 2020 Sep; 6: 100044. doi: [10.1016/j.ijchy.2020.100044](https://doi.org/10.1016/j.ijchy.2020.100044)
3. Lewington S, Clarke R, Qizilbash N. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002; 360(9349): 1903–1913.
4. Carretero, O.A, Oparil, S. Essential hypertension. Part I: definition and setiology. *Circulation*(2000); 101 (3): 329–33
5. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo Jr JL, Jones DW, Materson BJ, Oparil S, Wright Jr JT, Roccella EJ. Seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. *hypertension*. 2003 Dec 1;42(6):1206-52.
6. Angeli F, Reboldi G, Verdecchia P. The 2014 hypertension guidelines: implications for patients and practitioners in Asia. *Heart Asia*. 2015 Jul 1;7(2):21-5.
7. Chan LCK, Ware RS, Kesting J, Marczak M, Good D, Shaw JTE. Association Between Anthropometric Measures of Obesity and Insulin Resistance in a Self-Selected Group of Indigenous Australians. *Hear Lung Circ*. 2007;16(4):303–4.
8. WALTON CH. High blood pressure. *Manit Med Rev*. 1947;27(4):212–6.
9. Davy KP, Hall JE (2004). Obesity and hypertension: two epidemics or one? *.Am J Physiol Regul Integr Comp Physiol* ;286:803–813.
10. World Health Organization. *The world health report 2002: reducing risks, promoting healthy life*. World Health Organization; 2002.

11. Omima S A E. normal Sudanese values of blood pressure among apparently healthy Sudanese age 50 years and above in Wad Madni City. A thesis submitted in partial fulfillment for the requirements of MSc degree in human physiology. B.V.Sc University of Khartoum 2003.
12. Hago EY, Bashir AA, Abdalla IM et.al. Normal blood pressure in adult Sudanese in Khartoum state, Sudan. *Int J Health Sci Res.* 2020; 10(10):161-165
13. Kiefe CI, Williams OD, Bild DE, Lewis CE, Hilner JE, Oberman A. Regional disparities in the incidence of elevated blood pressure among young adults: the CARDIA Study. *Circulation.* 1997;96:1082–1088.
14. Reddy KS, Yusuf S: Emerging epidemic of cardiovascular disease in developing countries. *Circulation* 1998, 97(6):596-601.
15. Pearson TA: Cardiovascular disease in developing countries: myths, realities and opportunities. *Card Drugs Therapy* 1999, 13:95-104
16. Reckelhoff JF. Gender differences in the regulation of blood pressure. *Hypertension.* 2001 May;37(5):1199-208.
17. Soranno D, Prasad V, Oberfield DR, Greco A, Sivaraman N, Drucker W. Hypertension and virilization caused by a unique dexamethasone and androgen-secreting adrenal adenoma. *J Pediatr Endocrinol Metab.* 1999;12:215–220.
18. Talbott E, Guzick D, Clerici A, Berga S, Detre K, Weimer K, Kuller L. Coronary heart disease risk factors in women with polycystic ovary syndrome. *Arterioscler Thromb Vasc Biol.* 1995;15:821– 826.
19. Mattson LACG, Hamberger L, Samsioe G, Silfverstolpe G. Lipid metabolism in women with polycystic ovary syndrome: possible implications for an increased risk of coronary heart disease. *Fertil Steril.* 1984;42: 579–584.
20. Weiner CP, Lizasain I, Baylis SA, Knowles RG, Charles IG, Moncada S. Induction of calcium dependent NO synthase by sex hormones. *Proc Natl Acad Sci U S A.* 1994;91:5212–5216.

21. Goetz RM, Morano I, Calovini T, Studer R, Holtz J. Increased expression of endothelial constitutive NO synthase in rat aorta during pregnancy. *Biochem Biophys Res Commun.* 1994;205:905–910.
22. J. I. Schall, “Sex differences in the response of blood pressure to modernization,” *American Journal of Human Biology*, vol. 7, pp. 159–172, 1995.
23. W.W. Dressler and J. R. Bindon, “Social status, social context, and arterial blood pressure,” *American Journal of Physical Anthropology*, vol. 85, no. 18, p. 82, 1994.
24. V. Suman and S. Kapoor, “Effect of cold stimulus on blood pressure,” in *Contemporary Studies in Human Ecology: Human Factor, Resource Management and Development*, M. K. Bhasin and S. L. Malik, Eds., pp. 349–354, Kamle Raj Enterprises, 1998.
25. R. B. Singh, S. S. Rastogi, V. Rastogi et al., “Blood pressure trends, plasma insulin levels and risk factors in rural and urban elderly populations of north India,” *Coronary Artery Disease*, vol. 8, no. 7, pp. 463–468, 1997.
26. Mungreiphy NK, Kapoor S, Sinha R. Association between BMI, blood pressure, and age: study among Tangkhul Naga tribal males of Northeast India. *Journal of Anthropology.* 2011;2011.
27. Dua S, Bhuker M, Sharma P, Dhall M, Kapoor S. Body mass index relates to blood pressure among adults. *North American journal of medical sciences.* 2014 Feb;6(2):89.
28. Sorof J, Daniels S. Obesity hypertension in children. A problem of epidemic proportions. *Hypertension* 2002; 40: 441–447.