

Role of Cytokinin Oxidase/Dehydrogenase in Crop Improvement

Abstract

The phytohormone Cytokinin regulates various crucial functions of plant growth and development such as shoot apical meristem activity, flower development, vascular development etc. and also has negative role in lateral root development and root apical meristem activity. Cytokinin is degraded by cytokinin oxidase/dehydrogenase (*CKX*) irreversibly and reversible inactivation by glucosylation. Various homologous genes of *CKX* have been identified in Arabidopsis and in many other crop plants. The root specific expression of *CKX* showed positive results for larger root with higher lateral root numbers and higher root to shoot biomass. The site-specific manipulation of *CKX* gene has been done in crops like wheat, barley, maize, rice, legumes and horticultural crops to regulate the cytokinin levels in the particular tissues to enhance the yield, tolerance to various abiotic stresses (drought, salt, lower soil fertility), biofortification for many micronutrients (Zn, Fe) in the seed, propagation, etc. And the use of modern techniques in manipulation of *CKX* have been started to improve the crop plant and the outcome is very promising for the future application in commercial agricultural activities.

Key words: Cytokinin, cytokinin oxidase/dehydrogenase, drought, lateral root, yield

1. Introduction

The demand for the agricultural crop is increasing day by day as the global population increases (Godfay *et al.*, 2010). The current situation of the world is about 815 million people were undernourished reported by United Nations Food and Agriculture Organizations (FAO, 2016). The main cause for this undernourishment is the protein-energy malnutrition and micronutrient deficiency in food (UNICEF, 2015). In the current situation where the land is decreasing due to population outbreak and food demand is increasing, the more intense use of cropland could meet the demands but due to unpredictable and changing environment condition the problem is becoming worse (Godfray *et al.*, 2010). Natural hazards like drought, flood and biotic stresses are limiting the crop production (FAO, 2019).

In the current scenario of human population, the improvement in the majorly grown crop like cereals, legumes, oilseeds etc., faces numerous abiotic stresses during their life

cycle, is the solution for feeding the growing population (Swaminathan and Bhawani, 2013). The plants with larger root system can withstand in the drought and low nutrient soil conditions (Brown *et al.*, 1989; Saxena *et al.*, 1996; Kondo *et al.*, 1999; Steel *et al.*, 2006; Henry, 2012). For this, the cytokinin oxidase/dehydrogenase enzyme can be an important player and can be exploited to improve the crops productivity. This enzyme degrades the phytohormone cytokinin in specific tissue at specific time in the plant. It has been reported by several scientists that cytokinin negatively regulated the lateral root formation (Goodwin and Morris, 1979; Wightman *et al.*, 1980) and the mutant plant of Arabidopsis defective response for cytokinin showed the high branching of roots (Mason *et al.*, 2005; Reifler *et al.*, 2006; To *et al.*, 2008).

Cytokinin level in plants is regulated by various processes like reversible glycosylation, action of adenine phosphoribosyl transferase and also degradation by the cytokinin oxidase/dehydrogenase (*CKXs*) irreversibly (Sakakibara, 2006). The manipulation of level of cytokine in the barley, *Populus* and rape was used to improve the agronomic value by using the variants of cytokinin dehydrogenase (Ramireddy *et al.*, 2018a; Nehnevajova *et al.*, 2019) and lead to improve root growth and drought tolerance. Recently Khandal *et al.* (2020) reported that the increased activity of *CKX* in chickpea roots enhances the drought tolerance, seed yield and also concentration of micronutrients like zinc, iron, potassium and copper without compromising with the protein level in seed. The cytokinin is necessary for seed development (Letham, 1963) and it was also reported by Ashikari *et al.* (2005) in rice. In wheat the concentration of endogenous cytokinin was found high at the time of rapid endosperm nuclear and cell division in developing seed (Bennet *et al.*, 1973). The overexpression of *AtCKX3* reduces the formation of primordium in floral meristem leading lower number of flowers (Werner *et al.*, 2003).

In this review article, the information regarding cytokinin and its regulation through *CKX* at different stages and in different crops has collected and compiled for better understanding of *CKX* functions which can help in developing the crops with higher nutrition quality and can grow under limited water condition and low fertile soil without compromising the seed yield.

1.1 Cytokinin and its metabolism

The plant hormone cytokinin is synthesized by two processes either by an isopentenyltransferase (IPT) attaching on isoprenoid side chain of adenosine triphosphate or

adenosine diphosphate (ATP/ADP) synthesizing nucleotide of isopentenyl adenine (iP) and *tans-zeatin* (tZ) or by a tRNA-IPT leading, indirectly to the *cisZ*-type cytokinin. Cytokinin is activated by the LONELY GUY (LOG) by releasing the free base from the nucleotide forms. Cytokinin oxidase/dehydrogenase are the key player in destruction of cytokinin. O-glucosylation or N-glucosylation both can inactivate the cytokinins (Jameson, 2017; Kieber and Schaller, 2018; Worthen *et al.*, 2019; Chen *et al.*, 2020).

1.2 Role of cytokinin in growth and development

1.2.1 In the shoot tissues

In the shoots, the development of shoot apical meristem (SAM) is controlled by various factors like transcriptional factors, external signals and phytohormones specially cytokinins (Skoog and Miller, 1957). Werner *et al.*(2001; 2003) and Wybouw and De Rybel(2019) reported the highly decreased SAM in the plant deficient in cytokinin and suggested that cytokine is crucially required by the plant for developing and maintaining SAM. Similar observations were reported by various workers (Miyawaki *et al.*, 2006; Nishimura *et al.*, 2004; Higuchi *et al.*, 2004; Riefler *et al.*, 2006) in their experiments by mutating the multiple receptor or *IPT*gene which suggested that the cytokinin positively regulate the SAM activity (Waidmann *et al.*, 2019).

The growth of vascular cambium and radial growth of plant is also controlled by the cytokinin (Nieminen *et al.*, 2008; Matsumoto *et al.*, 2008). Experiment of Nieminen *et al.* (2008) in poplar and birch trees found that the genes of receptors of cytokinin and other cytokinin signaling genes were expressed in cambial zone, and also found reduced cambial activity in the transgenic plant made by targeted expression of *CKX* genes in the cambial cells.

Cytokinin has an antagonistic activity to the auxin in shoot by promoting the growth of axillary buds, in *Physcomitrella* iPR and extracellular iP are the common cytokinin required to induce bud formation (Schwartzenberg *et al.*, 2007).

The sink-source relations and leaf senescence is also regulated by cytokinin concentration in plant (Werner and Schmulling, 2009). This was reported by Werner *et al.* (2008) in tobacco plant deficit in cytokinin leads to reduction in chlorophyll synthesis and lower in sugar content and increased amount of starch in the shoot and also reduction in invertase activity in vacuole.

1.2.2 In the root tissues

Cytokinin positively regulate the vascular differentiation in the root meristem. The work on characterization of *wooden leg* (*wol*) allele of *CRE1/AHK4* (a dominant-negative allele), Mahonen *et al.* (2000) found that cytokinin is necessary for procambial cell division during embryogenesis to ensure vascular differentiation.

Cytokinin play a role in root apical meristem (RAM) unlike to SAM development, it negatively regulate the activity of RAM, this was proven by various scientist as their results suggest that plant develop a large RAM and rapid growth of roots in the condition where cytokinin deficit or lower signal output (Werner *et al.*, 2003; Mason *et al.*, 2005; Miyawaki *et al.*, 2006; Reiflier *et al.*, 2006; Heyl *et al.*, 2008; Zou *et al.*, 2018).

The lateral root formation is negatively regulated by cytokine; this was observed in several experiments by making transgenic and mutant plants (Werner *et al.*, 2003; Lohar *et al.*, 2004; To *et al.*, 2004; Mason *et al.*, 2005; Reiflier *et al.*, 2006). The lateral root formation was inhibited by cytokinin by blocking the first division of xylem-pole pericycle cells only (Laplaze *et al.*, 2007; Li *et al.*, 2006).

2. Characterization and utilization of cytokinin oxidase/dehydrogenase

Cytokinin oxidase/dehydrogenase is a key enzyme which regulates the cytokinin level in the plant, this was reported by various researchers in different crops- *Arabidopsis thaliana* (Werner *et al.*, 2003), maize (Brugiere *et al.*, 2003), rice (Ashikari *et al.*, 2005), pea (Gaudinova *et al.*, 2005) barley (Zalewski *et al.*, 2014), foxtail millet (Wang *et al.*, 2014), *Fragaria vesca* (Jiang *et al.*, 2016), and wheat (Ogonowska *et al.*, 2019). Cytokinin oxidase/dehydrogenase is encoded by a small gene family, in *Arabidopsis thaliana* seven homologous genes are present (Mameaux *et al.*, 2012). Till date it has not been reported in chickpea, only *CaCKX6* was characterized by Khandal *et al.* (2020). Summary of some crop improvement works by utilizing the *CKX* manipulation is presented in table 1.

2.1 In model plant *Arabidopsis thaliana*

The *Arabidopsis* plant has seven homologous of *CKX* gene and the tissue specific expression of them is different, Werner *et al.*, (2003) reported that *AtCKX1* and *AtCKX2* expressed in young tissues like shoot apex, *AtCKX4* expressed in stomatal precursor cells, root caps and in cells of trichomes, *AtCKX5* expressed in the procambial region of root

meristem, *AtCKX6* expressed in vascular tissues and *AtCKX3* expression was associated with cell cycle.

The *CKX7* was reported as localized in cytosol (Schmulling *et al.*, 2003) this was experimentally demonstrated by Kollmer *et al.* (2014), they fused the green fluorescent protein (GFP) in *CKX7* at c-terminus and this construct was expressed under control of Cauliflower mosaic virus 35S promoter in Arabidopsis plant. The fluorescence signal was detected under confocal microscopy and reported the expression was found in cytosol.

Overexpression of *AtCKX* under control of 35S promoter in Arabidopsis was analysed and found that rate of root elongation was 70-90% more than wild type in 35S:*AtCKX1* and 35S:*AtCKX3* expressing seedling and the lateral root formation was also higher than the wild type (Werner *et al.*, 2003).

Overexpression of *CKX* in Arabidopsis plant was examined for the stress tolerance and reported a higher salt stress tolerance in the plant overexpressing *CKX1* among all other *CKX*. And the drought stress tolerance was also observed in these mutant plant and found more drought tolerant than wild type (Nishiyama *et al.*, 2011).

2.2 In monocot plants

Five different type of *ZmCKX* genes have been characterized in maize (Morris *et al.*, 1999; Bilveu *et al.*, 2003). In another experiment, the maximum expression of *ZmCKX* was found in kernels, tassels and ears as detected by semi-quantitative real time PCR reported by Massonneau *et al.* (2004).

In barley, the real time PCR data analysis for *HvCKX* showed that the *HvCKX1* expressed in mature kernels, roots and leaves, while *HvCKX2* expression was found in roots and leaves but the *HvCKX3* was predominantly expressed in mature kernels (Galuszka *et al.*, 2004). The over expression of *HvCKX* under constitutive expressed 35S promoter was carried out in Arabidopsis and tobacco plant and found shorter internode and dwarf habit along with larger root system (Galuszka *et al.*, 2004). Root specific expression of *CKX1* or *CKX2* under control of root specific promoter pEPP, pPEP and pRET by *Agrobacterium* mediated transformation in barley was done and observed that these mutant line showing tolerance under drought condition and also accumulate more minerals (Ramireddy *et al.*, 2018a). The analysis of transgenic plants of barley expressing *CKX* specifically in root accumulate higher Zn concentration having larger root system (Ramireddy *et al.*, 2018b). *HvCKX1* and *HvCKX3*

knockout mutants of barley plant was analysed and reported that lower number of grains and grain weight observed in *CKX3* mutant lines (Gasparis *et al.*, 2019).

In wheat, the RNA interference mediated gene silencing done for *TaCKX2.2.1-3A* and the mutant plants were screened for yield traits. The results showed that grain number per spike was enhanced as the expression of *TaCKX2.2.1-3A* reduces in T3 (Li *et al.*, 2018) and the grain weight was found associated with *TaCKX2.2.1-3D* (Zhang *et al.*, 2012b). The crown root growth was found associated with the expression of *TaCKX* in the limited water environment (Ogonowska *et al.*, 2019)

Ashikari *et al.* (2005) reported in rice that the *Gn1a* QTL is associated with the increase in yield which is a gene for *CKX2*. Overexpression of this gene produces more yield but downregulation has opposite impact on yield. Li *et al.* (2016) used gene editing tool to change the form of *OsCKX2* gene resulted in increased number of flowers and panicle size. Enhanced expression of *OsCKX4* in an enhancer mutant line of rice showed that crown root initiation required the activity of *OsCKX4* as recorded more crown roots, more root growth and strong root gravitropic response. Over expression of *OsCKX4* mutant lines showed more crown root growth and RNAi mutant lines of *OsCKX4* produced lesser crown root growth (Gao *et al.*, 2014).

2.3 In dicot plants

In tobacco plants the *CKX1* was overexpressed under root specific promoter (Werner *et al.*, 2010) and reported an enlarged root system which provided tolerance to drought stress. In tomato plants, the accumulation of hydrogen peroxide (H_2O_2) is regulated by cytokinin, the H_2O_2 involved in the degradation of chlorophyll. The protein gel blot analysis represents that the *CKX35* was found in chlorotic leaves and reduces the cytokinin levels and hence increased amount of H_2O_2 and in the green leaves the *CKX37* was predominantly present which is associated with active cytokinin concentration and normal hydroperoxide level in the leaves (Cueno *et al.*, 2012).

In strawberry (*Fragaria vesca*), the real time PCR analysis showed that the expression of *FvCKXs* was higher under drought, salt stress, heat stress and ABA treatment in roots, leaves and young fruits, this suggests that cytokinin has a negative role in growth and development under the stress conditions (Jiang *et al.*, 2016).

Liu *et al.* (2013) reported 12 *BrCKX* and 13 *BrIPT* genes in chinese cabbage and the stress related stimuli were found in the promoter region of these genes detected through transcription level analysis and they confirmed that these *BrCKX* and *BrIPT* genes have role in drought and salinity stress. In oilseed rape 23 *BnCKX* genes were identified (Liu *et al.*, 2018) and real time PCR results suggests that *BnCKX5-1*, *5-2,6-1* and *7-1* could be associated with the pod development and length of siliqua length. The analysis of transgenic oilseed rape plants with overexpression of *CKX2* represent the higher root-shoot biomass and accumulation of higher concentration of various micro and macro nutrients like P, Ca, Mg, S, Zn, Cu, Mo and Mn. The transgenic plant showed enhanced growth of chlorophyll under S and Mg deficiency and plants were able to extract more amount of Cd and Zn from contaminated medium and soil (Nehnevajova *et al.*, 2019).

For horticultural and silvicultural crops, grafting is an important technique for propagation but a problem during grafting is very common that undesirable lateral bud emergence reduces the grafting efficiency. To overcome of this problem Li *et al.* (2017) develop a transgenic with expression of tryptophan-2-monooxygenase (*iaaM*) under root specific promoter (*SbUGT*), this arrangement solved the problem of success of grafting but the root elongation biomass were reduced. They develop one more transgenic using *CKX* under same promoter, the root biomass was found higher. They made the cross of these two and reported that the negative effect due to *iaaM* was neutralized by activity of *CKX* and this made the grafting successful without compromising root development. The promoter *SbUGT* was used to express the *AtCKX2* in *Populus* which reduces the root sprouts in field condition resolving the problem of root-sprout mediated transgene spread (Li *et al.*, 2019).

2.4 In legumes

It was reported by Lohar *et al.* (2004) that the *AtCKX3* under constitutive expression positively regulates the nodulation in legumes demonstrated in *Lotus japonicus*. In *Medicago truncatula* Jardinaud *et al.* (2016) expressed *AtCKX3* under epidermal promoter pEPI in medicago and found upregulation of nodulation factor when inoculated with *S. meliloti* and when it expressed with cortex-specific promoter (pCO) the infection was reduced (Rival *et al.*, 2012).

The cellular localization of *CKX3* was reported in growing nodules and cortical cell division of the nodule primordium for this the promoter of *CKX3* was fused with YFP

(pCKX::YFP) and the fluorescence signal was recorded in *Lotus japonicum*. Phenotypic evaluation of *CKX3* mutant plants indicates that *CKX3* activity positively regulates the nodulation and the activity of cytokinin dose opposite function (Reid *et al.*, 2016).

Le *et al.* (2012) reported 14 *GmIPT* and 17 *GmCKX* genes in soybean and comparative analysis of promoter sequence of *GmCKX* genes with Arabidopsis genome and suggested that *GmCKX* were related with abiotic stress like drought.

Recently Khandal *et al.* (2020) used the chickpea root specific promoter (*CaWRKY31*) to express the chickpea cytokinin oxidase/dehydrogenase 6 (*CaCKX6*) in *Arabidopsis thaliana* and chickpea. Analysis of the transgenic plants represent increase in lateral root number and root to shoot biomass, and these transgenic lines withstand in long term drought condition without compromising with nodulation and nitrogen fixation. Their results also clearly indicates that those line produced upto 25% more yield and also having more Zn, Fe, K and Cu in the seeds with similar protein content.

3. Conclusion and future prospective

In the past the understanding related to cytokinin oxidase/dehydrogenase (*CKX*) has been significantly improved. From the above mentioned data it is clear that *CKX* enzyme has a key role in crop growth and development, it can regulate cytokinin level which play very crucial role in meristematic activity, root development, seed formation, chlorophyll concentration etc. Targeted manipulation of *CKX* provides the energy to the plant to withstand under stress conditions and enhancement in the yield with more nutrition values. In future the use of *CKX* manipulation through breeding programme or through editing tool can be done for the crop improvement for the majorly grown crops like cereals, pulses, oilseeds to feed the growing population and *CKX* can contribute to “new green revolution” (Ashikari *et al.*, 2005). Now the future challenge for researcher is to find out natural electron acceptors of *CKX* enzyme.

List of abbreviation

ABA: Abscisic Acid
CKX: Cytokinin Oxidase/Dehydrogenase
H₂O₂: Hydrogen Peroxide
iP: Isopentyl Adenine
PCR: Polymerase Chain Reaction
QTL: Quantitative Trait Locus

RAM: Root Apical Meristem
SAM: Shoot Apical Meristem
YFP: Yellow Florescence Protein

References

- Ashikari, M., Sakakibara, H., Lin, S., Yamamoto, T., Takashi, T., Nishimura, A., Angeles, E.R., Qian, Q., Kitano, H. and Matsuoka, M.** (2005). Cytokinin oxidase regulates rice grain production. *Science*, 309(5735): 741-745
- Bennett, M.D., Rao, M.K., Smith, J.B. and Bayliss, M.W.** (1973). Cell development in the anther, the ovule, and the young seed of *Triticum aestivum* L. var. Chinese Spring. Philosophical Transactions of the Royal Society of London. B, *Biological Sciences*, 266(875): 39-81.
- Bilyeu, K.D., Cole, J.L., Laskey, J.G., Riekhof, W.R., Esparza, T.J., Kramer, M.D. and Morris, R.O.** (2001). Molecular and biochemical characterization of a cytokinin oxidase from maize. *Plant Physiology*, 125(1): 378-386.
- Brown, S.C., Gregory, J., Cooper, J.M. and Keatinge, J.D.H.** (1989). Root and shoot growth and water use of chickpea (*Cicer arietinum*) grown in dryland conditions: effects of sowing date and genotype. *The Journal of Agricultural Science*, 113(1): 41-49.
- Brugière, N., Jiao, S., Hantke, S., Zinselmeier, C., Roessler, J.A., Niu, X., Jones, R.J. and Habben, J.E.** (2003). Cytokinin oxidase gene expression in maize is localized to the vasculature, and is induced by cytokinins, abscisic acid, and abiotic stress. *Plant Physiology*, 132(3): 1228-1240.
- Chang, C., Lu, J., Zhang, H., Ma, C.X. and Sun, G.** (2015). Copy number variation of cytokinin oxidase gene *TACKX4* associated with grain weight and chlorophyll content of flag leaf in common wheat. *PLoS One*, 10(12).
- Chen, L., Zhao, J., Song, J. and Jameson, E.** (2020). Cytokinin dehydrogenase: a genetic target for yield improvement in wheat. *Plant Biotechnology Journal*, 18(3): 614-630.
- Cueno, M.E., Imai, K., Ochiai, K. and Okamoto, T.** (2012). Cytokinin dehydrogenase differentially regulates cytokinin and indirectly affects hydrogen peroxide accumulation in tomato leaf. *Journal of Plant Physiology*, 169(8): 834-838.

- Duan, J., Yu, H., Yuan, K., Liao, Z., Meng, X., Jing, Y., Liu, G., Chu, J. and Li, J.** (2019). Strigolactone promotes cytokinin degradation through transcriptional activation of Cytokinin Oxidase/Dehydrogenase 9 in rice. *Proceedings of the National Academy of Sciences*, 116(28): 14319-14324.
- FAO.** (2019). Food and Agriculture Organizations.
- FAOSTAT.** (2016). FAO Statistical Database FAOSTAT.
- Galuszka, Frébortová, J., Werner, T., Yamada, M., Strnad, M., Schmülling, T. and Frébort, I.** (2004). Cytokinin oxidase/dehydrogenase genes in barley and wheat: cloning and heterologous expression. *European Journal of Biochemistry*, 271(20): 3990-4002.
- Gao, S., Fang, J., Xu, F., Wang, W., Sun, X., Chu, J., Cai, B., Feng, Y. and Chu, C.** (2014). Cytokinin oxidase/dehydrogenase E4 integrates cytokinin and auxin signaling to control rice crown root formation. *Plant Physiology*, 165(3): 1035-1046.
- Gasparis, S., Przyborowski, M., Kala, M. and Nadolska-Orczyk, A.** (2019). Knockout of the *HvCKX1* or *HvCKX3* Gene in Barley (*Hordeum vulgare* L.) by RNA-Guided Cas9 Nuclease affects the regulation of cytokinin metabolism and root morphology. *Cells*, 8(8): 782.
- Gaudinová, A., Dobrev, I., Šolcová, B., Novák, O., Strnad, M., Friedecký, D. and Motyka, V.** (2005). The involvement of cytokinin oxidase/dehydrogenase and zeatin reductase in regulation of cytokinin levels in pea (*Pisum sativum* L.) leaves. *Journal of Plant Growth Regulation*, 24(3): 188-200.
- Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M. and Toulmin, C.** (2010). Food security: the challenge of feeding 9 billion people. *Science*, 327(5967): 812-818.
- Goodwin, B. and Morris, S.C.** (1979). Application of phytohormones to pea roots after removal of the apex: effect on lateral root production. *Functional Plant Biology*, 6(2), 195-200.
- Henry, A.** (2012). IRRI's drought stress research in rice with emphasis on roots: accomplishments over the last 50 years. *Plant Root*, 7: 92-106.

- Heyl, A., Ramireddy, E., Brenner, W.G., Riefler, M., Allemeersch, J. and Schmölling, T.** (2008). The transcriptional repressor *ARR1-SRDX* suppresses pleiotropic cytokinin activities in Arabidopsis. *Plant Physiology*, 147(3): 1380-1395.
- Higuchi, M., Pischke, M.S., Mähönen, A., Miyawaki, K., Hashimoto, Y., Seki, M., Kobayashi, M., Shinozaki, K., Kato, T., Tabata, S. and Helariutta, Y.** (2004). In planta functions of the Arabidopsis cytokinin receptor family. *Proceedings of the National Academy of Sciences*, 101(23): 8821-8826.
- Holubová, K., Hensel, G., Vojta, Tarkowski, Bergougnoux, V. and Galuszka** (2018). Modification of barley plant productivity through regulation of cytokinin content by reverse-genetics approaches. *Frontiers in Plant Science*, 9: 1676.
- Jameson, E.** (2017). Cytokinins. In: Encyclopedia of Applied Plant Sciences (2nd Edition) Volume 1 (Thomas, B., Murphy, D.J. and Murray, B.G., eds), *Elsevier* B.V391–402..
- Jardinaud, M.F., Boivin, S., Rodde, N., Catrice, O., Kisiala, A., Lepage, A., Moreau, S., Roux, B., Cottret, L., Sallet, E. and Brault, M.** (2016). A laser dissection-RNAseq analysis highlights the activation of cytokinin pathways by Nod factors in the *Medicago truncatula* root epidermis. *Plant Physiology*, 171(3): 2256-2276.
- Jiang, Y., Mi, X., Lin, Y., Wu, H., Gu, T., Ding, J. and Li, Y.** (2016). Evolution and expression patterns of cytokinin oxidase genes in *Fragaria vesca*. *Scientia Horticulturae*, 212: 115-125.
- Khandal, H., Gupta, S.K., Dwivedi, V., Mandal, D., Sharma, N.K., Vishwakarma, N.K., Pal, L., Choudhary, M., Francis, A., Malakar, and Singh, N.** (2020). Root-specific expression of chickpea cytokinin oxidase/dehydrogenase 6 leads to enhanced root growth, drought tolerance and yield without compromising nodulation. *Plant Biotechnology Journal* 18(11), 2225-2240.
- Kieber, J.J. and Schaller, G.E.** (2018). Cytokinin signaling in plant development. *Development*, 145(4), dev149344.
- Köllmer, I., Novák, O., Strnad, M., Schmölling, T. and Werner, T.** (2014). Overexpression of the cytosolic cytokinin oxidase/dehydrogenase (*CKX 7*) from Arabidopsis causes specific changes in root growth and xylem differentiation. *The Plant Journal*, 78(3): 359-371.

- Kondo, M., Murty, M.V.R., Aragonés, D.V., Okada, K., Winn, T. and Kwak, K.S.** (1999). Characteristics of the root system and water uptake in upland rice. *Genetic Improvement of Rice for Water-Limited Environments*, 117-134.
- Kumara Charyulu, D. and Deb, U.**, (2014). Proceedings of the" 8th International Conference viability of small farmers in Asia" 1-154.
- Laplaze, L., Benkova, E., Casimiro, I., Maes, L., Vanneste, S., Swarup, R., Weijers, D., Calvo, V., Parizot, B., Herrera-Rodriguez, M.B. and Offringa, R.** (2007). Cytokinins act directly on lateral root founder cells to inhibit root initiation. *The Plant Cell*, 19(12): 3889-3900.
- Latham, M.C.** (1997). Human nutrition in the developing world (No.29). *Food & Agriculture Org.*
- Le, D.T., Nishiyama, R., Watanabe, Y., Vankova, R., Tanaka, M., Seki, M., Ham, L.H., Yamaguchi-Shinozaki, K., Shinozaki, K. and Tran, L.S.** (2012). Identification and expression analysis of cytokinin metabolic genes in soybean under normal and drought conditions in relation to cytokinin levels. *PloSOne*, 7(8).
- Letham, D.S.** (1963). Zeatin, a factor inducing cell division isolated from *Zea mays*. *Life Sciences* (1962), 8: 569.
- Li, M., Li, X., Zhou, Z., Wu, , Fang, M., Pan, X., Lin, Q., Luo, W., Wu, G. and Li, H.** (2016). Reassessment of the four yield-related genes Gn1a, DEP1, GS3, and IPA1 in rice using a CRISPR/Cas9 system. *Frontiers in Plant Science*, 7: 377.
- Li, S., Liu, X., Zhou, X., Li, Y., Yang, W. and Chen, R.** (2019). Improving zinc and iron accumulation in maize grains using the zinc and iron transporter ZmZIP5. *Plant and Cell Physiology*, 60(9): 2077-2085.
- Li, W., Fang, C., Krishnan, S., Chen, J., Yu, H., Murphy, A.S., Merewitz, E., Katin-Grazzini, L., McAvoy, R.J., Deng, Z. and Zale, J.** (2017). Elevated auxin and reduced cytokinin contents in rootstocks improve their performance and grafting success. *Plant Biotechnology Journal*, 15(12): 1556-1565.
- Li, W., Zhai, L., Strauss, S.H., Yer, H., Merewitz, E., Chen, J., Wang, X., Zhuang, W., Fang, C., Chen, Y. and McAvoy, R.** (2019). Transgenic reduction of cytokinin

levels in roots inhibits root-sprouting in *Populus*. *Plant Physiology*, 180(4): 1788-1792.

Li, X., Mo, X., Shou, H. and Wu (2006). Cytokinin-mediated cell cycling arrest of pericycle founder cells in lateral root initiation of *Arabidopsis*. *Plant and Cell Physiology*, 47(8): 1112-1123.

Li, Y., Fu, X., Zhao, M., Zhang, W., Li, B., An, D., Li, J., Zhang, A., Liu, R. and Liu, X. (2018). A genome-wide view of transcriptome dynamics during early spike development in bread wheat. *Scientific Reports*, 8(1): 1-16.

Liu, Zhang, C., Ma, J.Q., Zhang, L.Y., Yang, B., Tang, X.Y., Huang, L., Zhou, X.T., Lu, K. and Li, J.N. (2018). Genome-wide identification and expression profiling of cytokinin oxidase/dehydrogenase (*CKX*) genes reveal likely roles in pod development and stress responses in oilseed rape (*Brassica napus* L.). *Genes*, 9(3): 168.

Liu, Z., Lv, Y., Zhang, M., Liu, Y., Kong, L., Zou, M., Lu, G., Cao, J. and Yu, X. (2013). Identification, expression, and comparative genomic analysis of the IPT and *CKX* gene families in Chinese cabbage (*Brassica rapa* ss *pekinensis*). *BMC Genomics*, 14(1): 594.

Lohar, D., Schaff, J.E., Laskey, J.G., Kieber, J.J., Bilyeu, K.D. and Bird, D.M. (2004). Cytokinins play opposite roles in lateral root formation, and nematode and rhizobial symbioses. *The Plant Journal*, 38(2): 203-214.

Lu, J., Chang, C., Zhang, H., Wang, S.X., Sun, G., Xiao, S.H. and Ma, C.X. (2015). Identification of a novel allele of *TaCKX6a02* associated with grain size, filling rate and weight of common wheat. *PloSOne*: 10(12).

Mähönen, A., Bonke, M., Kauppinen, L., Riikonen, M., Benfey, N. and Helariutta, Y. (2000). A novel two-component hybrid molecule regulates vascular morphogenesis of the *Arabidopsis* root. *Genes & Development*, 14(23): 2938-2943.

Mameaux, S., Cockram, J., Thiel, T., Steuernagel, B., Stein, N., Taudien, S., Jack, Werner, Gray, J.C., Greenland, A.J. and Powell, W. (2012). Molecular, phylogenetic and comparative genomic analysis of the cytokinin oxidase/dehydrogenase gene family in the *Poaceae*. *Plant Biotechnology Journal*, 10(1): 67-82.

- Mason, M.G., Mathews, D.E., Argyros, D.A., Maxwell, B.B., Kieber, J.J., Alonso, J.M., Ecker, J.R. and Schaller, G.E.** (2005). Multiple type-B response regulators mediate cytokinin signal transduction in Arabidopsis. *The Plant Cell*, 17(11): 3007-3018.
- Massonneau, A., Houba-Hérin, N., Pethe, C., Madzak, C., Falque, M., Mercy, M., Kopečný, D., Majira, A., Rogowsky and Laloue, M.** (2004). Maize cytokinin oxidase genes: differential expression and cloning of two new cDNAs. *Journal of Experimental Botany*, 55(408): 2549-2557.
- Matsumoto-Kitano, M., Kusumoto, T., Tarkowski, Kinoshita-Tsujimura, K., Václavíková, K., Miyawaki, K. and Kakimoto, T.** (2008). Cytokinins are central regulators of cambial activity. *Proceedings of the National Academy of Sciences*, 105(50): 20027-20031.
- Miyawaki, K., Tarkowski, Matsumoto-Kitano, M., Kato, T., Sato, S., Tarkowska, D., Tabata, S., Sandberg, G. and Kakimoto, T.** (2006). Roles of Arabidopsis ATP/ADP isopentenyltransferases and tRNA isopentenyltransferases in cytokinin biosynthesis. *Proceedings of the National Academy of Sciences*, 103(44): 16598-16603.
- Morris, R.O., Bilyeu, K.D., Laskey, J.G. and Cheikh, N.N.** 1999. Isolation of a gene encoding a glycosylated cytokinin oxidase from maize. *Biochemical and Biophysical Research Communications*, 255(2), 328-333.
- Nehnevajova, E., Ramireddy, E., Stolz, A., Gerdemann-Knörck, M., Novák, O., Strnad, M. and Schmölling, T.** (2019). Root enhancement in cytokinin-deficient oilseed rape causes leaf mineral enrichment, increases the chlorophyll concentration under nutrient limitation and enhances the phytoremediation capacity. *BMC Plant Biology*, 19(1): 83.
- Nieminen, K., Immanen, J., Laxell, M., Kauppinen, L., Tarkowski, Dolezal, K., Tähtiharju, S., Elo, A., Decourteix, M., Ljung, K. and Bhalerao, R.** (2008). Cytokinin signaling regulates cambial development in poplar. *Proceedings of the National Academy of Sciences*, 105(50): 20032-20037.
- Nishiyama, R., Watanabe, Y., Fujita, Y., Le, D.T., Kojima, M., Werner, T., Vankova, R., Yamaguchi-Shinozaki, K., Shinozaki, K., Kakimoto, T. and Sakakibara, H.**

- (2011). Analysis of cytokinin mutants and regulation of cytokinin metabolic genes reveals important regulatory roles of cytokinins in drought, salt and abscisic acid responses, and abscisic acid biosynthesis. *The Plant Cell*, 23(6): 2169-2183.
- Ogonowska, H., Barchacka, K., Gasparis, S., Jablonski, B., Orczyk, W., Dmochowska-Boguta, M. and Nadolska-Orczyk, A.** (2019). Specificity of expression of *TaCKX* family genes in developing plants of wheat and their co-operation within and among organs. *PloSOne*, 14(4).
- Ramireddy, E., Galuszka, and Schmölling, T. (2018b).** Zn-fortified cereal grains in field-grown barley by enhanced root cytokinin breakdown. *Plant Signaling & Behavior*, 13(11): e1530023.
- Ramireddy, E., Hosseini, S.A., Eggert, K., Gillandt, S., Gnad, H., von Wirén, N. and Schmölling, T.** (2018a). Root engineering in barley: increasing cytokinin degradation produces a larger root system, mineral enrichment in the shoot and improved drought tolerance. *Plant Physiology*, 177(3): 1078-1095.
- Reid, D.E., Heckmann, A.B., Novák, O., Kelly, S. and Stougaard, J.** (2016). CYTOKININ OXIDASE/DEHYDROGENASE3 maintains cytokinin homeostasis during root and nodule development in *Lotus japonicus*. *Plant Physiology*, 170(2): 1060-1074.
- Riefler, M., Novak, O., Strnad, M. and Schmölling, T.** (2006). Arabidopsis cytokinin receptor mutants reveal functions in shoot growth, leaf senescence, seed size, germination, root development, and cytokinin metabolism. *The Plant Cell*, 18(1): 40-54.
- Rival, de Billy, F., Bono, J.J., Gough, C., Rosenberg, C. and Bensmihen, S.** (2012). Epidermal and cortical roles of NFP and DMI3 in coordinating early steps of nodulation in *Medicago truncatula*. *Development*, 139(18): 3383-3391.
- Sakakibara, H.** (2006). Cytokinins: activity, biosynthesis, and translocation. Annual Review. *Plant Biology*, 57: 431-449.
- Saxena, N., Krishnamurthy, L. and Johansen, C.** (1993). Registration of a drought-resistant chickpea germplasm. *Crop Science*, 33(6): 1424-1424.

- Skoog, F. and Miller, C.** (1957). Chemical regulation of growth and organ formation in plant tissues cultured. *In Vitro Symp Soc Exp Biol*.
- Steele, K.A., Price, A.H., Shashidhar, H.E. and Witcombe, J.R.** (2006). Marker-assisted selection to introgress rice QTLs controlling root traits into an Indian upland rice variety. *Theoretical and Applied Genetics*, 112(2): 208-221.
- Swaminathan, M.S. and Bhavani, R.V.** (2013). Food production & availability-Essential prerequisites for sustainable food security. *The Indian Journal of Medical Research*, 138(3): 383.
- To, J. and Kieber, J.J.** (2008). Cytokinin signaling: two-components and more. *Trends in Plant Science*, 13(2): 85-92.
- To, J., Haberer, G., Ferreira, F.J., Deruere, J., Mason, M.G., Schaller, G.E., Alonso, J.M., Ecker, J.R. and Kieber, J.J.** (2004). Type-A Arabidopsis response regulators are partially redundant negative regulators of cytokinin signaling. *The Plant Cell*, 16(3): 658-671.
- UNICEF.** 2015. United Nations Children's Fund.
- Von Schwartzberg, K., Nunez, M.F., Blaschke, H., Dobrev, I., Novák, O., Motyka, V. and Strnad, M.** (2007). Cytokinins in the bryophyte *Physcomitrella patens*: analyses of activity, distribution, and cytokinin oxidase/dehydrogenase overexpression reveal the role of extracellular cytokinins. *Plant Physiology*, 145(3): 786-800.
- Waidmann, S. and Kleine-Vehn, J.** (2020). Asymmetric cytokinin signalling opposes gravitropism in roots. *Journal of Integrative Plant Biology*.
- Wang, W., Simmonds, J., Pan, Q., Davidson, D., He, F., Battal, A., Akhunova, A., Trick, H.N., Uauy, C. and Akhunov, E.** (2018). Gene editing and mutagenesis reveal inter-cultivar differences and additivity in the contribution of TaGW2 homoeologues to grain size and weight in wheat. *Theoretical and Applied Genetics*, 131(11): 2463-2475.
- Wang, Y., Liu, H. and Xin, Q.** (2014). Genome-wide analysis and identification of cytokinin oxidase/dehydrogenase (CKX) gene family in foxtail millet (*Setaria italica*). *The Crop Journal*, 2(4): 244-254.

- Werner, T. and Schmülling, T.** (2009). Cytokinin action in plant development. *Current Opinion in Plant Biology*, 12(5), 527-538.
- Werner, T., Holst, K., Pörs, Y., Guivarc'h, A., Mustroph, A., Chriqui, D., Grimm, B. and Schmülling, T.** (2008). Cytokinin deficiency causes distinct changes of sink and source parameters in tobacco shoots and roots. *Journal of Experimental Botany*, 59(10): 2659-2672.
- Werner, T., Motyka, V., Laucou, V., Smets, R., Van Onckelen, H. and Schmülling, T.** (2003). Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. *The Plant Cell*, 15(11): 2532-2550.
- Werner, T., Motyka, V., Strnad, M. and Schmülling, T.** (2001). Regulation of plant growth by cytokinin. *Proceedings of the National Academy of Sciences*, 98(18): 10487-10492.
- Werner, T., Nehnevajova, E., Köllmer, I., Novák, O., Strnad, M., Krämer, U. and Schmülling, T.** (2010). Root-specific reduction of cytokinin causes enhanced root growth, drought tolerance, and leaf mineral enrichment in Arabidopsis and tobacco. *The Plant Cell*, 22(12): 3905-3920.
- Wightman, F., Schneider, E.A. & Thimann, K.V.** (1980). Hormonal factors controlling the initiation and development of lateral roots: II. Effects of exogenous growth factors on lateral root formation in pea roots. *Physiologia Plantarum*, 49(3): 304-314.
- Wood, J.A. and Grusak, M.A.** (2007). Nutritional value of chickpea. *Chickpea Breeding and Management*, 101-142.
- Worthen, J.M., Yamburenko, M.V., Lim, J., Nimchuk, Z.L., Kieber, J.J. and Schaller, G.E.** (2019). Type-B response regulators of rice play key roles in growth, development and cytokinin signaling. *Development*, 146(13).
- Yeh, S.Y., Chen, H.W., Ng, C.Y., Lin, C.Y., Tseng, T.H., Li, W.H. and Ku, M.S.** (2015). Down-regulation of cytokinin oxidase 2 expression increases tiller number and improves rice yield. *Rice*, 8(1): 36.

Zalewski, W., Gasparis, S., Boczkowska, M., Rajchel, I.K., Kała, M., Orczyk, W. and Nadolska-Orczyk, A. (2014). Expression patterns of *HvCKX* genes indicate their role in growth and reproductive development of barley. *PLoS One*, 9(12).

Zhang, L., Zhao, Y. L., Gao, L. F., Zhao, G.Y., Zhou, R. H., Zhang, B. S. and Jia, J. Z. (2012b). *TaCKX6D1*, the ortholog of rice *OsCKX2*, is associated with grain weight in hexaploid wheat. *New Phytologist*, 195(3): 574-584.

Zou, X., Shao, J., Wang, Q., Chen, Zhu, Y. and Yin, C. (2018). Supraoptimal cytokinin content inhibits rice seminal root growth by reducing root meristem size and cell length via increased ethylene content. *International Journal of Molecular Sciences*, 19(12): 4051.

Table 1: Some important crop improvement works using *CKX* manipulation.

Plant	<i>CKX</i> variant	Action	Effect	Source
Arabidopsis	<i>CKX1</i> , <i>CKX2</i> , <i>CKX3</i> , <i>CKX4</i>	Overexpression	Retarded shoot growth and enhanced root growth	Nishiyama <i>et al.</i> (2011)
			Tolerant to salt stress Tolerant to drought stress	
Rice	<i>CKX7</i>	Overexpression	Root growth and xylem differentiation	Kollmer <i>et al.</i> (2013)
	<i>CKX2</i> (Gn1a)	<i>CKX</i> mutant	High in panicle branches	Ashikari <i>et al.</i> (2005)
		RNAi	High in grains/panicle High in grain number	Yeh <i>et al.</i> (2015)
	Overexpression	High in grain number		
	RNAi	Low in grain number High in tiller number		
<i>CKX2</i> (Gn1a)	CRISPR/Cas9	High in grain number High in grain weight No response for grain/panicle	Li <i>et al.</i> (2016)	
<i>CKX9</i>	CRISPR/Cas9	High in flower number	Duan <i>et al.</i> (2019)	
	Overexpression	High in tiller number Low in panicle size		

Barley	<i>HvCKX1</i>	RNAi	Low in grain number High in spike number in T4 High in grain number in T4	Zalewski <i>et al.</i> (20014)
	<i>HvCKX9</i> <i>HvCKX1</i>	RNAi RNAi	No effect at T4 High in spike number	Holubova <i>et al.</i> (2018)
		CRISPR/Cas9	High in grain number Low in 1000 grain weight High in yield No yield data provided	
	<i>HvCKX1</i>	CRISPR/Cas9	Limited effect	Gasparis <i>et al.</i> (2019)
	<i>HvCKX3</i>	CRISPR/Cas9	Low in grain number Low in grain weight	
	<i>CKX2</i>	Targeted overexpression in root	Tolerant in long term drought	Ramireddy <i>et al.</i> (2018)
Wheat			Biofortification of Zn, Fe and other micronutrients	Ramireddy <i>et al.</i> (2018b)
	<i>Ta2.2.1-3A</i>	RNAi	High in grain weight	Li <i>et al.</i> (2018)
			No change in spikelet number	
	<i>GW2</i>	CRISPR/Cas9 + TILLING; triple mutant on A, B, D sub-genome	High in grain size High in 1000 grain weight	Wang <i>et al.</i> (2018)
	<i>CKX2.2.1-3D</i>	Mutation association analysis	High in 1000 grain weight	Zhang <i>et al.</i> (2012b)
	<i>CKX2.1-3D</i>	Mutation associated analysis	High in grain size	Lu <i>et al.</i> (2015)
		High in grain weight High in grain filling High in grain weight	Chang <i>et al.</i> (2015)	
Tobacco	<i>CKX1</i>	Root specific overexpression	Tolerant to drought stress Higher accumulation of minerals Improved leaf	Werner <i>et al.</i> (2010)

<i>Medicago sativa</i>	<i>MsCKX</i>	Overexpression	chlorophyll content Tolerant to salt stress	Li <i>et al.</i> (2019)
Lotus	<i>CKX3</i>	qRT analysis	Helping in nodule development	Reid <i>et al.</i> (2016)
<i>Brassica</i>	<i>CKX5-1, 5-2, 6-1, 7-1</i>	qRT analysis	Development of siliques	Liu <i>et al.</i> (2018)
	<i>CKX2</i>	Overexpression	Enhanced root growth	Nehnevajova <i>et al.</i> (2019)
<i>Populus</i>	<i>CKX2</i>	Root specific overexpression	Enhanced chlorophyll content Accumulation of higher amount of Cd and Zn	Li <i>et al.</i> (2019)
			Inhibition of sprouts development	

UNDER PEER REVIEW