

Original Research Article

Effect of Segregated and Unsegregated urban solid waste drum composts on growth, yield and nutrient uptake by finger millet (*Eleusine coracana L.*)

Abstract: A greenhouse experiment was conducted to study the effect of segregated and unsegregated urban solid waste drum compost on plant height, number of leaves, ear head weight, ear head length, 1000 grain weight, grain yield and straw yield in finger millet (*Eleusine coracana L.*) during kharif-2015 in Department of Soil Science and Agricultural Chemistry, University of Agricultural Sciences, Gandhi Krishi Vignana Kendra, Bangalore (India). The result revealed that plant height, number of leaves, ear head weight, ear head length, 1000 grain weight, grain yield and straw yield and uptake content were significantly improved by the application of 100% NPK+ segregated urban solid waste compost (10 t ha⁻¹) followed by 100% NPK+ unsegregated urban solid waste compost (10 t ha⁻¹) as compared to the treatment which include only inorganics.

Keywords: Effects, segregated, unsegregated urban solid waste compost, Growth, Yield, Nutrient uptake.

Comment [S1]: Why this? You don't need it in an Abstract. This one can be in the introduction.

Introduction

Finger millet (*Eleusine coracana L.*) Gaertn is an important cereal that belongs to the grass family Poaceae, subfamily Chloridoideae. It is estimated that finger millet accounts for some 10 per cent of the 30 million tons of millet produced globally (Dida *et al.*, 2008). It is also known as Ragi or African millet ranks fourth in importance among millets in the world after sorghum (*Sorghum bicolor*), pearl millet (*Pennisetum glaucum*) and foxtail millet (*Setaria italica*). It is grown globally on more than 4 million hectares and is the primary food source for millions of people in tropical dry land regions. It also has nutritional qualities superior to that of rice and is on par with wheat (Latha *et al.*, 2005). The crop is also adapted to a wide range of tropical soils, ranging from red lateritic to sandy loams and black heavy Vertisols (Dida and Dvos, 2006). In India, it is cultivated on 1.8 m ha with average yields of 1.3 t ha⁻¹. In Karnataka, finger millet is grown in an area of 0.76 m ha producing 1.32 million tonnes with a yield of 1715 kg ha⁻¹ (FAO, Stat, 2014).

Urban solid waste can be defined as any solid or semi-solid substance or object resulting from human or animal activities, discarded as useless or unwanted materials. It is an extremely mixed mass of wastes, which may originate from household, commercial, industrial or agricultural activities. Municipal solid waste (MSW) composting, however, has proved to be a safe and effective way to accelerate the decomposition and stabilization of biodegradable components of bio-waste from MSW, leading to production of compost for soil amendment or as an organic nutrient source.

With accelerating the world shift to an urban future, municipal solid waste (MSW) is a significant by-product of urban lifestyle that is expanding faster than urbanization. According to World Bank research, there has been approximately 70% global growth in urban MSW, which leads nations to most difficulties (Meena *et al.* (2016). As a result of the high rates of organic waste generation and their open dumpsite in landfills, there are certain negative impacts on the environment, economy, and social life. One of the most eco-friendly methods for keeping organic contaminants out of landfills is composting Erguven G. *et al.*, 2020. Recycling waste through composting is an environmentally favorable substitute, which can be used as a source of plant nutrients (Almendro Candel *et al.*, 2019). The main objectives of sustainable waste management are resource conservation, protection of the environment, and human health. Goals also include preventing the export of issues related to trash into the future and maintaining long-term soil fertility (Brust G.E, 2019).

Comment [S2]: This way of writing your reference is different from those in the first paragraph. Choose one way.

Comment [S3]: Put comma(,) here.

Municipal solid waste contains high soil organic matter, which becomes more significant as an organic amendment for restoring soil fertility and enhancing soil biological, physical, and chemical properties. Compost made from municipal solid waste reduces the harmful effects of salt-affected soils and functions as a soil conditioner significantly enhancing crop production (Meena MD *et al.* 2019).

Thus, contributing to an improvement of crop yield and quality, Urban solid waste compost consists about 51 percent of organic waste, utilizing it in agriculture not only improves soil fertility but also can save chemical fertilisers, if it is utilised scientifically through segregation and composting which results in mitigating environment pollution. Keeping these points in view, the present investigation was studied aiming to partially replace chemical fertiliser with segregated and unsegregated urban solid waste compost and study its effect on growth paramaters, grain yield, straw yield and uptake in finger millet (*Eleusine coracana L.*)

Materials and Methods

A pot culture experiment was conducted to study the effect of segregated and unsegregated urban solid waste drum compost on growth parameters, grain yield, straw yield and uptake in finger millet (*Eleusine coracana L.*) variety GPU-48. The experiment was conducted in Department of Soil Science and Agricultural Chemistry, Gandhi Krishi Vignana Kendra, Bangalore (India) during kharif-2015. The pot (8 kg soil pot⁻¹) study where the soil was treated with segregated, unsegregated urban compost and farm yard manure (FYM).

The soil used in this experiment was treated with segregated and unsegregated compost that were prepared with the use of urban solid waste containing organic waste such as vegetables, fruits, flower, dry leaves and saw dust etc. were separated and transferred into drum composter. Drum composter is a method of preparation of compost using biodegradable organic waste in an air circulated horizontal rotatory drum container. Using drum composter, the compost is prepared within 75-80 days and it is called segregated urban solid waste compost which is having less concentration of heavy metals. The unsegregated urban solid

wastes containing both organic and solid waste are transferred into the drum composter without separation. The compost was prepared aerobically as followed in the segregated urban solid waste compost preparation. The chemical composition of segregated, unsegregated urban solid waste compost and farm yard manure (FYM) is presented in (Table 1).

The experiment was laid out in a Complete Randomized Design (CRD) with three replications. The treatments were T₁ - 100% NPK + FYM at 10 t ha⁻¹ (Package of Practice); T₂ - Segregated compost (10 t ha⁻¹); T₃ - Unsegregated compost (10 t ha⁻¹); T₄ - FYM (10 t ha⁻¹); T₅- 100 % NPK + Segregated compost; T₆ - 100% NPK + Unsegregated compost; T₇ - T₂ + 50% NPK; T₈- T₃ + 50% NPK; T₉ -T₄ + 50% NPK; T₁₀ - 50% NPK; T₁₁ - 100% NPK.

Statistical Analysis

The data obtained from the study were subjected to statistical analysis of variance method at 5 per cent level of significance as per the procedure given by Sundaraj *et al.* (1972).

Table 1. Physical, chemical and biological composition of segregated, unsegregated urban solid waste compost and FYM used in the pot culture experiment

Parameters	FYM	Segregated compost	Unsegregated compost
Physical properties			
Moisture (%)	20.12	23.65	22.03
Colour	Brown	Black	Black
Bulk density (g cm ⁻³)	0.98	1.01	1.04
Chemical properties			
pH (1:10)	7.2	7.35	7.84
EC (dS m ⁻¹)	1.22	1.12	1.51
Organic carbon (%)	17.77	29.16	24.17
C: N ratio	29.01	17.78	19.18
N (%)	0.61	1.64	1.22
P (%)	0.18	0.45	0.32
K (%)	0.52	1.11	0.91
Ca (%)	0.68	1.47	0.95
Mg (%)	0.27	0.74	0.56
S (%)	0.21	0.76	0.64
Zn (mg kg ⁻¹)	13.86	118.07	143.17
Cu (mg kg ⁻¹)	2.2	42.11	44.23
Fe (mg kg ⁻¹)	520.3	3529.11	3604
Mn (mg kg ⁻¹)	38.12	350.67	366.33
Ni (mg kg ⁻¹)	18.21	12.75	23.42
Cd (mg kg ⁻¹)	ND	ND	ND
Pd (mg kg ⁻¹)	ND	23.87	43.62
Cr (mg kg ⁻¹)	ND	12.53	21.34

Result and discussion

The results revealed that plant height and number of leaves/plant was significantly influenced by various treatment combinations at 30, 60 and 90 DAS. Significantly higher plant height of 31.47, 76.90, 110.82 cm and 8.58, 10.50, 11.17 was recorded in T₅ (100 % NPK + Segregated drum compost) which was found on par with T₆ (100 % NPK+ Unsegregated drum compost) 29.92, 73.65, 103.70 cm and 8.33, 10.33, 10.70. The lower nutrient content was recorded in T₄ (FYM alone @ 10 t ha⁻¹) is presented in (Table 2). The increase in plant height with segregated drum compost followed by unsegregated drum compost may be attributed to increase in cell multiplication and elongation due to increased supply of 100 per cent NPK + Segregated drum compost @ 10 t ha⁻¹. Similar results were found in finger millet crop which reported by Kibria *et al.* (2013) and Kavitha and Subramanian (2007). Sultana M. *et al.* (2020) compost serve as a source of energy for microorganisms, boost the availability and concentration of nutrients in the soil and release nutrients in a manner easily absorbed by plants its increases growth and yield of plant.

Number of leaves increased due to residual effect of 100 per cent NPK + Segregated drum compost, which might be due to increased availability of nitrogen, phosphorus, potassium and other nutrients throughout the crop growth stages through as a result of continuous mineralization and its uptake. Similar results found in cowpea were reported by Roohi *et al.* (2018).

Table 2. Effect of quality of segregated and unsegregated urban solid waste drum composts on growth of finger millet in different stages.

Treatments	Plant height (cm)			No. of leaves /plant		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T ₁	27.77	73.27	97.92	8.33	10.17	10.57
T ₂	22.68	66.6	82.77	8.00	8.57	9.07
T ₃	23.67	63.87	85.48	8.17	9.2	9.64
T ₄	19.72	66.18	80.3	7.17	8.27	8.53
T ₅	31.47	76.9	110.82	8.58	10.5	11.17
T ₆	29.92	73.65	103.7	8.33	10.33	10.7
T ₇	27.08	63.92	96.45	7.83	10.03	10.23
T ₈	25.23	73.15	93.7	8.17	10.04	10.14
T ₉	25.15	71.82	87.55	7.83	9.37	10.07
T ₁₀	22.87	64.38	83.18	7.67	9.03	9.31
T ₁₁	23.15	69.88	87.13	8.00	9.5	9.91
S. Em ±	1.35	2.53	2.84	0.21	0.39	0.28
CD @ 5 %	3.95	7.42	8.34	0.61	1.15	0.83

The result indicated that grain and straw yield of finger millet was increased significantly to a maximum yield of 8.10 and 14.48 g/pot in T₅ (100 % NPK + Segregated drum compost) followed by T₆ (100 % NPK + Unsegregated drum compost @ 10 t ha⁻¹) 7.91 and 13.63 g/pot. The lower grain and straw yield recorded 6.33 and 10.82 g/pot in T₄ (FYM alone @ 10 t ha⁻¹) is presented in (Table 3).

Comment [S4]: In the body of your work, the reference must mention page number.

Significantly higher grain and straw yield of finger millet was recorded on combined application of 100 per cent NPK + Segregate drum compost. This may be ascribed to enough nutrients being provided to the crop and thus improving soil physical, microbial, chemical and nutritional properties which encourages proliferous root system resulting in better absorption of water and nutrient from soil and thus resulting in higher grain and straw yield and nutrient uptake. Similar findings were reported by Queriemmi *et al.* (2021) MSW compost with sewage sludge compost acts as a source of nutrients, it has a beneficial residual effect on nutrient contents in soil, and finally yields (increases up to 77%). The results are in conformity with Bhanu Prakash *et al.* (2007) and (Dimambro *et al.*, 2007). Improvement in yield component of wheat and maize with the application of urban compost and sewage sludge was reported by Sushil kumar (2002). Lima *et al.* (2004) who concluded that the urban waste application contributes to increase the growth of corn plants.

Earhead and 1000 grain weight of finger millet crop varied significantly, the treatment T₅ which received (100 % NPK + Segregated drum compost) recorded significantly higher earhead weight of 11.43 (g hill⁻¹), 3.58 g. Which was on par with T₆ (100 % NPK + Segregated drum compost) 11.24 (g hill⁻¹), 3.48 g. The lower grain and straw yield recorded 9.38 (g hill⁻¹) and 2.84 g in T₄ (FYM alone @ 10 t ha⁻¹). Significantly higher ear head length and number of finger per earhead of finger millet of 5.40 cm and 5.60 was recorded in T₅ (100 % NPK + Segregated drum compost) on par by T₆ (100 % NPK + Unsegregated drum compost) 5.28 cm and 5.43. The lower ear head length and number of finger per earhead 4.55 cm and 4.17 was recorded in T₄ (FYM alone @ 10 t ha⁻¹) is presented in (Table 3).

Application of segregated drum compost waste compost significantly influenced the earhead length, earhead weight, 1000 grain weight and number of finger per earhead of finger millet crop. Significantly higher earhead length and earhead weight were recorded in segregated drum compost treatment which could be ascribed to the slow and steady rate of nutrient release into the soil to match the adsorption pattern of finger millet. Sayara *et al.* (2020) reported that compost application can increase soil organic carbon three-fold and microbial activity double.

By adding organic matter to the soil, compost improves soil macronutrient levels that support plant metabolism and raises long-term soil productivity. Composts containing high amounts of available nitrogen (N) often result in more rapid plant growth and yield, whereas composts with more N bound up in the organic fraction exhibit surplus growth response over successive seasons. Roohi, *et al* (2018) reported urban solid waste compost its ability to make nutrients (macro and micro-nutrients) more readily available to crop plants upon mineralization and supply of the nutrient to the crop throughout the vegetation period of crop result in increase the plant growth and yield.

Dimambro *et al.* (2007) reported increased number of ears with the application of MSW compost and sewage sludge in wheat. Saunshi *et al.* (2014) recorded increased number

of earheads, earhead length and earhead weight in finger millet with application of enriched liquid bio-digester manure.

Table 3. Effect of quality of segregated and unsegregated urban solid waste drum composts yield and yield attributes of finger millet at harvest

Treatments	Grain yield (g / pot)	Straw yield (g / pot)	Earhead weight (g hill ⁻¹)	1000 grain weight (g)	Earhead length (cm)	No. of finger per earhead
T ₁	7.81	12.62	10.73	3.45	5.15	5.24
T ₂	6.84	11.34	9.87	2.88	4.68	4.48
T ₃	6.44	10.70	9.68	2.86	4.63	4.33
T ₄	6.33	10.82	9.38	2.84	4.55	4.17
T ₅	8.10	14.48	11.43	3.58	5.40	5.60
T ₆	7.91	13.63	11.24	3.48	5.28	5.43
T ₇	7.48	12.06	10.70	3.36	4.88	5.17
T ₈	7.45	11.90	10.67	3.25	4.80	5.11
T ₉	7.23	11.58	10.28	3.13	4.75	4.84
T ₁₀	6.58	11.17	10.01	2.94	4.67	4.50
T ₁₁	7.11	11.72	10.20	3.05	4.71	4.67
S. Em ±	0.04	0.12	0.17	0.14	0.16	0.26
CD (5%)	0.11	0.35	0.49	NS	0.47	0.75

Significantly higher uptake of nitrogen, phosphorus, potassium, calcium, magnesium and sulphur in finger millet grain of 0.18, 0.02, 0.11, 0.046, 0.026 and 0.045 g/pot was recorded in T₅ (100 % NPK + segregated drum compost) followed by treatment T₆ (100 % N + Unsegregated drum compost) which recorded 0.16, 0.017, 0.09, 0.036, 0.019 and 0.035 g/pot. The lower uptake in grain was recorded in T₄ (FYM alone @ 10 t ha⁻¹) of 0.05, 0.008, 0.05, 0.012, 0.007 and 0.013 g/pot is presented in (Table 4).

This might be attributed to the treatment which sufficiently supplied the nutrients as per crop requirement and release of N from organic amendments through mineralization, which in turn resulted in increased yield and thereby increased uptake. The results are in line with the work done by Protima *et al.* (2023) reported an nutrient content in soil and uptake in rice increase as compost application rates increased. The benefits to the soil of adding organic matter as well as N with compost can increase longer-term soil productivity. Sultana *et al.* 2021 reported that MSW compost inoculated with *Trichoderma* has a great way to improve the soil nutrient status (N, P, K, and S), soil fertility, and crop productivity which left a long impact on the soil. Similar findings were observed by Roohi, *et al* (2018) segregated urban solid waste compost is increases major nutrient uptake due to the complimentary effect of organics upon mineralization increased major and secondary nutrients availability in the soil results in uptake by cowpea. Oueriemmi H, *et al* (2021) reported that nutrient content in soil

and uptake in rice tended to increase as MSW compost application rates increased in the previous crop. The application of MSW compost and green manure does not replace chemical fertilization but is used in association with fertilizers can satisfy the necessity of crop nutrients.

Similar findings were observed by Grigatti *et al.* (2017) organic manure increases P availability and uptake by plants. Sunitha *et al.*, (2010) reported that availability of secondary nutrient status (Ca, Mg and S) concentration and uptake of nutrients were significantly influenced by native as well as the applied sources. The increase in sulphur uptake may be attributed to residual effect of applied organic which might have resulted in slow and steady release of sulphur from native as well as the applied sources.

Mohammadreza *et al.* (2010) reported an increase nitrogen content and uptake in soyabean with the application of municipal solid waste compost. Praveen *et al.* (2012) also reported increase in P uptake in maize with the application phosphate solubilizing fungi and different phosphorus levels. Badr *et al.* (2006) also found that potassium uptake by maize roots improved markedly with inoculation of bacteria in the tested soils compared to corresponding controls. The present results are in consonance with the findings of Rostami *et al.* (2012).

Table 4. Effect of segregated and unsegregated urban solid waste compost on of major and secondary nutrients uptake of finger millet grain

Treatments	Finger millet Grain					
	Nutrient uptake (g / pot)					
	N	P	K	Ca	Mg	S
T ₁	0.15	0.013	0.08	0.025	0.016	0.025
T ₂	0.07	0.009	0.06	0.018	0.008	0.015
T ₃	0.05	0.008	0.05	0.016	0.01	0.015
T ₄	0.05	0.008	0.05	0.012	0.007	0.013
T ₅	0.18	0.02	0.11	0.046	0.026	0.045
T ₆	0.16	0.017	0.09	0.036	0.019	0.035
T ₇	0.14	0.011	0.07	0.021	0.013	0.022
T ₈	0.13	0.01	0.06	0.021	0.012	0.021
T ₉	0.13	0.008	0.06	0.019	0.008	0.020
T ₁₀	0.09	0.009	0.05	0.018	0.012	0.018
T ₁₁	0.13	0.012	0.06	0.022	0.01	0.017
S. Em ±	0.01	0.001	0.008	0.005	0.002	0.004
CD (5%)	0.03	0.004	0.024	0.014	0.006	0.011

Conclusion

The Application of Segregated urban solid waste drum compost to finger millet crop significantly improved the quality of growth, yield and nutrient uptake were noticed as compared to inorganic fertilizer alone. Nutrient uptake in grain in finger millet increases as compost application rates increased. The benefits to the soil of adding organic matter as well as N with compost can increase longer-term soil productivity. Composts with high levels of available N tend to show more immediate plant response in terms of growth, yield and uptake while compost with more N tied up in the organic fraction shows a carry-over growth response in subsequent seasons.

References

AKTAR SA, ISLAM MS, HOSSAIN MS, AKTER H, MAULA S, HOSSAIN SS. Effects of municipal solid waste compost and fertilizers on the biomass production and yield of BRR1 Dhan 50. *Progressive Agriculture*. 2018;29 (2):82-90.

ALMENDRO-CANDEL MB, NAVARRO-PEDREÑO J, LUCAS IG, ZORPAS AA, VOUKKALI I, LOIZIA P. The use of composted municipal solid waste under the concept of circular economy and as a source of plant nutrients and pollutants. *Municipal Solid Waste Management*. 2019;33-50

Comment [S5]: Read well MLA to master the way to put the reference.

BADR, M. A., SHAFEI, M. A. AND SHARAF, S. M., 2006, The dissolution of K and P-bearing minerals by silicate dissolving bacteria and their effect on sorghum growth. *Res. J. Agric. Biol. Sci.*, 2(1): 5-11.

BHANU PRAKASH, U. H., BHARGAVI, M. V., RAMAKRISHNA PARAMA, V. R. AND PREETHU, D.C., 2007, Bioremedial recycling of solid urban waste. *Proc. Int. Conf. Sust. Solid Waste Mngt.*, pp.289-295.

Comment [S6]: Here is APA's method. Follow only one method (MLA or APA) in an article.

LATHA, M. A, VENKATESWARA RAO, K. AND DASHAVANTHA REDDY, V., 2005, Production of transgenic plants resistant to leaf blast disease in finger millet (*Eleusine coracana* (L.) Gaertn.). *Pl. Sci.*, 169: 657-667.

DIDA, M. M. AND DVOS, K. M., 2006, Genome mapping and molecular breeding in plants; in cereals and millets (ed.) C Kole (Berlin: Springer-Verlag). 8: 333-343.

- DIMAMBRO, M. E., LILLYWHITE, R. D. AND RAHN, C. R., 2007, The physical, chemical and microbial characteristics of biodegradable municipal waste derived composts. *Compost Sci. Util.*, 15(4): 243-252.
- DOMÍNGUEZ M, PARADELONÚÑEZ R, PIÑEIRO J, BARRAL MT. Physicochemical and biochemical properties of an acid soil under potato culture amended with municipal solid waste compost. *International Journal of Recycling of Organic Waste in Agriculture*. 2019;8:171-178.
- GRIGATTI M, BOANINI E, MANCARELLA S, SIMONI A, CENTEMERO M, VEEKEN AH. Phosphorous extractability and ryegrass availability from bio-waste composts in a calcareous soil. *Chemosphere*. 2017;174: 722-31
- ERGUVEN G, AND KANAT G. Importance of solid waste management on composting, problems, and proposed solutions: The case of Turkey. *Avrupa Bilimve Teknoloji Dergisi*. 2020;19:66-71.
- FAO statistics division 2014, 04 August 2014.
- KAVITHA, R. AND SUBRAMANIAN, P., 2007a, Effect of enriched municipal solid waste compost application on growth, plant nutrient uptake and yield of rice. *J. Agron.*, 6(4): 586-592.
- KIBRIA, D. P. AND HAM, R.K., 2013, The effect of lignin and sugars to the aerobic composting of solid waste. *Waste management*, 23(5): 419-423.
- LIMA, J. S., DE QUEIROZ, J. E. G. AND FREITAS, H. B., 2004, Effect of selected and non-selected urban waste compost on the initial growth of corn. *Resources, Cons. Recycling.*, 42: 309-315.
- MALLIKARJUN, K., DEVAKUMAR, N., CHALAPATHI, N .V. AND GANGADAR ESHWAR, R. G., 2000, Integrated P management for sunflower (*Helianthus annuus*. L) in Alfisols. *Crop Res.*, 19(1): 23-27.
- MEENA MD, JOSHI PK, JAT HS, CHINCHMALATPURE AR, NARJARY B, SHEORAN P, SHARMA DK. Changes in biological and chemical properties of saline soil amended with municipal solid waste compost and chemical fertilizers in a mustard-pearl millet cropping system. *Catena*. 2016; 140:1-8.
- MOHAMMADREZA, M., ZOHRE, A., SEVDA, A. AND KOBRA, K., 2010, Fertilization of soybean plants with municipal solid waste compost under leaching and non-leaching conditions. *American-Eurasian J. Agric. Envi. Sci.*, 8(1): 55-59.

- PRAVEEN, P. M., KULIGOD, V. B., HEBSUR, N. S., PATIL, C. R. AND KULKARNI, G. N., 2012, Effect of phosphate solubilizing fungi and phosphorus levels on growth, yield and nutrient content in maize (*Zea mays*). *Karnataka J. Agric. Sci.*, 25(1): 58-62.
- PROTIMA, R.S., MD. MAZIBUR RAHMAN, ABDULLAH AL MAMUN, NAZMUN NAHAR AND MOHAMMAD M.H.T., 2013, Application of Municipal Solid Waste Compost and Green Manure Exerted Residual Effects on Soil Nutrient Content and Plant Nutrient Uptake in Rice. *International Journal of Plant & Soil Science.*,35(8):158-170
- ROSTAMI, S. V., PIRADASHTI, H., BAHMANYAR, M. A. AND ALALEH MOTAGHIAN, 2012, Response of soyabean (*Glycine max L.*) yield and nutrient uptake in three consecutive years application of municipal solid waste compost. *Int. J. Agri. Crop. Sci.*, 4(8): 468-473.
- ROOHI, H.C. PRAKASHA AND H.M. MEENA. Residual Effect of Segregated and Unsegregated Urban Solid Waste Compost on Yield and Nutrient Uptake by Cowpea (*Vigna unguiculata L.*) *Int.J.Curr.Microbiol.App.Sci.* (2018);7(8): 2028-2336.
- ROOHI, H.C. PRAKASHA AND H.M. MEENA, 2018, Residual Effect of Segregated and Unsegregated Urban Solid Waste Compost on Yield and Nutrient Uptake by Cowpea (*Vigna unguiculata L.*) *Int.J.Curr.Microbiol.App.Sci.*,7(8): 2028-2336.
- SAUNSHI, S., REDDY, V. C., MALLIKARJUN. AND RAJESH RAWAL., 2014, Influence of enriched bio-digester liquid manure on growth and yield of finger millet. *The Bioscan.*, 9(2): 613-616.
- SAYARA, T. SALIMIA, R. B., HAWAMDE, F. AND SANCHEZ, A. 2020, Recycling of organic wastes through composting: Process Performance and Compost Application in Agriculture., *Agronomy.*,10(11):1838.
- SESHADRI REDDY, S., SHIVARAJ, B. AND REDDY, V.C., 2004, Nutrient uptake and agronomic efficiency of groundnut as influenced by different organic manures. *Karnataka J. Agric. Sci.*, 17(4): 670-675.
- SUNDARAJ, N., NAGARAJU, S., VENKATARAMU, M, N. AND JAGANNATH, M. L., 1972, Design and analysis of field experiments. *Univ. Agric. Sci. Bangalore.*
- SUSHIL KUMAR, NAVINDU GUPTA, AKOLKAR, A. B., JOSHI, H. C. AND JAIN, M.C., 2002, Studies on quality and application of MSW compost and sewage sludge in Agriculture. 2: 2nd Int. Agron. Cong., pp. 1052-1053.

- SULTANA M, JAHIRUDDIN M, ISLAM MR, RAHMAN MM, ABEDIN MA, SOLAIMAN ZM. Nutrient-enriched municipal solid waste compost increases yield, nutrient content, and balance in rice. *Sustainability*. 2021; 13(3):1047.
- SULTANA M, JAHIRUDDIN M, ISLAM MR, RAHMAN MM, ABEDIN M. Effects of nutrient-enriched municipal solid waste compost on yield and nutrient content of cabbage in alluvial soil. *Asian Journal of Soil Science and Plant Nutrition*. 2020; 6(4):32-42.
- SUNITHA, B.P., PRAKASHA, H.C AND GURUMURTHY, K.T. 2010. Influence of organics, inorganics and their combination on availability, content and uptake of secondary nutrients by rice crop (*oryza sativa* L.) in bhadra command, Karnataka. *Mysore Journal of Agricultural Sciences*, 44(3): 509-516
- OUERIEMMI H, KIDD PS, TRASAR-CEPEDA C, RODRÍGUEZ-GARRIDO B, ZOGLAMI RI, ARDHAOUI K, PRIETO-FERNÁNDEZ Á, MOUSSA M. Evaluation of composted organic wastes and farmyard manure for improving the fertility of poor sandy soils in arid regions. *Agriculture*. 2021;11(5):415.