

Assessment of Physico-chemical properties of soil from different villages of Sunder Nagar block of Mandi district of Himachal Pradesh, India

Abstract

One of the most popular phrases in agriculture is the term “Soil Health and Soil Quality.” Agricultural sustainability is being dependent on soil health. Soil quality assessment is of paramount importance to know the appropriate management practices to be adopted for sustainable crop production. Soil samples were collected from Sunder Nagar block of Mandi district on Oct. 2022 in three depths viz. 0-15,15-30 and 30-45 cm and analysed for their Physico-chemical parameters by using standard laboratory techniques. Soil health parameters were analyzed and the founding state that, texture of soil is sandy loam. Soil reaction was neutral to slightly alkaline with soil EC mostly below $<1.0 \text{ dS m}^{-1}$ significantly affect for better crop production. In the case of organic carbon, nitrogen and phosphorus content of research area was found to be slightly low to medium while the range of potassium range was medium to high. The concentration levels of certain micronutrients were analyzed and the results indicate that zinc levels were low to slightly medium, copper levels were low to medium, iron levels were medium to high and manganese levels were medium to slightly high. For promoting the soil health and soil quality use of organic manure is recommended for fertilizer management.

Keywords: Soil health, Sunder Nagar block, Physico-chemical properties, Texture, etc.

Introduction

Himachal Pradesh is in the western Himalayas. Covering an area of 55,673 square kilometers (21,495 sq m), it is a mountainous state. Most of the state lies on the foothills of the Dhauladhar Range. At 6,816 m, Reo Purgyl is the highest mountain peak in the state of Himachal Pradesh.

Sunder Nagar, Mandi District, Himachal Pradesh – $17^{\circ}5'018''$, India ($11^{\circ}0'7.506''$)

76°5'6.552'' E) Co-ordinates: 31°23'2.63'' N 76°6'06.05'' E to 32°0'7.604'' N 77°3'7.726'' E Maximum elevation: 6.405 m, Minimum Elevation:272 m, Average elevation :1.945 m.


Soil is a dynamic, 3-dimensional natural body of the landscape developed from the weathering of rocks through various pedogenic processes, composed of mineral and organic materials, possessing a defined set of physical, chemical and biological qualities, having variable depth covering the earth's surface, and providing a medium for terrestrial plant growth. The rate of soil deterioration is influenced by land use patterns, soil types, terrain and climate variables. Inappropriate land use is one of these variables that accelerates the deterioration of soil physico-chemical and biological quality. (**Bindu *et al.*, 2022**).

The physical and chemical characteristics of soil play a big role in the plants ability to extract water and nutrients. High quality soils not only produce better food and fiber, but also help to establish natural ecosystem and enhance air and water quality. The physical properties of soil depend upon the shape, structure, size, pore space, amount of organic matter and mineral composition of soil. The chemical properties of the soil are the interactions of various chemical constituents among soil particles and soil solution. The physical and chemical properties are soil texture, bulk density, water holding capacity, soil structure, soil colour, pH, electrical conductivity, cation exchange capacity, organic carbon and soil nutrients (macro and micro) (**Griffiths *et al.*, 2010**).

Agroclimatic zones of Himachal Pradesh

Himachal is in the western Himalayas. Covering an area of 55,673 square kilometres (21,495 sqm), it is a mountainous state. Most of the state lies on the foothills of the Dhauladhar Range. At 6,816 m, Reo Purgyl is the highest mountain peak in the state of Himachal Pradesh. Sundar Nagar, Mandi District, Himachal Pradesh – 175018, India (11.07506 76.56552) Co-ordinates: 31.23263 76.60605 32.07604 77.37726, Maximum elevation: 6.405 m, Minimum Elevation:272 m, Average elevation :1.945 m.

Materials and Methods



Central Survey of India

Map1: Study area

Table 1: Sampling Sites

S. No.	Village	Latitude (°N)	Longitude (°E)
1	JUGAHAN	11°3'41.947"	76°36'12.401"
2		11°3'27.392"	76°35'39.627"
3		11°0'3.791"	76°35'31.246"
4	JARAL	11°5'39.462"	76°37'16.399"
5		11°5'27.892"	76°37'8.677"
6		11°5'50.95"	76°37'19.459"
7	DHANOTU	11°9'20.75"	76°39'50.745"
8		11°9'9.605"	76°39'20.937"
9		11°9'4.147"	76°38'55.266"

Study Area

Soil samples were collected from three different villages *i.e.*, Jugahan, Jaral, Dhanotu block of Sunder Nagar in Mandi district. From each village of soil samples were taken from three different areas. Soil samples were collected from the highland, middle land and lowland fields that are used for crop production.

Soil Sampling

Soil samples were collected from three different depths *i.e.*, 0-15, 15-30 and 30-45 cm. Most root activity and fertilizer applications are generally restricted to 30-45 cm depth. Total number of soil samples were collected 27. First the surface area of the sampling spot was cleared out. Weeds, leaves, stones were removed. A 'V' shaped pit of 15 cm depth was dug out with the help of garden *hoe/spade*, depth was measured by a meter scale and 1 to 2 cm slice of soils were collected using garden *hoe/ spade/khurpi*. The soil was put in a clean white paper and was evenly spread out. Foreign materials like roots, stones, pebbles and gravels were removed. Then the soils were mixed thoroughly and quartered into four equal parts. Two opposite soil quarters were discarded and the remaining two were mixed up.

Each soil sample was spread on a clean white paper sheet in the shade and air dried at room temperature. The large lumps/clods were crushed/broken to its ultimate soil particle using a wooden mallet. The powdered soil was sieved through 2 mm sieve. The soil material was collected in a clean cloth or polythene bag and was labelled properly for laboratory analysis.

Table 2: Analysis of Physico-chemical Properties of Soil

Sr. No.	Particulars	Scientist name	Methods	Unit
I.	Physical properties			
1.	Bulk density	Muthuvel <i>et al.</i> , 1992	Measuring cylinder	Mg m ⁻³
2.	Particle density			Mg m ⁻³
3.	Pore space			%
4.	Water holding capacity			%
5.	Colour	Munsell, 1971	Munsell colour charts	Colour
6.	Texture	Bouyoucos, 1927	Bouyoucos hydrometer	Texture

	(Sand, Silt, Clay %)			(Sand, Silt, Clay %)
II	Chemical properties			
1.	Soil pH (1:2.5)	Jackson, 1958	pH meter	
2.	Electrical conductivity (1:2.5)	Wilcox, 1950	digital conductivity meter	dS m ⁻¹
3.	Organic carbon	Walkley and Black, 1947	Walkley and Black Wet oxidation method	kg ha ⁻¹
4.	Available nitrogen	Subbiah and Asija, 1956	Modified alkaline permanganate oxidation method	kg ha ⁻¹
5.	Available phosphorus	Olsen <i>et al.</i> , 1954.	Olsen's extraction followed by Spectrophotometric method	kg ha ⁻¹
6.	Available potassium	Toth and Prince, 1949	Neutral normal ammonium acetate extraction followed by Flame photometric method	kg ha ⁻¹

Result and Discussion

Bulk density

The highest bulk density was recorded in Dhanotu (V₃F₂) is 1.38, 1.40 and 1.43 (Mgm⁻³) and lowest value of bulk density recorded in Jugahan (V₁F₁) 1.16, 1.19 and Dhanotu (V₃F₃) 1.22 (Mg m⁻³) at depth 0-15, 15-30 and 30-45 cm. Similar results were earlier reported by Kumari *et al.*, 2013 and Pravin *et al.*, 2013.

Particle density

The highest particle density was found in Dhanotu (V₃F₁) 2.49, 2.496 and 2.498 (Mg m⁻³) and lowest value of particle density found in Jaral (V₁F₁) 2.261, 2.265 and 2.267 (Mg m⁻³) at depth 0-15, 15-30 and 30-45 cm. Similar results were reported by Kumari *et al.*, 2013 and Pravin *et al.*, 2013.

Pore space

The highest pore space was recorded in Dhanotu (V₃F₁) was 49.2, 48.9 and 47.5 % and lowest value of pore space recorded in Dhanotu (V₃F₂) is 41.7, 41.2 and (V₃F₃) 40.2 % at depth 0-15, 15-30 and 30-45 cm. These results were similar with the findings of Ratnam *et al.*, 2001.

Water holding capacity

The highest water holding capacity (%) was recorded in Dhanotu (V_3F_1) 47.2, 46.9 and 45.5 % and lowest value of water holding capacity (%) found in Dhanotu (V_3F_2) 39.7, 39.2 and 38.2 % at depth 0-15, 15-30 and 30-45 cm. Similar trends were observed by Venkateswarlu *et al.*, 1995 and Pulakeshi *et al.*, 2014.

pH

The highest pH values was recorded in Jugahan (V_1F_2) 7.63, 7.66 and 7.68 and lowest value of pH Dhanotu (V_3F_2) 6.94, 7.02 and 7.24 at depth 0-15, 15-30 and 30-45 cm. similar trend was observed by Kekane 2015; Patel 2015 and Kumari *et al.*, 2005

EC

The highest values was recorded in EC Dhanotu (V_3F_3) 0.325, Jugahan (V_1F_1) 0.217 and (V_1F_3) 0.490 ($dS\ m^{-1}$) and lowest value of EC Dhanout (V_3F_1) 159, (V_3F_3) 0.140 and 0.308 ($dS\ m^{-1}$) at depth 0-15, 15-30 and 30-45 cm. These results were in accordance with the findings of Krishnamurthy *et al.*, 2007.

Organic carbon

The highest value of organic carbon percent was found in soil of Jaral (V_2F_2) 0.381, 0.377 and 0.372 % and lowest value of OC Jaral (V_2F_3) 0.314, 0.308 and 0.305 % at depth 0-15, 15-30 and 30-45 cm. These results were similar with the findings of Ratnam *et al.* (2001).

Available nitrogen

The highest value of available nitrogen was found in soil of Jaral (V_2F_2) 246.39, 241.11 and Jughan (V_1F_3) 234.38 $kg\ ha^{-1}$ and lowest value in soil of Jaral (V_2F_3) 216.18, 209.38 and 200.37 $kg\ ha^{-1}$ with depth 0-15, 15-30 and 30-45 cm. Similar observations of reported by Bandyopadhyay *et al.*, 2004.

Available phosphorus

The highest value of available phosphorus was found in soil of Dhanotu (V_3F_2) 19.26, 17.26 and 14.88 $kg\ ha^{-1}$ and lowest value in soil of Jugahan (V_1F_2) 10.88, 8.23 and 7.26 $kg\ ha^{-1}$ with depth 0-15, 15-30 and 30-45 cm. Similar observations of high phosphorus content were reported by Bandyopadhyay *et al.*, 2004.

Available potassium

The maximum value of available potassium was found in soil of Dhanotu (V_3F_2) 157.77, 153.48 and 149.38 $kg\ ha^{-1}$ and minimum value in soil of Jaral (V_2F_3) 125.63, 118.48 and 130.53 $kg\ ha^{-1}$ with depth 0-15, 15-30 and 30-45 cm. Similar observations of high potassium content were reported by Bandyopadhyay *et al.*, 2004.

Available iron

The highest value of Iron is found in soil of Dhanotu (V_3F_1) 22.48, Jugahan (V_1F_3) 21.38 and (V_1F_3) 21.30 ppm and lowest value in soil of Jaral (V_2F_1) 17.27, 15.58 and 13.69 ppm with depth 0-15, 15-30 and 30-45 cm. The pH range is generally between 7.0 to 8.5, which is considered slightly alkaline. Similarly result reported by Shukla *et al.*, 2015.

Available manganese

The highest value of Manganese is found in soil of Dhanotu (V_3F_1) 18.91, (V_3F_2) 16.80 and 15.23 ppm and lowest value in soil of Jaral (V_2F_1) 8.55, 7.39 and 6.01 ppm with depth 0-15, 15-30 and 30-45 cm respectively. Similarly result reported by Shukla *et al.*, 2015.

Available zinc

The highest value of Zinc is found in soil of Jugahan (V_1F_2) 3.17, (V_1F_3) 3.18 and Dhanotu (V_3F_3) 2.55 ppm and lowest value in soil of Dhanotu (V_3F_2) 1.58, 1.52 and 1.49 ppm with depth 0-15, 15-30 and 30-45 cm. Similarly result reported by Shukla *et al.*, 2015.

Available copper

The highest value of copper was found in soil of Jugahan (V_1F_2) 3.56, (V_1F_3) 3.23 and 2.75 ppm and minimum value in soil of Dhanotu (V_3F_2) 1.53, 1.51 and 1.02 ppm with depth 0-15, 15-30 and 30-45 cm. The value of copper (ppm) is found invaried from 1.63-7.54. Similarly result reported by Shukla *et al.*, 2015.

Table 3: Bulk density (Mg m^{-3}), particle density (Mg m^{-3}), pore Space (%) and water holding capacity (%) of soil at different depth.

Name of village and Farmer's field		Bulk density (Mg m^{-3})			Particle density (Mg m^{-3})			Pore Space (%)			Water holding capacity (%)		
		0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
JUGAHAN	V₁F₁	1.16	1.19	1.3	2.273	2.275	2.279	47.10	46.10	45.20	45.1	44.1	43.1
	V₁F₂	1.24	1.26	1.29	2.284	2.288	2.289	48.80	47.10	45.20	46.8	45.1	43.2
	V₁F₃	1.17	1.2	1.23	2.272	2.274	2.276	44.77	43.80	42.20	42.7	41.8	40.2
JARAL	V₂F₁	1.3	1.33	1.35	2.381	2.385	2.389	46.60	47.80	43.50	44.6	42.8	41.5
	V₂F₂	1.37	1.39	1.4	2.363	2.365	2.389	45.60	44.40	43.90	44.6	42.4	41.9
	V₂F₃	1.22	1.25	1.27	2.261	2.265	2.267	47.40	46.10	45.20	45.4	44.1	43.2
DHANOTU	V₃F₁	1.24	1.27	1.3	2.49	2.496	2.498	49.90	48.90	47.50	47.2	46.9	45.5
	V₃F₂	1.38	1.4	1.43	2.452	2.456	2.458	41.70	41.20	40.90	39.7	39.2	38.2
	V₃F₃	1.17	1.2	1.22	2.471	2.473	2.477	42.70	41.70	40.20	40.7	39.7	38.9
F- test		S	S	S	S	S	S	S	S	S	S	S	S
S.Em. (\pm)		0.01	0.02	0.01	0.03	0.03	0.03	0.64	0.43	0.64	0.59	0.38	0.55
C. D. (P=0.05)		0.04	0.06	0.05	0.10	0.11	0.11	1.92	1.28	1.90	1.75	1.14	1.65

Table 4: pH (w/v), EC (dSm⁻¹) and organic carbon (%) of soil at different depth.

Name of village and Farmer's field		pH(w/v)			EC (dS m ⁻¹)			Organic carbon (%)		
		0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
JUGAHAN	V₁F₁	7.34	7.45	7.64	0.25	0.26	0.29	0.367	0.361	0.354
	V₁F₂	7.39	7.40	7.68	0.40	0.43	0.45	0.323	0.317	0.311
	V₁F₃	7.35	7.37	7.38	0.28	0.30	0.31	0.348	0.341	0.337
JARAL	V₂F₁	7.31	7.43	7.58	0.24	0.26	0.29	0.328	0.321	0.316
	V₂F₂	7.42	7.43	7.45	0.29	0.31	0.33	0.381	0.377	0.372
	V₂F₃	7.47	7.49	7.53	0.39	0.38	0.36	0.314	0.308	0.305
DHANOTU	V₃F₁	7.32	7.41	7.28	0.23	0.25	0.26	0.359	0.352	0.348
	V₃F₂	7.28	7.46	7.24	0.20	0.22	0.24	0.339	0.333	0.329
	V₃F₃	7.34	7.45	7.51	0.25	0.26	0.30	0.345	0.340	0.337
F- test		NS	NS	NS	NS	NS	NS	S	S	S
S.Em. (±)		-	-	-	-	-	-	0.005	0.004	0.004
C. D. (P=0.05)		-	-	-	-	-	-	0.017	0.010	0.012

Table 5: Available nitrogen (kg ha⁻¹), available phosphorus (kg ha⁻¹) and available potassium (kg ha⁻¹) of soil at different depth.

Name of village and Farmer's field		Available nitrogen (kg ha ⁻¹)			Available phosphorus (kg ha ⁻¹)			Available potassium (kg ha ⁻¹)		
		0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
JUGAHAN	V₁F₁	246.36	235.38	231.24	16.91	13.32	10.23	151.08	144.98	142.63
	V₁F₂	220.48	213.36	209.44	10.88	8.23	7.26	132.31	127.56	120.66
	V₁F₃	242.41	238.96	234.38	14.32	12.98	11.22	135.00	130.09	126.21
JARAL	V₂F₁	239.88	234.46	227.31	16.28	12.32	9.68	150.11	145.38	141.27
	V₂F₂	246.39	241.11	233.97	15.32	13.66	11.78	142.22	137.42	132.98
	V₂F₃	216.18	209.23	200.37	14.31	12.76	10.84	125.63	118.48	130.53
DHANOTU	V₃F₁	243.73	237.89	231.23	18.26	16.34	12.98	157.77	153.48	149.38
	V₃F₂	240.01	236.38	232.98	19.26	17.26	14.88	156.63	152.36	148.39
	V₃F₃	241.71	237.86	233.48	17.87	15.58	12.26	152.67	148.23	144.49
F- test		S	S	S	S	S	S	S	S	S
S.Em. (±)		3.13	3.38	3.72	0.26	0.20	0.19	2.47	1.96	1.42
C. D. (P=0.05)		9.30	10.06	11.06	0.79	0.60	0.56	7.36	5.83	4.21

Table 6: Available iron (ppm), available manganese (ppm), available zinc (ppm) and available copper (ppm) of soil at different depth.

Name of village and Farmer's field		Available iron (ppm)			Available manganese (ppm)			Available zinc (ppm)			Available copper (ppm)		
		0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
JUGAHAN	V₁F₁	21.62	20.60	19.10	14.25	11.63	10.75	2.46	2.44	2.34	2.7	2.59	2.32
	V₁F₂	21.38	20.84	19.88	11.26	10.49	9.73	3.71	2.53	2.32	3.56	2.88	2.62
	V₁F₃	21.3	21.38	21.30	9.03	8.71	7.28	3.43	3.18	2.16	3.41	3.23	2.75
JARAL	V₂F₁	17.27	15.58	13.69	8.55	7.39	6.01	2.44	1.7	1.55	2.3	2.2	1.79
	V₂F₂	20.84	19.15	17.00	10.67	9.31	8.54	2.71	2.49	2.23	2.7	2.32	2.32
	V₂F₃	17.87	16.72	13.84	9.24	9.17	7.02	2.5	2.35	2.01	2.48	2.17	2.03
DHANOTU	V₃F₁	22.48	19.01	15.12	18.91	15.43	10.92	2.4	2.32	1.79	1.85	1.63	1.29
	V₃F₂	19.79	19.79	19.56	17.58	16.8	15.23	1.58	1.52	1.49	1.53	1.51	1.02
	V₃F₃	19.01	18.64	16.95	17.9	12.510	12.3	3.05	2.72	2.55	3.05	2.72	2.55
F- test		S	S	S	S	S	S	S	S	S	S	S	S
S.Em. (±)		0.27	0.33	0.29	0.20	0.17	0.13	0.03	0.04	0.02	0.02	0.02	0.04
C. D. (P=0.05)		0.81	0.98	0.87	0.60	0.52	0.40	0.10	0.12	0.08	0.08	0.06	0.11




Fig. 1: Bulk density (Mg m^{-3}), and particle density (Mg m^{-3}) of soil at different depth.




Fig. 2: Pore space (%) and water holding Capacity of soil at different depth.




Fig. 3: Organic carbon (%) of soil at different depth.




Fig. 4: Available nitrogen (kg ha⁻¹), and available potassium (kg ha⁻¹) of soil at different depth.




Fig. 5: Available phosphorus (kg ha⁻¹) of soil at different depth.




Fig. 6: Available iron (ppm), and available manganese (ppm) of soil at different depth.




Fig. 7: Available zinc (ppm) and available copper (ppm) of soil at different depth.

Conclusion

It is concluded that the soils samples were moderately to strongly alkaline in reaction and non-saline. The physical properties of both surficial and sub-surficial soils are normal as the bulk density value is optimum. The Water Holding Capacity is medium. The overall fertility status of the soils was low, medium and high in nitrogen, phosphorus and potassium respectively. As the soils were calcareous and strongly alkaline, there is need for application of any acid forming amendment and organic materials to alleviate the nutrient deficiency and improve productivity.

Acknowledgements

The authors are grateful to the Hon'ble Vice chancellor SHUATS, Department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute, for taking their keen interest and encouragement to carry out the research work.

Reference

- Bindu, A. H., Hasan, A., Thomas, T., David, A. A. and Singh, A. K. (2022) Assessment of Physico-Chemical Properties of Soil from Different Blocks of Prakasam District, Andhra Pradesh, India. *International Journal of Plant and Soil Science*, **34**(19): 340-349.
- Bouyoucos, G.L, (1927) The hydrometer as a new method for the mechanical analysis of soils. *Soil Sci.* **23**: 343-353.
- Fisher, R. A. and Yates (1960) *Statistical method for research worker Oliver and Boyd Ltd.* Edin. burgh and London.10.
- Jackson, M. L. (1958) Soil chemical analysis Prentice Hall of India Ltd. New Delhi. 219-221.
- Kekane, S. S., Chavan, R. P., Shinde, D. N., Patil, C. L. and Sagar, S. S. (2015) A review on physico-chemical properties of soil. *International Journal of Chemical Studies*, **3**(4), 29-32.
- Lindsay, W. L. and Norvell, W. A. (1978) Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Science Society of America Journal*; **42**: 421-428.
- Munsell, A. H. (1971) Munsell's description of his colour system, from a lecture to the American Psychological Association. *American Journal of Psychology* **23**(2): 236-244.

- Muthuvel, P., Udayasoorian, C., Natesan, R. and Ramaswamy, P. P. (1992) Introduction to Soil Analysis, *Tamil Nadu Agricultural University Coimbatore-641002*.
- Olsen, S. R., Cole, C. V., Watanabe, F. S. and Dean, L. A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate (NaHCO₃), *U.S.D.A. Circular. 93(9)*: 1-19.
- Patel, K. P. (2015) Analysis of Soil Quality Using Physico-Chemical Parameters of Shehra Taluka District: Panchmahals (Gujarat). *Indian journal of applied sciences*, **5(9)**, 466-468.
- Subbiah, B. V. and Asijja, E. C. (1956) A rapid procedure for estimation of available nitrogen in soil. *Current Science* **25(8)**: 259-260.
- Tewari, G., Khati, D., Rana, L., Yadav, P., Pande, C., Bhatt, S. and Joshi, P. K. (2016) Assessment of Physicochemical Properties of Soils from Different Land Use Systems in Uttarakhand, India. *Journal of Chemical Engineering and Chemistry Research*, **3(11)**: 1114-1118.
- Toth, S. J. and Prince, A. L. (1949) Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by flame photometer technique. *Soil Sci.*, **6(7)**: 439-445.
- Walkley, A. and Black, I. A., (1947) Estimation of soil organic carbon by the chromic acid titration method. *Soil Science*. **4(7)**: 29-38.
- Wilcox, L.V. (1950) Electrical conductivity. *Am. Water Works Assoc. J.* **4(2)**: 775-776.