

# Review Article

## Iron toxicity in plants: A Review

---

### ABSTRACT

Iron (Fe) is an essential micronutrient for plant growth and development, but excessive iron uptake can cause iron toxicity, leading to damage to plant cell membranes, reduced growth, yield, and overall health. This review paper discusses the issue of iron toxicity in plants, a common problem that affects crops such as rice, soybean, wheat, vegetables and is a common issue in Southeast Asia, Brazil, Africa, Australia, and the United States. Iron toxicity is more likely to occur in soils with high pH, high organic matter, or elevated levels of available iron. It discusses the various mechanisms that cause iron toxicity in plants, such as competition with other essential elements, oxidative stress, and changes in gene expression and phytohormones. The excess iron ions can cause damage to the root cells and the plasma membrane, leading to oxidative stress and increased production of reactive oxygen species (ROS). ROS can cause damage to cellular components, such as lipids, proteins, and DNA, leading to the death of root cells. Plants have specific mechanisms to take up iron, including transport proteins that are responsible for moving  $Fe^{2+}$  across the plasma membrane of root cells. Furthermore, it discusses the impact of iron toxicity on plant growth and development, including stunted growth, reduced root development, decreased water and nutrient uptake, and reduced photosynthetic activity. Highlights the importance of proper management of iron levels in soils to prevent iron toxicity and promote healthy plant growth.

*Keywords:* Iron toxicity, Cell membrane, Yield, Reducing growth, Oxidative stress

### 1. INTRODUCTION

Iron (Fe) is an essential micronutrient for plant growth and development, playing a critical role in various physiological processes, such as photosynthesis, respiration, and nitrogen metabolism [1]. Plants need to maintain iron (Fe) in the concentration of  $10^{-4}$  to  $10^{-9}$  M to achieve optimal growth [2]. However, excessive iron uptake can lead to iron toxicity, causing damage to the plant's cell membranes, reducing growth, yield, and affecting the overall health of the plant [3]. Iron toxicity can occur in different soils, including those with high pH, high levels of organic matter, or elevated levels of available iron [4]. These conditions lead to an increase in the concentration of free Fe ions, which can bind to cellular components, disrupting cellular function and leading to plant stress. It is a common problem in many parts of the world, affecting crops such as rice, soybean, wheat, maize, and vegetables. Some of the regions with high iron toxicity in plants include Southeast Asia, Brazil, Africa, Australia, and United States. Iron toxicity can occur in both soil-grown and hydroponic systems, and it is especially common in calcareous soils, which have high pH levels [1].

Iron toxicity can occur in different plants if the levels of iron in the soil are too high [3]. Some plants are more sensitive to iron toxicity than others, such as most fruit trees, tomatoes, and peppers. On the other hand, some plants are less sensitive to iron toxicity and can tolerate high levels of iron in the soil, such as most grasses, some tree species like oak, pine, and eucalyptus [1]. High levels of iron in the soil can also lead to the formation of

toxic compounds, such as hydroxyl radicals, that can cause oxidative damage to plant cells. The plant's antioxidant system, including enzymes such as superoxide dismutase, catalase, and ascorbate peroxidase, helps to reduce the levels of ROS in the plant cells and protect against oxidative stress [3]. Additionally, excess Fe accumulation can disrupt the nutrient balance of plants, leading to deficiencies in other essential nutrients such as zinc, copper, and manganese. Nutrient deficiencies can further increase the effects of Fe toxicity, leading to reduced plant growth, chlorosis, and necrosis. It can have a significant impact on crop productivity and quality [5]. In addition to reducing plant growth, it can also lead to a decline in yield and quality of harvested crops. The presence of excess Fe in the soil can also have negative effects on soil microbial communities, further impacting plant growth and productivity.

## 2. Impact on plant growth and development

Iron toxicity in plants is believed to be associated with the regulation of iron uptake and transport. Iron uptake in plants is facilitated by specific transporters that are regulated by the concentration of iron in the soil [6]. It is a serious problem for plants as it can negatively affect various aspects of their growth and development. It can cause stunted growth and reduced root development, which leads to decreased water and nutrient uptake and reduced photosynthetic activity [7]. In addition, iron toxicity can result in decreased chlorophyll content, which affects the plant's ability to perform photosynthesis, leading to reduced growth and productivity. In severe cases, leaf necrosis may occur, leading to the death of the plant [8]. Leaf yellowing is another common symptom of iron toxicity, which is associated with chlorosis [1]. Furthermore, iron toxicity can cause reduced seed production, which affects plant reproduction and survival [5].

## 3. Mechanisms of iron toxicity in plants

Iron toxicity in plants occurs when the concentration of iron in the soil solution exceeds the plant's ability to absorb and transport it to the roots. The form of iron that is taken up by plants is usually  $\text{Fe}^{2+}$  (ferrous iron) ion.  $\text{Fe}^{3+}$  (ferric iron) is the most common form of iron in aerobic soils, but it is not easily absorbed by plant roots because it forms insoluble precipitates [9]. In anaerobic soils,  $\text{Fe}^{2+}$  is more prevalent as it is formed by reduction of  $\text{Fe}^{3+}$  by microorganisms, and this form of iron can be more easily absorbed by plant roots. Plants have specific mechanisms to take up iron, including transport proteins that are responsible for moving  $\text{Fe}^{2+}$  across the plasma membrane of root cells. Plants also secrete compounds called "siderophores," which bind to  $\text{Fe}^{3+}$  and help to solubilize it, making it more available for uptake by plant roots [10]. Once inside the plant, iron is used in several processes, including the formation of chlorophyll for photosynthesis and in the production of enzymes that are involved in respiration and nitrogen fixation.

The excess iron ions can cause damage to the root cells and the plasma membrane, leading to oxidative stress and increased production of reactive oxygen species (ROS). ROS can cause damage to cellular components, such as lipids, proteins, and DNA, leading to the death of root cells [11]. Excess iron can also affect the uptake of other essential elements, such as phosphorus and zinc, by competing with the transporters that facilitate their uptake by the roots. This interference can lead to nutrient imbalances, reduced growth, and overall plant health [1].

Recent research studies have shown that excess iron can lead to alterations in the expression of genes involved in iron uptake, transport, and homeostasis [12]. These changes in gene expression can further increase the physiological and biochemical changes

associated with iron toxicity. Another mechanism involved in iron toxicity in plants is the production of phytohormones, such as ethylene and auxin. Excess iron can stimulate the production of ethylene and auxin in plants, leading to various physiological and biochemical changes [13]. These changes can further contribute to the toxic effects of excess iron on plant growth and development.

#### **4. Effects of iron toxicity on plant**

Iron toxicity can have detrimental effects on plant growth and development, [14] found that elevated iron concentration caused a boost in the antioxidant enzyme activity in both roots and leaves, but lowered the levels of photosynthetic pigments, leaf gas exchange, and overall plant growth. Similarly, [15] reported that iron toxicity can cause poor growth, tillering, reduction in crop spike number, spikelet sterility, and flowering delay or even failure and severe yield reductions, associated with leaf discoloration. It can reduce crop yields dramatically, such as that of rice, by up to 78% in West Africa.

The presence of excess iron in submerged paddy fields with low pH, often leads to iron toxicity in rice, which can disrupt cell homeostasis and impair growth and yield due to the significant increase in ferrous ion concentration as reported by [16]. Iron toxicity symptoms in rice manifest due to the excessive absorption of  $Fe^{2+}$  by the roots, which is then transported upwards to the leaves through acropetal translocation. This results in an increased production of harmful oxygen radicals, which can harm cell structural components and disrupt physiological functions [17]. Excess iron can trigger a rise in polyphenol oxidase activity, resulting in the synthesis of oxidized polyphenols. Additionally, iron toxicity can induce leaf bronzing and decrease root oxidation capacity. Excessive levels of  $Fe^{2+}$  ions increase the generation of ROS ( $O_2^{\cdot-}$ ,  $H_2O_2$ ,  $^1O_2$ ,  $HO_2$ ,  $OH^{\cdot}$ ,  $OH^-$ , and RO), which are extremely toxic and reactive and can cause protein oxidation, lipid peroxidation, carbohydrate, and DNA damage, which can result in cell death [18]. Fe toxicity in the medicinal plant *Bacopa monnieri* L. leads to the production of MDA in the leaves and shoots [19].

Iron toxicity can have adverse effects on the uptake and balance of other vital nutrients in plants, as reported by [20]. Their study revealed that iron toxicity reduced the uptake of nitrogen (N), phosphorus (P), and potassium (K). The reduced uptake of these essential nutrients can cause nutrient imbalances, leading to growth inhibition and reduced yield.

#### **5. Strategies for mitigating iron toxicity**

Iron toxicity in plants can be a significant problem in areas with high soil iron levels. To mitigate its effects, several strategies have been developed. One approach is to modify the soil environment by adjusting the soil pH can reduce the availability of iron to the plant. For example, adding lime to alkaline soils or sulphur to acidic soils can help to lower the soil pH and decrease the uptake of iron [8]. Another strategy is to select iron-tolerant plant species. certain plant species are more tolerant to iron toxicity and can be used in areas with high soil iron levels. Additionally, chelation can be used to reduce the toxicity of excess iron. Chelation is the process of binding metal ions with chelating agents, such as organic molecules called chelators or chelating agents [34]. The resulting complex formed between the chelator and the metal ion is called a chelate. In biological systems, chelation is involved in the transport and storage of metal ions, such as iron, copper, and zinc, and is critical for the proper functioning of enzymes and other proteins that require these metals as cofactors. Chelation can also be used to remove excess metal ions [35].

**Table 1: Effects of iron toxicity on different crops**

Crop	Iron Concentration in Plant	Effect	Reference
Wheat	>100 mg kg <sup>-1</sup>	Inhibition of root, shoot growth, reduced yield, and oxidative stress	[21]
Rice	>300 mg kg <sup>-1</sup>	Leaf bronzing, stunted growth, reduction in yield, and biomass	[22]
Cherry	>1300 mg kg <sup>-1</sup>	Reduced relative growth rate (RGR) of roots and leaves, leaf chlorosis and darkening of roots	[23]
Soybean	>30 mg kg <sup>-1</sup>	Leaf chlorosis, stunted growth, reduced yield, reduced biomass, oxidative stress, and change in gene expression	[24]
Arabidopsis	>350 μM Fe L <sup>-1</sup>	Reduced growth, chlorosis, and oxidative stress	[25]
Pepper tree	>250 μM Fe L <sup>-1</sup>	Leaf chlorosis, dark roots, and plant growth reduction	[26]
Green gram	>200 μM Fe L <sup>-1</sup>	Seed germination is significantly reduced	[27]
Cucumber	>10 μM Fe L <sup>-1</sup>	Decrease in chlorophyll content, growth, and yield	[28]
Common reed	>50 mg L <sup>-1</sup>	Reduced growth of the seedlings	[29]
Cowpea	>400 mg L <sup>-1</sup>	Retarded growth, decreased chlorophyll concentration, reduced yield and oxidative stress.	[30]
Pea	>40 mg L <sup>-1</sup>	Reduced growth, chlorosis, and yield reduction	[31]
Potato	>0.1 Mm Fe	Reduced growth, decreased chlorophyll concentration, reduced tuber yield, deterioration of quality by lowering the concentration of sugars, starch.	[32]
Ashwagandha	>200 mM Fe	Thickened cell wall, folding and shrinkage of cell wall, damage of cell membranes	[33]

According to [1], chelating agents like EDTA or DTPA can be added to the soil to solubilize excess iron and reduce its toxicity. Limiting waterlogging can also help to mitigate the effects of iron toxicity. As reported by [36], waterlogging can exacerbate iron toxicity by reducing oxygen levels in the soil and increasing the concentration of iron ions in the soil

solution. Finally, using iron-efficient cultivars can be an effective strategy for reducing the effects of iron toxicity. [37] investigated the effectiveness of using chelating agents, such as citric acid and ethylenediamine tetra acetic acid (EDTA), to reduce the bioavailability of iron in the soil and reduce the symptoms of iron toxicity in rice plants. Other studies have focused on genetic engineering to increase the plant's tolerance to iron toxicity, such as by overexpressing genes involved in iron uptake and transport [38].

The use of iron-deficient soils or hydroponic solutions has also been shown to reduce the symptoms of iron toxicity. Additionally, the application of iron-solubilizing bacteria, such as *Bacillus* spp., has been shown to increase the bio-availability of iron in the soil and reduce the symptoms of iron toxicity in various crops [39]. These approaches have shown promise in mitigating iron toxicity in plants and could be used to improve crop yield and reduce the negative effects of iron toxicity on plant growth and development.

#### 4. CONCLUSION

Iron toxicity is a widespread problem that can seriously affect plant growth and development in soils with high iron concentrations. This condition leads to reduced productivity and increased susceptibility to diseases. Several studies have highlighted the damaging effects of iron toxicity on photosynthesis, chlorophyll content, antioxidant defence systems, and nutrient uptake and balance in plants. To alleviate the negative effects of iron toxicity, several strategies have been suggested. One of the most effective ways is to modify soil conditions by adjusting the pH and/or adding organic matter, such as compost or manure, to the soil. Another strategy is to use iron-tolerant plant species, which have developed mechanisms to withstand high levels of iron in their environment. Other methods include chelation, limiting waterlogging, and using iron-efficient cultivars. Despite these strategies, further research is necessary to gain a better understanding of the mechanisms of iron toxicity in plants and to develop new and more effective approaches for mitigating its negative effects. Research could help identify the genetic and physiological factors that influence iron tolerance in plants and develop better soil management practices for iron-contaminated soils.

#### REFERENCES

- 1 Meharg A. Marschner's Mineral Nutrition of Higher Plants. Edited by P. Marschner. Amsterdam, Netherlands: Elsevier/Academic Press (2011), pp. 684, US \$124.95. ISBN 978-0-12-3849052. *Experimental agriculture*. 2012 Apr;48(2):305-.
2. Kim SA, Guerinot ML. Mining iron: iron uptake and transport in plants. *FEBS letters*. 2007 May 25;581(12):2273-80.
3. Kumar V, Sinha AK, Makkar HP, Becker K. Dietary roles of phytate and phytase in human nutrition: A review. *Food chemistry*. 2010 Jun 15;120(4):945-59.
4. Alloway BJ, editor. *Heavy metals in soils: trace metals and metalloids in soils and their bioavailability*. Springer Science & Business Media; 2012 Jul 18..
5. Kumari S, Kumar R, Chouhan S, Chaudhary PL. Influence of Various Organic Amendments on Growth and Yield Attributes of Mung Bean (*Vigna radiata* L.). *International Journal of Plant & Soil Science*. 2023 May 13;35(12):124-30. <https://doi.org/10.9734/ijpss/2023/v35i122975>
6. Morrissey J, Guerinot ML. Iron uptake and transport in plants: the good, the bad, and the ionome. *Chemical reviews*. 2009 Oct 14;109(10):4553-67.

7. Peña-Olmos JE, Casierra-Posada F, Olmos-Cubides MA. The effect of high iron doses (Fe<sup>2+</sup>) on the growth of broccoli plants (*Brassica oleracea* var. *Italica*). *Agronomía Colombiana*. 2014 Apr;32(1):22-8.
8. Zahra N, Hafeez MB, Shaukat K, Wahid A, Hasanuzzaman M. Fe toxicity in plants: Impacts and remediation. *Physiologia Plantarum*. 2021 Sep;173(1):201-22.
9. Krohling CA, Eutrópico FJ, Bertolazi AA, Dobbss LB, Campostrini E, Dias T, Ramos AC. Ecophysiology of iron homeostasis in plants. *Soil Science and Plant Nutrition*. 2016 Jan 2;62(1):39-47.
10. Kermeur N, Pédrot M, Cabello-Hurtado F. Iron Availability and Homeostasis in Plants: A Review of Responses, Adaptive Mechanisms, and Signaling. *Plant Abiotic Stress Signaling*. 2023 Mar 22:49-81.
11. Onyango DA, Entila F, Dida MM, Ismail AM, Drame KN. Mechanistic understanding of iron toxicity tolerance in contrasting rice varieties from Africa: 1. Morpho-physiological and biochemical responses. *Functional Plant Biology*. 2018 Oct 29;46(1):93-105.
12. Thomine S, Vert G. Iron transport in plants: better be safe than sorry. *Current opinion in plant biology*. 2013 Jun 1;16(3):322-7.
13. Zaid A, Ahmad B, Jaleel H, Wani SH, Hasanuzzaman M. A critical review on iron toxicity and tolerance in plants: role of exogenous phytoprotectants. *Plant micronutrients: Deficiency and toxicity management*. 2020:83-99.
14. Delias DS, Da-Silva CJ, Martins AC, de Oliveira DS, do Amarante L. Iron toxicity increases oxidative stress and impairs mineral accumulation and leaf gas exchange in soybean plants during hypoxia. *Environmental Science and Pollution Research*. 2021 Nov 17:1-2.
15. Audebert A, Sahrawat KL. Mechanisms for iron toxicity tolerance in lowland rice. *Journal of Plant Nutrition*. 2000 Nov 1;23(11-12):1877-85.
16. Aung MS, Masuda H. How does rice defend against excess iron?: Physiological and molecular mechanisms. *Frontiers in Plant Science*. 2020 Aug 7;11:1102.
17. Becker M, Asch F. Iron toxicity in rice—conditions and management concepts. *Journal of Plant Nutrition and Soil Science*. 2005 Aug;168(4):558-73.
18. Boruah K, Bharali A. Physiological basis of iron toxicity and its management in crops. *Recent Advances in Crop Physiology*. 2015 Jan;2:203-24.
19. Sinha S, Saxena R. Effect of iron on lipid peroxidation, and enzymatic and non-enzymatic antioxidants and bacoside-A content in medicinal plant *Bacopa monnieri* L. *Chemosphere*. 2006 Mar 1;62(8):1340-50.
20. Sahrawat KÁ. Iron toxicity in wetland rice and the role of other nutrients. *Journal of plant nutrition*. 2005 Jan 2;27(8):1471-504..
21. Setter TL, Waters I, Sharma SK, Singh KN, Kulshreshtha N, Yaduvanshi NP, Ram PC, Singh BN, Rane J, McDonald G, Khabaz-Saberi H. Review of wheat improvement for waterlogging tolerance in Australia and India: the importance of anaerobiosis and element toxicities associated with different soils. *Annals of Botany*. 2009 Jan 1;103(2):221-35.
22. Khairullah I, Saleh M, Alwi M. Increasing productivity of rice through iron toxicity control in acid sulfate soils of tidal swampland. *InIOP Conference Series: Earth and Environmental Science 2021 Feb 1 (Vol. 648, No. 1, p. 012151)*. IOP Publishing.
23. de Oliveira Jucoski G, Cambraia J, Ribeiro C, de Oliveira JA, de Paula SO, Oliva MA. Impact of iron toxicity on oxidative metabolism in young *Eugenia uniflora* L. plants. *Acta physiologicae plantarum*. 2013 May;35:1645-57.
24. Gülser F, Yavuz Hİ, Gökkaya TH, Sedef M. Effects of iron sources and doses on plant growth criteria in soybean seedlings. *Eurasian Journal of Soil Science*. 2019;8(4):298-303.
25. Li B, Sun L, Huang J, Göschl C, Shi W, Chory J, Busch W. GSNOR provides plant tolerance to iron toxicity via preventing iron-dependent nitrosative and oxidative cytotoxicity. *Nature communications*. 2019 Aug 29;10(1):3896.

26. Dobbss LB, Dos Santos TC, Pittarello M, de Souza SB, Ramos AC, Busato JG. Alleviation of iron toxicity in *Schinus terebinthifolius* Raddi (Anacardiaceae) by humic substances. *Environmental Science and Pollution Research*. 2018 Apr;25:9416-25.
27. Verma L, Pandey N. The effect of iron toxicity on seed germination and early seedling growth of green gram (*Vigna radiata* L. Wilczek). *Int J Sci Res*. 2017;6(8):1427-30.
28. Waters BM, Troupe GC. Natural variation in iron use efficiency and mineral remobilization in cucumber (*Cucumis sativus*). *Plant and Soil*. 2012 Mar;352:185-97.
29. Batty LC, Younger PL. Effects of external iron concentration upon seedling growth and uptake of Fe and phosphate by the common reed, *Phragmites australis* (Cav.) Trin ex. Steudel. *Annals of Botany*. 2003 Dec 1;92(6):801-6..
30. IFIE JE, IFIE-ETUMAH SO, IKHAJIAGBE B. Physiological and biochemical responses of selected cowpea (*Vigna unguiculata* (L.) Walp.) accessions to iron toxicity. *Acta Agriculturae Slovenica*. 2020 Feb 18;115(1):25-38.
31. Nenova V. Effect of iron supply on growth and photosystem II efficiency of pea plants. *Gen Appl Plant Physiol*. 2006 Jan;32:81-90.
32. Chatterjee C, Gopal R, Dube BK. Impact of iron stress on biomass, yield, metabolism and quality of potato (*Solanum tuberosum* L.). *Scientia horticulturae*. 2006 Mar 16;108(1):1-6.
33. Zhang Y, Wang Q, Xu C, Sun H, Wang J, Li L. Iron (Fe<sup>2+</sup>)-induced toxicity produces morphological and physiological changes in roots in *Panax ginseng* grown in hydroponics. *Toxicological & Environmental Chemistry*. 2016 Jul 2;98(5-6):630-7.
34. Dey S, Regon P, Kar S, Panda SK. Chelators of iron and their role in plant's iron management. *Physiology and Molecular Biology of Plants*. 2020 Aug;26(8):1541-9.
35. Kim JJ, Kim YS, Kumar V. Heavy metal toxicity: An update of chelating therapeutic strategies. *Journal of Trace elements in Medicine and Biology*. 2019 Jul 1;54:226-31.
36. Khabaz-Saberi H, Setter TL, Waters I. Waterlogging induces high to toxic concentrations of iron, aluminum, and manganese in wheat varieties on acidic soil. *Journal of Plant Nutrition*. 2006 Jun 1;29(5):899-911.
37. Chouhan S, Kumari S, Kumar R, Chaudhary PL. Climate Resilient Water Management for Sustainable Agriculture. *International Journal of Environment and Climate Change*. 2023 May 10;13(7):411-26. <https://doi.org/10.9734/ijecc/2023/v13i71894>
38. Koźmińska A, Wiszniewska A, Hanus-Fajerska E, Muszyńska E. Recent strategies of increasing metal tolerance and phytoremediation potential using genetic transformation of plants. *Plant biotechnology reports*. 2018 Feb;12:1-4.
39. Athukorala AS. Solubilization of micronutrients using indigenous microorganisms. *Microbial Technology for Sustainable Environment*. 2021:365-417.