

Original Research Article

Genetic variability, heritability and genetic advance studies in field pea (*Pisum sativum* L.) for yield and its attributing traits

Formatted: Font: Italic

ABSTRACT

An investigation was carried out with 48 genotypes of pea obtained through line x tester mating analysis and were tested using a randomized block experimental design with three replications at Tirhut College of Agriculture, Dholi, Dr. Rajendra Prasad Central Agricultural University, Bihar. The result showed highly significant difference for all the 14 fourteen characters under study. The variability, heritability and genetic advance as percent mean were studied for all the characters. High genotypic coefficient of variation was observed for the traits number of secondary branches, plant height, number of pods per plant, harvest index, nodules fresh weight and nodules dry weight. High heritability coupled with high genetic advance estimates was recorded for number of secondary branches, plant height, pod length, number of seeds per pod, number of pods per plant, 100 seed weight, grain yield per plant, harvest index, nodules fresh weight and nodules dry weight indicating the role of additive gene action and selection for these traits could be is reliable.

Comment [SA1]: Indicate the purpose of the study first then about the material and method

Keywords: genetic variability, genetic advance, heritability, Peas, Pisum traits, yield

Formatted: Font: Italic

INTRODUCTION

Pulses play a crucial role, especially in developing nations like India where the majority of the population is vegetarian. Pulses belong to the family *Leguminosae* and subfamily *Papilionoideae*. For a vegetarian diet, pulses are an excellent substitute of protein as they are nutritive and economical. They are also rich sources of amino acids especially lysine. Because of the ease of their simple digestion of the seeds, 22-24% protein content, and increased demand for amino-rich base substances for livestock feed or ancillary goods for human nutrition, these pulses are becoming important in day-to-day life (1).

Peas (*Pisum sativum* L.) ($2n=14$), is one of the ~~that is among the~~ temperate pulse crops, is a leguminous plant with in Papilionoideae subfamily and a member of the dicotyledons class. ~~$2n=14$ is the chromosome number found in the pea. It is~~ extensively cultivated in cooler climates and is native to South Western Asia ~~and can endure~~. ~~Due to the fact that it is primarily a cold weather crop, it can~~ endure light frost. Field peas (*Pisum sativum* L. var. *arvense*), also referred to as dry peas, and garden peas (*Pisum sativum* L. var. *hortens*), commonly known as table peas, are the two types of peas which are traditionally grown.

Genetic diversity in pea is ~~still~~ crucial for crop breeding programmes to be successful, since the diversity of the crop influences the extent to which it may be improved through selection. Additionally, the relative magnitudes of the phenotypic and genotypic variation coefficients provide a solid indication of the degree of population diversity (2). ~~Estimating the characteristics of variability, particularly heritability and genetic advance, are crucial indications for improving characters through selection. Though selection for highly heritable characters is more efficient for a successful breeding strategy.~~ Considering this, an effort was made in the present research to evaluate the degree of variability, heritability, and genetic advance for various pea characteristics.

MATERIALS AND METHODS:

~~Seven lines (females) namely, VL 42, RFPG 111, TRCP 8, HUPT 1810, KPF 14-29, DDR-23, IPFD 19-9 and five testers (males) namely, IPF-18-7, PANT P 436, PANT P 462, APARNA and HFP 1545 of field pea, chosen due to their phenotypic variability, were hybridized during in Line x tester fashion during rabi. 2021-22. The current study was conducted~~ at Tirhut College of Agriculture, Dholi, Dr. RPCAU, Bihar. ~~The resultant 35 hybrids were evaluated along with parents and check variety RACHANA, the research was carried out. The experiment was performed during the rabi of 2021-2022 to generate crosses and 2022-2023, ----- design with ----- replications. Each entry was raised in a plot size of ----- sq.m. to evaluate the parents and crosses.~~ Dholi is 55 meters above MSL and is geographically located at 25.9° N latitude and 85.9° E longitude. The soil condition is sandy loam with a modest level of organic matter and an alluvial origin. The weather during the growing season was ideal for normal growth and development of field pea crop.

~~Seven lines (females) namely, VL 42, RFPG 111, TRCP 8, HUPT 1810, KPF 14 29, DDR 23, IPFD 19 9 and five testers (males) namely, IPF 18 7, PANT P 436, PANT P 462, APARNA and HFP 1545 of field pea were chosen due to their phenotypic variability. In~~

Comment [SA2]: Redraft

Comment [SA3]: ???

Formatted: Font color: Text 1

~~rabi2020-2021, 35 crosses were produced utilizing seven lines and five testers. Therefore, the 35 crosses and their twelve progenitors, including the popular check variety RACHANA, were utilized as the experimental materials.~~

Estimation of heritability

Heritability in broad sense was estimated following the formula as suggested by Johnson *et al.* (1955):

$$\text{Heritability (h}^2\text{) (\%)} = \frac{\sigma^2_g}{\sigma^2_p} \times 100$$

Heritability in narrow sense was calculated according to the formula

$$\text{Heritability (ns) (\%)} = \frac{2\sigma^2_{gca}}{2\sigma^2_{gca} + \sigma^2_{sca} + \sigma^2_e} \times 100$$

Where,

σ^2_g = genotypic variance

σ^2_p = phenotypic variance

σ^2_e = environment variance

As suggested by (3) heritability (h^2) estimates were categorized as:

Low: 0 - 30%; Medium: 30 – 60 %; High: Above 60%

As suggested by Robinson (1966) for narrow sense heritability were categorized as:

Low: 5-10%; Medium: 10-30%; High: more than 30%

Genetic Advance (GA)

Genetic advance refers to the expected gain or improvement in the next generation by selecting superior individuals under certain amount of selection pressure. From the heritability estimates the genetic advance was estimated.

$$GA = h^2(bs) \cdot \sigma_p \cdot K$$

Comment [SA4]: Author of the formula

Where,

$h^2(\text{bs})$ = Heritability in broad sense

K = Selection differential, constant value is 2.06 at 5% selection intensity

σ_p = Phenotypic Standard Deviation

Genetic Advance as per cent of Mean (GAM)

In order to visualize the relative utility of genetic advance among the characters, genetic advance as per cent of mean was computed.

$$\text{GAM} = \frac{\text{GA}}{\bar{X}} \times 100$$

Where,

GA = Genetic Advance

X = General mean of the character

The range of genetic advance as per cent of mean was classified Low (=less than 10%); Moderate (=10-20%) and High (= more than 20%)

as suggested by Johnson *et al.* (1955)

~~Low = less than 10%; Moderate = 10-20%; High = more than 20%~~

RESULTS AND DISCUSSION

The analysis of variance (Table 1) for 48 pea genotypes with regards to yield traits ~~has~~ revealed ~~that their significance of mean squares for all traits suggesting existence of sufficient variability and hence presenting scope for t amount of variation that could be further used in the further selection breeding programmes. The mean sum of squares of all the fourteen characters included in the study revealed to be highly significant among the treatments that indicates there is variation and these genotypes can be selected further. Whereas the replication mean sum of squares of all the traits revealed to be non significant which indicates that there is reduction in the experimental error among the genotypes.~~ Various genetic variability parameters like genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), broad sense heritability, genetic advance (GA) and genetic

advance as percentage of mean (GAM) for ~~fourteen-14~~ characters are ~~furnished-represented~~ in the Table 2.

Days to 50% flowering:For the following trait PCV and GCV values were found to be 3.90 and 4.21 respectively. Heritability value (85%) was found to be very high whereas genetic advance value was low (4.98). GAM value was found to be low (7.46).

Number of primary branches:Moderate values of GCV and PCV for the above trait was observed to be 10.07 and 13.56 respectively. Moderate level of heritability (55%) and low level of genetic advance (1.11) values were observed. GAM value was found to be moderate (15.42).

Number of secondary branches:High values of GCV and PCV for the above trait was observed to be 21.35 and 22.65 respectively. Very high level of heritability (88%) and low level of genetic advance (6.19) values were observed. GAM value was found to be high (41.45).

Plant height:High values of GCV and PCV for the above trait was observed to be 38.08 and 38.46 respectively. Very high level of heritability (98%) and high level of genetic advance (70.40) values were observed. GAM value was found to be high (77.67).

Days to maturity:Low values of GCV and PCV for the above trait was observed to be 2.81 and 4.09 respectively. Medium level of heritability (47%) and low level of genetic advance (4.71) values were observed. GAM value was found to be low (3.99).

Pod length:Moderate values of GCV and PCV for the above trait was observed to be 13.35 and 15.22 respectively. High level of heritability (76%) and low level of genetic advance (1.56) values were observed. GAM value was found to be high (24.13).

Number of seeds per pod:Moderate value of GCV (18.85) and high value of PCV (22.8) for the above trait was observed respectively. High level of heritability (68%) and low level of genetic advance (1.58) values were observed. GAM value was found to be high (32.10).

Number of pods per plant:High values of GCV and PCV for the above trait was observed to be 27.81 and 28.31 respectively. Very high level of heritability (96%) and moderate level of genetic advance (15.16) values were observed. GAM value was found to be high (56.29).

100 Seed weight:Moderate value of GCV (18.94) and high value of PCV (20.19) for the above trait was observed respectively. Very high level of heritability (87%) and low level of genetic advance (5.26) values were observed. GAM value was found to be high (36.60).

Grain yield per plant: Moderate values of GCV and PCV for the above trait was observed to be 15.48 and 17.02 respectively. Very high level of heritability (82%) and low level of genetic advance (5.04) values were observed. GAM value was found to be high (29.03).

Harvest index: High values of GCV and PCV for the above trait was observed to be 22.92 and 24.50 respectively. Very high level of heritability (87%) and moderate level of genetic advance (13.18) values were observed. GAM value was found to be high (44.19).

Number of nodules per plant: Low value of GCV (9.79) and moderate value of PCV (11.32) for the above trait was observed respectively. High level of heritability (74%) and low level of genetic advance (3.41) values were observed. GAM value was found to be moderate (17.46).

Nodules fresh weight: High values of GCV and PCV for the above trait was observed to be 39.45 and 40.29 respectively. Very high level of heritability (95%) and low level of genetic advance (1.26) values were observed. GAM value was found to be high (79.58).

Nodules dry weight: High values of GCV and PCV for the above trait was observed to be 38.64 and 39.33 respectively. Very high level of heritability (96%) and low level of genetic advance (0.38) values were observed. GAM value was found to be high (78.20).

~~Based on the above categorization, all the fourteen traits are divided and t~~The findings revealed that high PCV and GCV values ~~are-were~~ recorded ~~in-for~~ characters like number of secondary branches, plant height, number of pods per plant, harvest index, nodules fresh weight and nodules dry weight suggesting scope for substantial improvement through selection of these traits in later generations. These findings are in a line to the findings reported by (4,5,6).

Similarly moderate PCV and GCV values ~~are-were~~ recorded ~~for~~ traits like number of primary branches, pod length, number of seeds per pod, 100 seed weight, grain yield per plant and number of nodules per plant suggesting considerable amount of variability among the genotypes. These findings are in a line to the findings reported by (7,8,9,10).

Low magnitude of PCV and GCV was noted for the characters like days to 50% flowering and days to maturity representing the existence of low variability and selection of these traits are may not be worthwhile in early generations. These findings are in a line to the findings reported by (11,12,13,14).

High heritability coupled with high genetic advance estimates was recorded for number of secondary branches, plant height, pod length, number of seeds per pod, number of pods per plant, 100 seed weight, grain yield per plant, harvest index, nodules fresh weight

Comment [SA5]: The entire content seems to be a direct lift from results chapter of thesis. Discussion totally lacking

Comment [SA6]: Influence of environment should be taken in to consideration. How is the value of PCV as compared to GCV?

and nodules dry weight indicating the role of additive gene action and selection for these traits is reliable. These findings are in a line to the findings reported by (15, 16, 17).

High heritability and moderate genetic advance were recorded for number of primary branches and number of nodules per plant indicating the presence of both additive and non-additive gene action. These findings are in a line to the findings reported by (18,19). While high level of heritability and low genetic advance for days to 50% flowering and days to maturity indicating the role of non-additive genes in the inheritance of these traits. These findings are in a line with the findings reported by (20,21,22) and improvement of these traits through mere selection may not be fruitful and is not useful therefore breeder should go for heterosis breeding or recurrent selection could be resorted to, for improvement of these characters.

CONCLUSION:

As all the traits under evaluation significantly differed from one another, the current study on variance revealed significant variability among all the genotypes. The characters viz., number of secondary branches, plant height, pod length, number of pods per plant, nodule fresh weight and nodules dry weight showed high value for GCV, heritability and genetic advance as percentage of mean. Therefore, additive gene effects control these traits. Thus, when selecting pea cultivars to increase yield, these traits should be taken into consideration.

Comment [SA7]: redraft

Table 1. Analysis of variance for fourteen-14 characters in field pea

Comment [SA8]: Units for traits

S. No	Character	Range		Mean sum of squares			C.V. %
		Min	Max	Replication (d.f = 2)	Treatments (d.f = 47)	Error (d.f = 94)	
1	Days to 50% flowering	61.66	73.33	1.89	21.58**	1.12	1.58

Formatted Table

2	Number of primary branches	5.33	9.33	1.02	2.03**	0.43	9.08
3	Number of secondary branches	8.33	20.33	3.88	31.82**	1.28	7.57
4	Plant height	49.66	162.33	6.88	3598.93**	24.04	5.41
5	Days to maturity	110	124.66	13.77	45.55**	12.32	2.97
6	Pod length	4.93	8.9	0.05	2.47**	0.22	7.31
7	Number of seeds per pod	3.33	6.66	1.02	3.01**	0.40	12.85
8	Number of pods per plant	15	47.66	4.69	170.54**	2.03	5.28
9	100 seed weight	9.89	21.55	2.55	23.25**	1.01	7.00
10	Grain yield per plant	11.74	24.78	1.69	23.28**	1.51	7.06
11	Harvest index	17.9	50.24	18.44	147.12**	6.66	8.64
12	Number of nodules per plant	15.93	23.8	1.13	12.27**	1.23	5.67
13	Nodules fresh weight	0.35	2.95	0.039	1.21**	0.02	8.17
14	Nodules dry weight	0.17	0.92	0.004	0.11**	0.001	7.34

** Significant at p = 0.01 % level

UNDER PEER REVIEW

Table 2. Estimates of variability parameters in fourteen field pea genotypes**Comment [SA9]:** Values may be restricted to 2 decimal places

S. No	Character	Genotypic variance	Phenotypic variance	GCV (%)	PCV (%)	Heritability (%)	Genetic Advance	Genetic Advance as % of mean
1	Days to 50% flowering	6.8220	7.9447	3.9081	4.2174	85	4.9858	7.4601
2	Number of primary branches	0.53	0.96	10.07	13.56	55	1.11	15.42
3	Number of secondary branches	10.18	11.45	21.35	22.65	88	6.19	41.45
4	Plant height	1191.63	1215.66	38.08	38.46	98	70.40	77.67
5	Days to maturity	11.07	23.39	2.81	4.09	47	4.71	3.99
6	Pod length	0.75	0.97	13.35	15.22	76	1.56	24.13
7	Number of seeds per pod	0.86	1.27	18.85	22.8	68	1.58	32.10
8	Number of pods per plant	56.17	58.19	27.81	28.31	96	15.16	56.29
9	100 seed weight	7.41	8.42	18.94	20.19	87	5.26	36.60
10	Grain yield per plant	7.25	8.76	15.48	17.02	82	5.04	29.03
11	Harvest index	46.82	53.47	22.92	24.50	87	13.18	44.19
12	Number of nodules per plant	3.67	4.91	9.79	11.32	74	3.41	17.46
13	Nodules fresh weight	0.39	0.41	39.45	40.29	95	1.26	79.58
14	Nodules dry weight	0.035	0.036	38.64	39.33	96	0.38	78.2

Where, GCV- genotypic coefficient of variance, PCV - phenotypic coefficient of variance.

REFERENCES

1. Dabholkar A.R. 2001. Elements of biometrical genetics. *Concept Publishing Company*, New Delhi, pp. 188-195.
2. Singh, S., Verma, V., Singh, B., Sharma, V.R and Kumar, M. 2019. Genetic variability, heritability and genetic advance studies in pea (*Pisum sativum* L.) for quantitative characters. *Indian J. Agric. Res.*, 53 (5): 542-547
3. Johnson, H.W., Robinson, H.F and Comstock, R.E. 1955. Estimates of genetic and environmental variability in soybean. *Agronomy Journal*. 47 (7): 314-318.
4. Naveen Kumar, J., Ajeet Kumar, G., Hiralal, D. and Roopa, L.G. 2012. Genetic Variability Analysis in Field Pea (*Pisum sativum* L.). *International Journal of Science and Research*. 4 (1): 2006-2010.
5. Meena, B.L., Das, S.P, Meena, S.K., Kumari, R., Devi, A.G and Devi H.L. 2017. Assessment of GCV, PCV, heritability and genetic advance for yield and its components in field pea (*Pisum sativum* L.). *Int. J. curr. microbiol. appl. sci.* 6(5):1025-33.
6. Bahadur, L and Devi, B. 2021. Estimation of correlation and path analysis coefficient among yield and yield attribution traits of field pea (*Pisum sativum* L.). *J pharmacogn. phytochem.* 10(1):1696-9.
7. Jeberson, M.S., Shashidhar, K.S and Iyanar, K. 2016. Estimation of genetic variability, expected genetic advance, correlation and path analysis in field pea (*Pisum sativum* L.). *Electron. J Plant Breed.* 7(4):1074-8.
8. Pathak, V.N., Pandey, R.K., Verma, S.P., Ray, J., Singh, B and Jee, C. 2019. Studies of genetic variability, heritability and genetic advance for yield contributing traits in field pea (*Pisum sativum* L.). *J pharmacogn. phytochem.* 8(4):2587-2589.
9. Pratap, V., Sharma, V and Kamaluddin, G.S. 2021. Assessment of genetic variability and relationship between different quantitative traits in field pea (*Pisum sativum* var. *arvense*) germplasm. *Legum. Res.* 1:6.
10. Pujari, P. K., Salam, J. L., Sao, A., Mandavi, N. C. and Singh, D. P. 2021. Study of genetic variability in field pea (*Pisum sativum* L.). *Journal of PharmacognosyPhytochemistry*, 10 (1): 1053-1055.
11. Afreen, S., Singh, A. K., Moharana, D. P., Singh, V., Singh, P. and Singh, B. 2017. Genetic evaluation for yield and yield attributes in garden pea (*Pisum sativum*

Comment [SA10]: Uniform format may be followed. In some cases, journal names are abbreviated. In some, they are not

- var. *hortense* L.) under North Indian gangetic plain conditions. *International Journal of Current Microbiology Applied Sciences*, 6 (2): 1399-1404.
12. Singh, J and Dhall, R. K. 2018. Genetic variability parameters of yield and quality attributes in vegetable pea (*Pisum sativum* L.). *Genetika*, 50(1):153-170.
 13. Kumar, N., Mishra, S., Pandey, S., Mishra, D. P. and Pandey, V. P. 2019. Studies on phenotypic and genotypic coefficient of variation (PCV and GCV) between different characters in Pea (*Pisum sativum* L.) genotypes in sodic condition. *Journal of Pharmacognosy Phytochemistry*, 8(1): 349-351.
 14. Kalapchieva, S., Kosev, V and Vasileva, V. 2020. Genetic and phenotypic assessment of garden pea (*Pisumsativum* L.) genotypes. *Basrah Journal of Agricultural Sciences*, 33(1): 107-121.
 15. Rai M, Verma A, Kumar R, Vishwanath., Multivariate genetic analysis of pea (*Pisum sativum*). *Veg. Sci.* 2006; 33(2):149-154.
 16. Tiwari G, Lavanya GP. 2012. Genetic variability, character association and component analysis in F4 generation of field pea (*Pisum sativum* var. *arvense* L.). *Karnataka J. Agric. Sci.* 25(2):173-175.
 17. Kumar B, Kumar A, Singh AK, Lavanya GR. 2013. Selection strategy for yield and maturity in field pea (*Pisum sativum* L. *arvense*). *Global Sci. Res.*1(1):129-133.
 18. Georgieva N, Ivelina N, Valentin K. 2016. Evaluation of genetic divergence and heritability in pea (*Pisum sativum* L.). *J BioSci. Biotechnol.* 5(1):61-67.
 19. Ahmad, H.B., Rauf, S., Rafiq, M., Mohsin, A.U and Iqbal, A. 2014. Estimation of genetic variability in pea (*Pisum sativum* L.). *J Glob. Innov. Agric. Soc. Sci.* 2(2):62-64.
 20. Dar, S.A., Ishfaq, A., Ali, G., Pir, F.A and Abu M. 2013. Study on genetic variability, heritability and genetic advance in pea (*Pisum sativum* L.). *Ann. Hort.* 6(1):161-163.
 21. Singh, N and Dhillon, G.S. 2012. Genetic variability for pod yield and its contributing traits in pea. (*Pisum sativum* L.). *Haryana J Hort. Sci.* 33(3):300-301.
 22. Sharma, V.K and Sharma, B.B. 2013. Genetic variability, heritability and genetic advance studies in garden pea under mid hill condition of Garhwal Himalaya. *Environmental and Ecology.* 31:296-301.