

**Response of spring green gram cultivars to (*Vigna radiata* L.)
integrated nutrient management in Bundelkhand region of Uttar Pradesh**

ABSTRACT

A field experiment was conducted to study the effect of integrated nutrient management on growth and yield of *spring* mungbean (*Vigna radiata* (L.) Wilczek) cultivars in Bundelkhand region of Uttar Pradesh during season of 2019 at Banda University of Agriculture and Technology, Banda (Uttar Pradesh). The experiment was laid out in asymmetrical factorial randomized block design with 6 treatment combinations of integrated nutrient management practices with 2 varieties. Results revealed that cultivar “Shikha” suits to be the better over “Samrat” in respect of growth, yield attributes, seed yield and economics. Additionally, INM practices had better effect on growth, yield attributes compared to sole application of inorganic fertilizers and vermicompost treatments. 100% RDF + 100% vermicompost had highest seed yield (771 kg/ha), closely followed by 75% RDF + 50% vermicompost (741 kg/ha) compared to rest of treatments. Among the INM treatments, application of 50% RDF blended with 50% vermicompost recorded at par values of growth, yield attributes and seed yield over the application of 100% RDF + 100% vermicompost and 75% RDF + 50% vermicompost. On an average, INM treatments noted 69.89, 14.07 and 43.39% higher net returns as well as 68.92, 21.2 and 51.57% more benefit: cost ratio over the control, 100% RDF and 100% vermicompost, respectively. Therefore, application of 50% RDF + 50% vermicompost (1.25 t/ha) was optimum under the existing condition of Bundelkhand.

Key words: Mungbean, Integrated nutrient management, Yield and Economics

Introduction

Pulses are the important sources of proteins, with high fiber content and low glycemic index. [Pulses](#) occupies the predominant position in discussions related to food and nutritional security. India has appreciably 35% and 26% share in Global area and production, respectively, and the largest producer and consumer of pulses. Realizing the importance of Pulses for human diet, year 2016 was declared as international year of pulses. However, the growth in production and yield of pulses has lagged the growth rate of country's population

and this resulted a decline in per capita availability of pulses from 60.7 g/day during triennium ending 1951 to 55.9 g/day (P) during 2018 (Agricultural Statistics at a Glance, 2018). The Bundelkhand is known to be the pulse basket of India which comprises 7 districts from Uttar Pradesh and 6 districts from Madhya Pradesh.

In Uttar Pradesh, Mungbean occupies 89.5 thousand hectares of area with production of 44.1 thousand metric tonnes and a productivity of 478 kg/ha during the same year. Alone Bundelkhand region of Uttar Pradesh occupies 33.7% in area and 21.0% in total production (Anonymous, 2019). Average yield of this crop in the country is very low, which is primarily due to substandard methods of cultivation, poor crop stands, imbalanced nutrition, lack of high yielding varieties, etc.

The production and productivity of this crop is highest but due to lack of technological intervention in this region and high use of land reforms in spring season the yields and its attributing character lacks behind. Thus, an experiment was conducted at BUAT with the hypothesis that Response of spring green gram cultivars to (*Vigna radiata* L.) integrated nutrient management in Bundelkhand region of Uttar Pradesh

Formatted: Font: Italic

Materials and Methods

The experimental farm lies between 25°31'36.98" North latitude and 80°21'31.92" East longitude at an elevation of 123 MSL with hot and semi-arid climatic zone. Average rainfall of this region ranges [what is the range, please mention](#)... however, its distribution pattern is uncertain. Initially pH, organic carbon, available nitrogen, available phosphorous and available potassium of the soil samples (0-15cm) under the experimental field results 7.2, 0.41%, 211 kg/ha, 8.85 kg/ha and 215 kg/ha respectively.

The experiment was laid out in an asymmetrical factorial randomized block design with the 12 treatments and each treatment were replicated thrice to take the mean value. Treatments comprised the combination of two cultivars viz., Shikha (V_1) and Smart (V_2) and six INM treatments with absolute control viz.,

Control (N_0), 100% recommended dose of fertilizer-RDF (N_1), 100% Vermicompost-VC @ 2.5 t/ha (N_2), 100% RDF+100% VC @ 2.5 t/ha (N_3), 75% RDF+50% VC @ 1.25 t/ha (N_4) and 50% RDF + 50% VC @ 1.25 t/ha (N_5). Organic fertilizer (vermicompost) was applied during the final land preparation. Seeds were initially treated by *rhizobium* PSB @ 3 kg/ha mixed with 50 kg well decomposed FYM was broadcasted over the entire experimental land during last ploughing. Mung bean was sown on 1

March 2019 @ 20 kg/ha with 30 cm row spacing. Other recommended practices were adopted till harvesting (15 May, 2019) was adopted. The data in this paper represents the average value of three randomly selected plants from every plot at 30 and 45 DAS for growth analysis and a harvest for yield attributes. The mature pods were picked when they attained their full maturity.

Results and Discussion

The results invoked that the variety 'Shikha' produced numerically high values of Nodulation/plant and branches/plant with almost yield attributes viz., number of pods/plant, seeds/pod and 1000-seed weight variety 'Shikha' over variety 'Samart' (Table – 1). The significant difference among varieties over control plot were attributed might be due to genomic makeup which truly indicated gross photosynthesis production (references). Additionally, seed yield, net return and benefit: cost ratio were recorded comparable values with variety 'Shikha' with corresponding figures of 652 kg/ha, Rs. 23346 /ha and 0.97, respectively (Table 1). Among fertilizer treatments (N₃) had maximum number of branches/plant, number of root nodules/plant and its dry weight, followed by 75% RDF + 50% vermicompost and 50% RDF + 50% vermicompost, respectively. The least values of Nodulation/plant and branches/plant were recorded with the unfertilized plot. In addition to above, it has also been observed that INM treatments recorded at par compared to control, 100% RDF and 100% vermicompost treatments for almost yield and its attributing characters 46.45, 11.78 and 24.63% respectively (Table 1). It might be due to slow mineralization of organic manure and non-availability of required nutrients resulted in setback of crop growth at early stage of crop growth and thus affected the crop yield (references). The plot receiving no fertilizer (control) recorded the lowest values of seed yield (400 kg/ha) of mung bean.

Table 1. Effect of integrated nutrient management practices on growth, yield attributes and seed yield of mung bean cultivars

Formatted: Indent: Left: 0", Hanging: 3.3 ch, First line: -3.3 ch

| Treatment | Branches/ plant | Nodulation/plant at 30 DAS | | Pods per Plant | Seeds per pod | Test weight | Seed yield (kg/ha) |
|-------------|--------------------|-------------------------------|-----------------|-------------------|------------------|----------------|-----------------------|
| | | Counts | Dry weight (mg) | | | | |
| Variety (V) | | | | | | | |
| V1: Shikha | 5.92 | 45.85 | 73.18 | 17.33 | 5.40 | 37.36 | 652 |

| | | | | | | | |
|-----------------------------|-------|-------|-------|-------|------|-------|-----|
| V2:Smart | 4.36 | 37.42 | 64.88 | 15.04 | 4.78 | 35.98 | 635 |
| SE(d)± | 0.53 | 4.87 | 9.23 | 0.84 | 0.22 | 2.41 | 14 |
| CD(P=0.05) | NS | NS | NS | 1.76 | 0.46 | NS | NS |
| INMtreatments (N) | | | | | | | |
| N0;Control | 4.63 | 25.65 | 47.29 | 9.17 | 4.33 | 29.11 | 400 |
| N1;100%RDF | 5.28 | 40.92 | 64.59 | 14.18 | 5.08 | 37.77 | 659 |
| N2;100% VC(2.5t/ha) | 4.78 | 34.32 | 57.17 | 13.51 | 4.65 | 34.72 | 563 |
| N3;100%RDF+100% VC | 6.28 | 51.48 | 86.59 | 24.91 | 6.19 | 40.29 | 771 |
| N4; 75%RDF +50%VC 5t/ha | 6.01 | 50.54 | 80.77 | 18.75 | 5.20 | 39.64 | 741 |
| N550% RDF+50% VC (1.25t/ha) | 5.56 | 46.92 | 77.77 | 16.62 | 5.11 | 38.50 | 729 |
| SE(d)± | 0.925 | 8.44 | 15.99 | 1.46 | 0.38 | 4.18 | 25 |
| CD(P=0.05) | NS | 17.63 | NS | 3.05 | 0.81 | NS | 35 |
| Interactions | NS | NS | NS | NS | NS | NS | NS |

DAS= Days after sowing; RDF= Recommended dose of fertilizer; VC= Vermicompost; NS=Non significant

Table2. Effectofintegratednutrientmanagementpracticesoneconomicsof mungbeancultivars.

| Treatment | Grossreturn (₹/ha) | Netreturn (₹/ha) | B:Cratio |
|-----------------------------|-----------------------|---------------------|----------|
| Variety(V) | | | |
| V1:IPM2-3 | 46969 | 23346 | 0.97 |
| V2:IPM205-7 | 45738 | 22091 | 0.92 |
| SE(d)± | 1044 | 1045 | 0.04 |
| CD(P=0.05) | NS | NS | NS |
| INMProtocols(N) | | | |
| N0;Control | 28784 | 8684 | 0.43 |
| N1;100%RDF | 47457 | 24783 | 1.09 |
| N2;100% VC(2.5t/ha) | 40525 | 16327 | 0.67 |
| N3;100%RDF+100% VC(2.5t/ha) | 55529 | 28757 | 1.75 |
| N4;75%RDF+50% VC(1.25t/ha) | 53346 | 29049 | 1.19 |
| N5;50%RDF+50% VC(1.25t/ha) | 52482 | 28713 | 1.21 |
| SE(d)± | 1809 | 1810 | 0.076 |
| CD(P=0.05) | 3776 | 3778 | 0.15 |
| Interactions | NS | NS | NS |

Grossreturn,netreturnandbenefit:costratiowere higher values recorded with variety

'Shikha' being a part variety 'Smart' with corresponding figure of ₹ 23346/ha, ₹ 22091/ha and 0.97, respectively (Table 2). Among the different INM treatments, the mean effect of INM irrespective of doses (mean of N₃, N₄ and N₅) registered significantly higher gross (₹ 53785/ha) and net return (₹ 28839/ha) to the tune of 11.24 and 24.65% gross and 14.07 and 43.39% more net return compared to sole application of 100% RDF and 100% vermicompost, respectively (Table 2). The benefit: cost ratio (1.75) was higher 100% RDF and 100% vermicompost treatments. The superiority in gross and net returns with these treatments was mainly attributed to their higher yield of green gram. However, significantly lower gross and net returns as well as benefit: cost ratio was obtained with the control plot. Among the INM treatments, the maximum gross (₹ 55529/ha) and benefit: cost ratio (1.75) found in 100% RDF+100% vermicompost (1.25t/ha), where net return (₹ 29049/ha) was recorded from the application of 75% RDF+50% vermicompost (1.25t/ha), closely followed by two other INM treatments i.e. N₃ and N₅.

Conclusion

The present research work reveals that the application of 50% RDF+50% vermicompost (1.25t/ha) to *spring* mung bean varieties is more beneficial for higher growth, yield and income as compared to other remaining integrated nutrient management practices studied.

References

- Agricultural Statistics at a Glance, 2018. Directorate of Economics and Statistics, Department of Agriculture, Cooperation and Farmers Welfare, Ministry of Agriculture and Farmers Welfare.
- Anonymous. 2019. Ministry of Agriculture and Farmers Welfare, Department of Agriculture, Cooperation and Farmers Welfare, Directorate of Economics and Statistics, Fourth Advance Estimates of Production of Food Grains for 2017-18.
- Arsalan, Ahmed Shoaib, Chauhdary Nawaz Junaid and Sarwar Muhammad. 2016. Effect of vermicompost and phosphorus on crop growth and nutrient uptake in mung bean. *Journal of Applied Agriculture and Biotechnology*, **1(2)**:38-46.
- Bhavya G, Chandra Shaker K, Jayasree G and Reddy M. Malla. 2018. Nutrient uptake and yield of

greengram (*Vigna radiata* L.) as influenced by phosphorus fertilization, organic manures and biofertilizers. *International Journal of Chemical Studies*, **6(3)**:32-35.

Chaudhary SK, Singh SP, Singh Y and Dharminder. (2014). Influence of integrated use of fertilizers and manures on SRI grown rice (*Oryza sativa*) and their residual effect on succeeding wheat (*Triticum aestivum*) in calcareous soil. *Indian Journal of Agronomy*, **59(4)**:527-533.

Gadi Parvati, Dawson Joy and M Shankar. (2017). Effect of different organic manures, inorganic fertilizer and growth regulator on yield attributes and yield. *International Journal of Current Research*, **9(06)**:52385-52389

Jawhara A.L., -Owied A.L (2016). Comparative study of synthetic fertilizers and organic manures on some mungbean (*Vigna radiata* (L.) Wilczek) genotypes I-Effect on growth parameters and photosynthesis productivity. *American-Eurasian Journal Agriculture and Environmental Science*, **16** (4): 666-676.

Krishnaprabu S. 2018. Integrated approach in nutrient management of greengram on nutrient uptake and economics. *International journal of Pure and Applied Bioscience*, **6(5)**:832-834.

Niranjan K. Rajeev, Pal Amit, Gupta H.K., Verma K Manish and Gond Prasad Dadhwal (2010). Performance of different vermicomposts on yield and yield components of mung bean in major soils of Bundelkhand region. *India Journal Ecophysiology and Occupational Health*, **10(1)**: 61-70