

Original Research Article

**Influence of Biofertilizers and Zinc on the Growth and Yield of Barley**

**ABSTRACT**

The field experiment entitled "Influence of Biofertilizers and Zinc on the Growth and Yield of Barley" was conducted during *Rabi* season, 2022 at Crop Research Farm in the Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj Uttar Pradesh. The treatment consisted of Biofertilizers [*Azotobacter*, PSB and *Azotobacter*+ PSB], Zn (20, 25 and 30 kg/ha) and control. The experiment was layout in Randomized Block Design (RBD) with 10 treatments and replicated thrice. The soil in the experimental area was sandy loam with pH (8.0), Organic Carbon (0.62%), Available N (225 kg/ha), Available P (38.2 kg/ha) and Available K (240.7 kg/ha). Application of *Azotobacter* + PSB and Zn 30 kg/ha recorded growth parameters and yield attributes like, highest plant height (107.63 cm), maximum number of tillers/running row meter (66.32), higher Plant dry weight (22.31g), maximum number of effective tillers/m<sup>2</sup> (278.85), maximum number of grains/spike (52.27), higher grain yield (4.29t/ha) and straw yield (6.72t/ha).

**Keywords:** Barley, Biofertilizers, Zinc, *Azotobacter*, PSB, Growth and Yield attributes.

**Introduction**

Barley (*Hordeum vulgare* L.) is the world's 4<sup>th</sup> most essential cereal crop after wheat, rice and maize with a share of about 7% of the global cereals production and 15% of coarse grains consumption. Barley is grown throughout the temperate, tropical and subtropical regions of the world and can be successfully grown in adverse climatic conditions of drought, salinity and alkalinity due to its wider adaptability (Neelam et al., 2018).

Barley is generally grown on marginal and sub-marginal land because of its low inputs requirement. Barley grain is also valued for smothering and cooling effect on the body for easy digestion and as a source of vitamin B complex. Besides these conventional uses, it is an important industrial crop used as raw material for beer, whisky and brewing industries. Each 100 g of barley grain comprise 10.6 g protein, 2.1 g fat, 64.0 g carbohydrate, 50.0 mg calcium, 6.0 mg iron, 31.0 mg vitamin B<sub>1</sub>, 0.1 mg vitamin B<sub>2</sub> and 50.0 µg folate (Vaughan et

*al.*, 2006). In India, barley is mainly grown in the northern plains and concentrated in the states of Uttar Pradesh, Haryana and Rajasthan. In India, barley was cultivated on 609 thousand ha area with 1818 thousand t of production at an average productivity of 29.88 q/ha. In India, Rajasthan is the largest state having more than 52 % in production and 46% area followed by Uttar Pradesh. In Uttar Pradesh, barley was cultivated on 159.0 thousand ha area with 498.0 thousand tonnes of production at an average productivity of 31.32 q/ha(IWBR, 2020-21).

Comment [a1]: Uttar

Climate change, increased population pressure, harmful environmental effects on agricultural fields, and ongoing chemical use result in a decrease in organic carbon, a decrease in the microbial flora of soil, an increase in acidity and alkalinity, and a hardening of the soil. These negative effects ultimately result in a shortage of food production. To overcome the situation, new mechanism must be developed to meet the increased food demands with sustainable food production that has the potentiality to provide adequate food nutrition without hampering the fields. "Biofertilizer" is one such method for addressing agricultural needs.

*Azotobacter* are abiotic, naturally occurring soil microorganisms that bind atmospheric nitrogen that is unavailable to plants and play a significant role in the nitrogen cycle in nature. Inoculation with *Azotobacter* has been found to reduce the requirement of chemical fertilizer upto 50 per cent (Soleimanzadeh and Gooshchi, 2013). *The most prevalent heterotrophic free-living bacterium, Azotobacter, is important for crop production.* Bio-fertilizer normally contains microorganisms which are having particular function such as N<sub>2</sub> fixation by *Azospirillum* and phosphorus solubilisation by P solubilizing bacteria from the soil and fertilizer which are to be available for plants (Saraswati and Sumarno, 2008).

"Several enzyme systems that control different metabolic processes in plants depend on zinc. It contributes to the synthesis of auxins, which are plant growth regulators. Zinc is also vital for the oxidation processes in plant cells and helps in the transformation of carbohydrates and regulates sugar in plants" Zinc helped to increase leaf area, chlorophyll content in leaves, uptake of total Zinc availability in soil agronomic efficiency, grain and stover yield of sorghum"(Jakhadet *et al.*, 2023).

## 2. MATERIALS AND METHODS

This experiment was laid out during the *Rabi* season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The crop research farm is situated at

25° 39' 42" N latitude, 81° 67' 56" E longitude and at an altitude of 98 m above mean sea level. The experiment was laid out in Randomized Block Design Which consisted of ten treatments with T<sub>1</sub> – *Azotobacter* + Zn20 kg/ha, T<sub>2</sub> – *Azotobacter* + Zn25 kg/ha, T<sub>3</sub> – *Azotobacter* + Zn30 kg/ha, T<sub>4</sub> - PSB + Zn20 kg/ha, T<sub>5</sub> - PSB + Zn25 kg/ha, T<sub>6</sub> - PSB + Zn30 kg/ha, T<sub>7</sub> - *Azotobacter* + PSB + Zn20 kg/ha, T<sub>8</sub> - *Azotobacter* + PSB + Zn25 kg/ha, T<sub>9</sub> - *Azotobacter* + PSB + Zn30 kg/ha, T<sub>10</sub> - Control (NPK 80-30-20 Kg/ha). Seeds are sown at a spacing of 23 cm × 5 cm to a seed rate of 100 kg/ha. The recommended dose of nitrogen (80 kg/ha), phosphorus (30 kg/ha) and potassium (20 kg/ha) in the form of Urea, DAP and MOP, respectively and Biofertilizer and zinc were applied as per the treatments. Data recorded on different aspects of the crop, viz., growth, yield attributes were subjected to statistically analysis by analysis of variance method. (Gomez and Gomez, 1976) and economic data analysis mathematical method.

## **RESULT AND DISCUSSION:**

### **Growth parameters**

#### **Plant height (cm)**

The data revealed that a significantly and higher plant height (107.68cm) was recorded in treatment 9 [*Azotobacter* + PSB + Zn30 kg/ha]. However, treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (106.00 cm), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (103.70 cm), were found to be statistically at par with treatment 9. The significant and higher plant height was observed with the application of Zinc might be due to Zinc involves in biosynthesis of indole acetic acid (IAA) which helps in better development of growth attributes. Similar result was reported by (Ganapathy *et al.* 2006). Further increase in plant height observed with the application of biofertilizers (20g/kg seed) might be due increase levels of biofertilizers application to improve growth by increasing the supply or availability of primary nutrients to the host plant. Crop development may have been aided by the higher endogenous nitrogen content brought on by inoculation. These results were supported by Aechraet *et al.* (2020).

#### **Number of tillers/running row meter**

The data revealed that a significantly and maximum number of tillers/running row meter (66.32) was recorded in treatment 9 [*Azotobacter* + PSB + Zn30 kg/ha]. However, treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (64.93), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (63.46) were found to be statistically at par with treatment 9. The application of biofertilizers resulted in the substantial and highest number of tillers/running row meter, which may be

attributable to the higher amount of biofertilizer application that has been shown to boost cereal tiller output, initially determined by rate of auxiliary bud growth and later on growth of individual tiller, which is markedly influenced by environmental factors (temperature, photoperiod etc.) as well as availability of mineral nutrients and photosynthates. The results are in close conformity with the findings of **Kekatpureet al., (2021)**.

#### **Plant dry weight (g)**

The data revealed that a significantly and maximum plant dry weight (22.31 g) was recorded in treatment 9 [*Azotobacter* + PSB + Zn30 kg/ha]. However, treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (21.65 g), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (21.25 g), were found to be statistically at par with treatment 9. Zinc in the soil and its participation in numerous enzymatic activities, as well as its position as a catalyst in various development processes and in the generation of hormones, may be the cause of the considerable and greater plant dry weight that was found with the application of zinc (30 kg/ha) and protein synthesis which results in increasing growth. Similar results were reported by (**Shekhawatet al., 2017**). Further increase in dry weight observed with the application of biofertilizers (20g/kg seed) might be due to biofertilizers stimulates activation of hormones which helps in shoot and root elongation and high dry matter production, similar results were observed by **Aktharet al., (2018)**.

#### **B. Post-harvest observation**

##### **Number of effective tillers/m<sup>2</sup>**

The data revealed that Treatment 9 [*Azotobacter* + PSB + Zn30 kg/ha] was recorded significant and maximum of effective tillers/m<sup>2</sup> (278.85) which was superior over all other treatments. However, treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (249.63), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (238.10), treatment 6 [PSB + Zn30 kg/ha] (236.98), treatment 5 [PSB + Zn 25 kg/ha] (234.85) was found to be statistically at par with the treatment 9. The application of biofertilizers (20 g/kg seed) resulted in a significant and maximum number of effective tillers/m<sup>2</sup>, which may be attributable to the increased availability of nitrogen for plant uptake. This growth promotion of plant characteristics, such as increased tillering, led to an increase in the number of effective tillers. These can be corroborated by similar findings by **Yadav et al. (2011)**.

### **Number of grains/spike**

The data revealed that Treatment 9 [*Azotobacter* + PSB + Zn30 kg/ha] was recorded a significant and maximum number of grain/spike (52.27) which was superior over all other treatments. However, treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (51.48), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (50.62), treatment 6 [PSB + Zn30 kg/ha] (50.04), treatment 5 [PSB + Zn 25 kg/ha] (49.37), treatment 4 [PSB + Zn20 kg/ha] (48.86) was found to be statistically at par with the treatment 9. Significant increase in number of grains/spike is due to increase in the availability of *Azotobacter* and PSB bio fertilizer inoculation by which more seeds were produced due to increased rate of production, similar results were found by (Joshi and Chilwal 2018).

### **Grain Yield (t/ha):**

The data revealed that aTreatment 9 [*Azotobacter* + PSB + Zn30 kg/ha], was recorded significantly maximum Grain yield (4.29 t/ha) which was superior over all other treatments. However, treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (4.20 t/ha), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (4.11 t/ha), treatment 6 [PSB + Zn 30 kg/ha] (4.06 t/ha), treatment 5 [PSB + Zn 25 kg/ha] (4.09 t/ha), was found to be statistically at par with the treatment 9. The significant and maximum seed yield was observed with the application of Biofertilizers (20g/kg seed) might be due increase levels of biofertilizers application increased yield attributes of barley might be due to release of growth hormones by various biofertilizers. Similar findings for yield attributes were reported by Diman and Dubey (2017). Significant and higher seed yield obtained with the application of zinc (25kg/ha) might be due to zinc improves the source and sink relationship due to increased translocation of photosynthates towards reproductive system (Sammuauria et al., 2010).

### **Straw yield (t/ha):**

The data revealed that aTreatment 9 [*Azotobacter* + PSB + Zn30 kg/ha] was recorded significantly maximum Straw yield (6.72t/ha) which was superior over all other treatments. However, the treatment 8 [*Azotobacter* + PSB + Zn25 kg/ha] (6.38 t/ha), treatment 7 [*Azotobacter* + PSB + Zn20 kg/ha] (6.22 t/ha) was found to be statistically at par with the treatment 9. Significant and higher Stover yield was obtained with the application of Zinc (25kg/ha) might be due to the Zinc increase in yields attributed to the fact that because of favorable nutritional environment in rhizosphere and higher absorption of nutrients by plant leading to the increased photosynthetic efficiency and production of assimilates. Similar

results were also reported by **Khan *et al.*, 2010**. Application of *Azotobacter* (20 g/kg) resulted in an even higher maximum straw yield. Significant Application of biofertilizer, which increases grain and straw yield, was the cause of the increase in straw yield. Similar results were reported by (**Patel *et al.* 2017**).

## CONCLUSION

Based on the finding it is concluded that seed inoculation with *Azotobacter* and PSB along with the application of Zn (30kg/ha) produce significantly higher plant height, dry weight, number of tillers/running row meter, number of effective tillers/m<sup>2</sup>, number of grains/spike, Grain yield and straw yield.

UNDER PEER REVIEW

**Table: 1 Effect of Biofertilizers and Zinc on growth of Barley.**

S.No.	Treatment combinations	At 120 DAS		
		Plant height (cm)	Number of tillers/running row meter	Dry weight (g/plant)
1.	<i>Azotobacter</i> + Zn 20 kg/ha	90.77	53.87	18.72
2.	<i>Azotobacter</i> + Zn 25 kg/ha	92.80	56.23	19.25
3.	<i>Azotobacter</i> + Zn 30 kg/ha	98.00	58.58	19.73
4.	PSB+ Zn 20 kg/ha	99.10	60.43	20.15
5.	PSB+ Zn 25 kg/ha	100.70	61.11	20.53
6.	PSB+ Zn 30 kg/ha	102.67	62.36	20.89
7.	<i>Azotobacter</i> + PSB+ Zn 20 kg/ha	103.70	63.46	21.25
8.	<i>Azotobacter</i> + PSB+ Zn 25 kg/ha	106.00	64.93	21.65
9.	<i>Azotobacter</i> + PSB+ Zn 30 kg/ha	107.63	66.32	22.31
10.	Control	88.27	50.25	18.15
	F-test	S	S	S
	SEm(±)	3.48	2.07	0.78
	CD(p=0.05)	10.36	6.16	2.34

**Table: 2 Effect of Biofertilizers and Zinc on yield attributes and yield of Barley.**

S.No.	Treatment combination	Number of effective tillers/m <sup>2</sup>	Number of grains/spike	Grain Yield (t/ha)	Straw Yield (t/ha)
1.	<i>Azotobacter</i> + Zn 20 kg/ha	204.86	45.51	3.77	4.99
2.	<i>Azotobacter</i> + Zn 25 kg/ha	207.75	46.35	3.87	5.64
3.	<i>Azotobacter</i> + Zn 30 kg/ha	213.16	47.78	3.90	5.74
4.	PSB+ Zn 20 kg/ha	224.51	48.86	3.77	5.82
5.	PSB+ Zn 25 kg/ha	234.85	49.37	4.09	5.92
6.	PSB+ Zn 30 kg/ha	236.98	50.04	4.06	6.06
7.	<i>Azotobacter</i> + PSB+ Zn 20 kg/ha	238.10	50.62	4.11	6.22
8.	<i>Azotobacter</i> + PSB+ Zn 25 kg/ha	249.63	51.48	4.20	6.38
9.	<i>Azotobacter</i> + PSB+ Zn 30 kg/ha	278.85	52.27	4.29	6.72
10.	Control (RDF)	188.25	42.49	3.58	4.78
	F-test	S	S	S	S
	SEm(±)	15.84	1.54	0.12	0.21
	CD(p=0.05)	47.07	4.59	0.36	0.64

## REFERENCE

1. Aechra, S., Meena, R. H., Jat G., Sharma, J., Doodhwal, K. and Jat, H. (2020). Effect of Biofertilizers and Split Application of Vermicompost on Productivity and Profitability of Wheat (*Triticumaestivum* L.) Crop in Clay Loam Soils. *International Journal of Current Microbiology and Applied Sciences* **9**(4): 1129-1139.
2. Akhtar, N., Ramani. V. B., Yunus, M., Vala, Femi. (2018). Effect of Different Nutrient Management Treatments on Growth, Yield Attributes, Yield and Quality of Wheat (*Triticumaestivum* L.). *International Journal of Current Microbiology and Applied Sciences* **7**:3473-3479.
3. Dhiman, S. and Dubey, Y.P. (2017). Effect of biofertilizers and inorganic fertilizers on yield attributes, yield and quality of *Triticumaestivum* and *Zea mays* in an acid alfisol. *International Journal of Current Microbiology and Applied Sciences* **6**(7): 2594-2603.
4. Ganapathy, B. A. and Savalgi, V. P. (2006). Effect of micronutrients on the performance of *Azospirillum* *Brasiliense* on the nutrient uptake, growth and yield in maize crop. *Karnataka Journal Agricultural Sciences*. **19**(1):66-70.
5. Gomez, K. A. and Gomez, A. A. (1976). Statistical procedures for agriculture Research, 2<sup>nd</sup> Edition, John Wiley and Son, New York, 680p.
6. Joshi, G. and Chilwal, A. (2018). Effect of integrated nutrient management on growth parameters of baby corn (*Zea mays* L.). *International Journal of Advances in Agricultural Science and Technology*, **5**(7): 216-225.
7. IIWBR. (2020-21). Progress Report, All India Coordinated Wheat and Barley Improvement Project. ICAR-Indian Institute of Wheat and Barley Research, Karnal, Haryana **6**: pp-1.1
8. Jakhad, A. and Debbarma, V. (2023). Influence of biofertilizer and zinc on growth, yield and economics of sorghum (*Sorghum bicolor* L.). *International Journal of Plant and Soil Science*, **35**(8), 90-97.
9. Kekatpure, K. and Chaturvedi, D. P., (2021) Growth and Yield Response of Wheat (*Triticumaestivum* L.) in Relation to the Use of Varieties and Bio-Fertilizer. *Indian Journal of Pure Applied Bioscience* **9**(6), 53-57.
10. Khan MB, Farooq M, Hussain M, Shanawaz, Shabir G. (2010). Foliar application of micronutrients improves the wheat yield and net economic return. *International Journal of Agriculture and Biology*. **12**: 953-956.
11. Neelam, S.B., Khippal, A. and Mukesh and Satpal. (2018). Effect of different nitrogen levels and biofertilizers on yield and economics of feed barley. *Wheat and Barley Research* **10**(3): 214-218.

12. Patel PR, Patel BJ, Vyas KG and Yadav B. (2017). Effect of integrated nitrogen management and bio-fertilizer in *Kharif* pearl millet (*Pennisetum glaucum* L.). *Advance Research Journal Crop Improvement*. **5**(2):122-125.
13. Sammauria, R. and Yadav, R. S. (2010). Response of pearl millet (*Pennisetum glaucum*) to residual fertility under rainfed conditions of arid region of Rajasthan. *Indian Journal of Dryland Agricultural Research and Development*. **25**(1):53-60.
14. Saraswati, R. and Sumarno, (2008). Application of soil microorganisms as component of agriculture technology. *Iptek Dibrang Pangan*, **3**, 41.
15. Shekhawat, P. S. and Kumawat, N. (2017). Response of zinc fertilization on production and profitability of pearl millet (*Pennisetum glaucum*) under rainfed condition of Rajasthan: Zinc fertilization for improving production and profitability of pearl millet. *Journal of Agri Search*. **4**(4):251- 254.
16. Soleimanzadeh, H. and Gooshchi, F. (2013). Effect of Azotobacter and nitrogen chemical fertilizer on yield and yield component wheat (*Triticum aestivum* L.). *World Applied Science Journal* **21**(8): 118.
17. Vaughan, J.G., Judd, P.A. and Bellamy, D. (2006). The Oxford Book of Health Foods. Available online at <http://books.google.co.in/books> pp. 37.
18. Yadav, D. D., Verma C K, Singh B P and Shanker S (2011). Role of biofertilizers in relation to nitrogen levels on growth and yield of wheat (*Triticum aestivum* L.). *Crop Research* **42**: 23-26.