

Review Article

Quality Enhancement In Forage Crops Of Cereals And Legumes

ABSTRACT

This article focus on quality improvement in forage crops, specifically cereals, legumes, and grasses. As milk and many other by-products obtained from dairy farms, poultry, etc are getting into demand, high-quality forage for livestock production is well recognized, and efforts to improve forage quality can have significant economic and environmental benefits. And an overview of current strategies for quality improvement, including breeding and genetic selection, management practices, and how these practices affect the quality of forage crops. Use of different breeding methods more over like synthetic cultivars, recurrent selection, etc. the use of biotechnology tools such as RNAi interference, tissue culture, and marker-assisted selection in the particular crop that has been developed for increasing the nutritional value, proteins that present in forage crops, and to decrease the harmful chemicals. Additionally, the paper discusses the challenges associated with quality improvement in forage crops and potential solutions. And about different types of grasses used for different types of cattle as supplements. Limitations that cause obstruction to improving the quality of forage crops. digestibility and how to increase the quality and protein content in milk and cattle. And nutrition that needs to be present in prescribed quantities in forage crops. Overall, this paper highlights the need for continued research and innovation in the field of forage crop quality improvement to support sustainable agriculture and meet the increasing demand for high-quality livestock feed.

Keywords: factors affecting, limitations, digestibility, nutritional value

Introduction

Forage crops are the plants or products of dry fodder used to feed ruminants for milk, dairy products, and meat production. However, the area under forage agriculture in India was 13.8 million hectares in 2019, whereas the total area under forage cultivation worldwide was 1.11 billion hectares in 2020, according to the most recent information available from the Food

Comment [u1]: explain the type according to the introduction

Comment [u2]: Enter the related nutrients

Comment [u3]: enter the source of the theor

and Agriculture Organization (FAO) of the United Nations. It's crucial to remember that these numbers might alter over time depending on changes in farming techniques, the weather, and other variables. Forage quality has a wide range of definitions. Agronomy experts and animal science researchers have recently learned to rely on ruminant animals' biological responses to make accurate assessments of the quality of fodder. High-grade forage is described to be leafy, fine-stemmed, green, sweet-smelling, rich in amino acids, low in fiber, and delicious (Moore, 1980). Forage crops are a key source of feed for livestock; hence, forage crop, quality enhancement is important for the livestock business (Moore, 1980). The health of the cattle that consume forage crops is intimately correlated with the quality of those crops. Forage crops were very recently domesticated, with major landrace and quality improvement in forage breeding beginning in the early 20th century (Harlan, 1992). Numerous research studies have looked at various methods, such as choosing high-quality plant varieties, applying fertilizer, and using growth regulators, to enhance the quality of fodder crops. Grasses (Poaceae), herbaceous legumes, or fodder cereals are used as forage crops (Fabaceae). Some tree legumes, like River tamarind (*Leucaena leucocephala*) and mulga (*Acacia aneura*), are also cultivated in desert areas and sward (Muir *et al.*, 2011). The predominant grasses in temperate regions include orchard grass (*Dactylis spp*), bent grass (*Agrostis spp*), fescue (*Festuca spp*), ryegrass (*Lolium spp*), and hybrids of this date back to the 1970s. Cereals are inexpensive and produce a lot of dry stuff, hence they are widely employed in livestock nutrition (Ghanbari-Bonjar and Lee, 2003). The variety of cereals cultivated for fodder, such as maize, oats, pearl millet, and rye, has gained relevance in the diets of ruminant animals (Eskandari *et al.*, 2009). The third-most significant cereal crop worldwide, maize (*Zea mays*), is used as food, animal feed, and forage. In contrast to sorghum or other fodder, feeding maize at any stage of growth is harmless and poses no risk from hydrocyanic or oxalic acid (Dahmardeh *et al.*, 2009). Oats (*Avena sativa L.*) have distinct benefits over other fodder species due to their high production potential, nutritional value, and great ability for regeneration, especially during the first few months of winter (Alipatra *et al.*, 2013). In the lower Midwest countries pastures with a combination of annual ryegrass (*Lolium multiflorum Lam*) and cereal rye (*Secale cereale L*) have become popular for grazing in the late winter and early spring (Altom *et al.*, 1996). The annual ryegrass and cereal rye pastures offer several alluring qualities. The two forage plants produce 70% of their yearly development while many tree species are either dormant or not experiencing severe environmental stress, which is perhaps the most significant (Altom *et al.*, 1996), (Zhang *et al.*, 1997), (Balandier *et al.*, 2000). During the low summer months of

May to July and in conjunction with other fodder crops during the kharif and summer seasons, pearl millet is a potential crop for the provision of green fodder. During the dry season, when there is a shortage of green fodder and grazing, cattle in marginal production areas are also fed with pearl millet's dry fodder and straw. As a result, it is typically cultivated in locations where the climate, particularly rainfall, temperature, and soil fertility, is too severe for the growth of other cereals (Gulia *et al.*, 2007).

Comment [u4]: enter theoretical sources

Forage legumes such as alfalfa, clover, and soybean, are high in amino acids, whereas grasses are rich in energy from both structural and non-structural carbohydrates (Humphreys, 1997). Because of its excellent nutritional content and productivity, alfalfa is the feed crop that is most often grown worldwide (Skot. *et al.*, 2007). The fodder crop feed differs or changes depending on the country and soil conditions available. Similar patterns may be seen for pearl millet in Rajasthan, where the crop's straw is a significant source of animal feed (Kelley *et al.*, 1996). Red clover is a popular feed legume in the United States, and it is often used as a cover crop to improve the soil environment (Taylor, 2008).

Certain tropical locations combine tropical legumes, largely from Asia, with grass species, especially from Africa and the Americas, as demonstrated by pasture growth in tropical Queensland, Australia (Walker and Weston, 1990). Have been thoroughly assessed and widely applied in cattle production systems. Forage legumes are ideal for use as roughage in the diet of animals because of their abundance of protein, vitamins, and minerals, whether they are fresh, dried, or stored (Vasiljevic *et al.*, 2009). They are distinguished as roughage because of their high fiber content (above 18% crude cellulose). In comparison to concentrate, these feeds have a smaller proportion of accessible (digestible) energy per unit weight or volume, and the majority of energy is present as cellulose or hemicellulose (Jonker *et al.*, 2010).

Factors affecting and improving forage quality

Environmental, genetic, and management factors

The quality of forage crops changes according to climate, soil quality, and water availability. Nearly all cool-season forage species are suitable for growing the humid regions, except perhaps for reed canary grass *Phalaris arundinacea* L (Sheaffer *et al.*, 1990).

Forage crops can achieve a certain percentage of their potential productivity in a given area depending on factors like temperature and rainfall, but the primary force that directly

determines how much can be produced is solar radiation through photosynthesis (Allard *et al.*, 1991). Legume forage plants seek to reduce light penetration through the canopy by displaying their leaf area more horizontally than grasses. Similar to this, the leaf area index (LAI), which measures the number of leaf blades per unit area of the soil surface, is lower for legumes around 4-5 than for grasses about 6-8 and is necessary to maximum radiation interception. Because of this, it is essential for the management of legume-grass combinations that the legume does not shadow the grass, even though the quality would be greater. Plants grow better forage when they are in the shadow (Buxton *et al.*, 1993). Although this is not always true (Allard *et al.*, 1991).

Comment [u5]: enter the source of the theor

According to research conducted by Vough and Marten 1971, forage quality is often enhanced by water stress, while heat stress has little impact on it. According to Pembleton alfalfa grown under water stress had greater quality (IVDMD) than alfalfa grown in normal water circumstances, and better quality under stress was brought about by a delay in development (Pembleton *et al.*, 2010). According to a study by Reid, the level of oxalate in (*Atriplex halimus*) plants can be affected by soil moisture levels (Reid and Jung, 1974). The research suggests that high soil moisture can lead to alterations in the oxalate level in these plants. One of the most important inputs for the production of crops is water. Not only does it directly affect crop performance, but it also indirectly does so by affecting the availability of nutrients, the timing of cultural activities, and other variables. With no irrigation, forage sorghum yields varied from 38.3 t per ha to 88.4 t per ha with 56 mm of irrigation (Nabati and Moghadam, 2010). According to Abdel, splitting the same amount of irrigation water into more frequent irrigation resulted in a greater benefit (Abdel *et al.*, 1982). The absorption of N, P, and K as well as the production of dry matter were enhanced when three to four irrigations were applied to barley throughout the active tillering, flag leaf, and milk phases (Wahab and Singh, 1983). Soybean produced the driest matter when irrigated with a ratio of 0.6 IW: CPE (Veeramani *et al.*, 2000). The yields of grains, fodder, and straw are influenced by genomic and non-genetic factors, which vary between and within crop species. Crop varieties with higher grain and feed yields, as well as better straw quality, may be chosen or bred. According to research, wheat crop residue quality varies (Doyle *et al.*, 1987), Rice (Walli *et al.*, 1988), barley (Ramanzin *et al.*, 1986), oats (Shand *et al.*, 1988), finger millet (Subba Rao *et al.*, 1993), sorghum (Badve *et al.*, 1993), and maize (Harika and Sharma *et al.*, 1994). The proportion of variation in grain and fodder production, as well as straw digestibility that is attributable to genetic factors versus non-genetic components such as environmental

conditions, crop management practices, and after-harvest methods, can vary depending on the crop species and the specific genotype within a given crop species (Pembleton *et al.*, 2010).

Application of N up to 120 kg per ha resulted in higher levels of dry matter, CP, and green forage while lowering NDF levels (Bebawi, 1988), (Patil *et al.*, 1992). As per the research conducted by Patel (Patel *et al.*, 1993), the impact of phosphorus or potassium on the production and quality of fodder is not as significant as that of nitrogen. The study shows that the application of P₂O₅ did not affect the protein content or yield, indicating that the production and quality of fodder are not greatly influenced by the application of phosphorus. Crude fiber decreased with increased N, while treatment of P had little effect (Humphreys *et al.*, 2010). Temperate forage legumes have also seen similar rates of genetic improvement as ryegrass. Research indicates that breeding advancements in yield and quality attributes for these legumes have also been estimated to be around 3.8 and 4.0% per WSC and annual dry matter yield maximized during that time period, respectively. These rates of improvement have been observed since the early 20th century (Woodfield and Brummer, 2001); (Ali and Rawat, 1986).

According to research on various cropping systems, drought-tolerant crops like forage sorghums or pearl millet should be mono cropped during dry years as they are drought-tolerant varieties (Yadav and Kumar, 1997). According to statistics on green forage yields, barley was the plant that tolerated soil salinity the best, followed by oats, sorghum, pearl millet, Egyptian clover, and maize (Ismaeil *et al.*, 1993). Salinity reduced seed germination and early seedling development in sorghum. Pre-treatments with CaCl₂ and ZnSO₄ also boosted sorghum sowing rate and percentage (Assefa and Ledin, 2001). Oats can withstand wet circumstances better than the majority of other cereals forage (Liu *et al.*, 2015). The quality of oats can be increased by inducing the water stress conditions and use of inorganic fertilizers than organic. **the use of an N:P ratio of 112:45 with manure increases the quality of forage oats.** Enhancing the nutritional value of fodder crops requires certain agronomic and breeding techniques, some of which are mentioned in Table 1.

Comment [u6]: should be completed with the results of these plants

Forage crop	Mineral nutrients status	Strategy used	Mechanism of action	References
White clover	Increased mineral uptake (shoots: Mg and Cu; roots: Ca,	administration of sodium nitroprusside, a	Increase the activity of the plasma membrane enzymes V-H+-	(Nawaz <i>et al.</i> , 2016)

Comment [u7]: table do not cut

	Mg, and Fe)	NO donor, exogenously	ATPase and H ⁺ -ATPase in the root and shoot.	
Fodder maize	enhanced se (36%), complete basic amino acids (40%), CP (47%), crude fibres (10%), and extract without nitrogen (10%).	Foliar Se supplementation	The osmoprotectants that se accumulates help to keep the turgor pressure constant. activate the antioxidant defense system to control physiological and biochemical processes.	(Contreras-Govea <i>et al.</i> , 2010)
Fodder maize, sorghum	a lower NDF and a higher net energy	Plant population decline and N fertilizer application	increase the effectiveness of resource use and reduce energy waste.	(Hack <i>et al.</i> , 2019)
White sweet clover	higher amounts of N and P in the root zone and stem	the bacterium Rh meliloti and Arbuscular Mycorrhizal Fungi (AMF) inoculation	Inoculants increase physiological nitrogen fixing and phosphorus accessibility in a reciprocal manner.	(Binder <i>et al.</i> , 2020)
Cereal Rye + Corn	decreased NDF (10%), elevated observed N recovery (84%), enhanced CP (29%), and decreased net lactation energy (10%).	pumping soil with liquid dairy cow excrement.	Enhance soil capacity to hold water and N and P restoration.	(Gallego & Giraldo <i>et al.</i> , 2016)

Table.1 Represents the soil and agronomic methods applied to enhance the nutritional value of fodder.

Comment [u8]: preferably above the table

Limitations:

The crop protein content, amino acid composition, and fiber content all affect how nutritious they are. There is genetic heterogeneity in the protein content and acid profile of cereals and legumes used as feed crops, according to several studies (Wang *et al.*, 2012). Having limited genetic diversity it is difficult for selecting desirable traits. While being of excellent quality, alfalfa protein is not well used (Sheaffer *et al.*, 1990). The rapid breakdown of soluble crude protein in alfalfa can lead to a disadvantage in terms of its quality, as it may result in the wastage of high-quality protein in the rumen. Furthermore, the breakdown of alfalfa amino

acid can cause an increase in the solidity of the rumen fluid, which can put animals at risk of pasture bloat (Kaur *et al.*, 2018). Another important aspect that has an impact on the quality of cereal and legume fodder crops is the environment. The nutritional value of various crops can be impacted by environmental conditions including temperature, light, and water availability. High temperatures, for instance, can lower the protein level of grains and legumes, and environmental factors that the forage crop are the same with every crop(Hou *et al.*, 2020). The quality of cereal and legume feed crops can also be impacted by management techniques, including fertilization, irrigation, and harvesting. While nitrogen fertilization can raise these crop protein content, overusing nitrogen can lower or increase the quality of their fodder crops(Hou *et al.*, 2020). The genetic diversity in maize is limited because of the narrow genetic base of commercial maize varieties. This limits the potential to improve quality traits in forage maize and reduces the quality traits, such as digestibility, which can lead to a reduction in yield and other agronomic traits (Osman, 2018). Environmental factors such as Drought stress can lead to reduced yields and lower-quality forage maize and pearl millet(Bondar *et al.*, 2018). Improper harvesting can result in increased levels of lignin and reduced digestibility(Divya *et al.*, 2017). While in pearl millet poor storage, conditions can result in mold and spoilage, leading to reduced quality forage(Dhaliwal *et al.*, 2020). Forage oats contain anti-nutritional factors, such as phytate, which can reduce the bioavailability of minerals, such as iron and zinc These factors can also have an impact on the digestibility of forage oats, reducing their nutritional value(Zhu *et al.*, 2016). In rye such as ergot alkaloids, which can be toxic to livestock (Asay *et al.*, 1985). Pest and disease attack is common in forage clover. Common pests include clover root weevils, which can reduce yield and quality, and aphids, which can transmit viruses and reduce plant growth(Vogel and Pederson, 1993).

Breeding Methods for quality improvement of forage crop

Forage breeders must overcome obstacles in breeding fodder crops, like inherited and polymorphic variability, polyploid nature, and polygenic regulation of agronomic and nutritional qualities. Traditional breeding methods have been successful in improving forage crops, but they are time-consuming(Oram and Culvenor, 1994). Before producing a synthetic cultivar with a variable number of parents, some type of recurrent selection technique is frequently made use of because many forage species are perennial plants(Gates *et al.*, 199). believe that groups of half-sibling families chosen under space-planted and sward conditions will be important for Phalaris to make up recurrent selection. Breeding white clover and alfalfa mostly uses phenotype recurrent selection (Collard and Mackill, 2008). This has been

Comment [u9]: complete the theoretical source

accomplished using both traditional breeding methods and genetic engineering, in which fodder crops have had genes linked with enhanced nutritional quality put into them(Hanna, 1993).

Future breeding will make use of the chances offered by new technology to accurately examine and alter genotype and phenotype. Using DNA profiling and marker-assisted selection, more focused breeding plans may be created for a larger variety of qualities. The raising of fodder crop nutritional quality is one area that has drawn a lot of interest. Forage crops with high quantities of protein, fiber, and energy have been the focus of research since these nutrients are crucial for the growth and well-being of cattle (Beever and Reynolds, 1994).

Breeding fodder crops to be more precisely adapted to animal demands is a difficult task. It found that advances in NIR near infrared reflectance technology enable breeders to assess a variety of feed quality criteria more quickly, particularly those related to the number and quality of animals(Marshall *et al.*, 2016).

In the majority of our breeding fodder crops, synthetic cultivars have been the primary breeding product. These only sparingly employ the heterosis that emerges from broad crossings across several gene pools (Burton and Mullinix, 1998). A better-regulated application of hybrids might lead to major gains in many forages. This tactic is significantly affecting the rates of maize breeding advancement Some perennial fodder plants, such as Bermuda grass, have had their heterosis taken advantage of successfully by vegetative multiplication(Kumar *et al.*, 2021). According to Dudley and Lambert, 1992 maize grain has undergone 90 cycles of selection for either high or low CP concentration, as well as high or low oil concentration, resulting in consistent progress over time. Plant growth-promoting rhizobacteria (PGPR) improves plant growth and yield Panchagavya, which contains macro and micronutrients, growth-regulating chemicals, and useful microorganisms, might help provide appropriate plant nutrients, providing proper plant nutrients, hence boosting fodder quality and production(Brown *et al.*,1980),(Harrington, 1952) outlined certain procedures for the artificial hybridization of oats. Oat crosses can be developed outdoors, indoors, or in a greenhouse. Excellent plants to form crosses can be found in conditions that are ideal for plant development and growth. Although growth chamber crosses can be performed at any time, the majority of greenhouse crosses take place during the winter when it is simple to maintain the cooler temperatures needed for the best outcomes. Change the photoperiod and

Comment [u10]: complete the theoretical source

make the light more intense, specifically during the cloudy and brief winter days, when additional light is needed. Excellent outcomes have been attained in the greenhouse at Urbana, Illinois, using metal halide lighting and a 13-hour photoperiod. According to Harrington, 1952(Kumar *et al.*, 2012). the pedigree system has been especially effective for rust resistance in oat breeding. By crossing the pearl millet x Napier hybrid the dry matter potential of both plants can be combined. To create the inter-specific hybrid commercially, CMS pearl millet and napier pollinators are planted in a 1:1 ratio Sukanya *et al.*, 2001). Pandey *et al.*, 2019 examined 30 hybrids for the number of leaves per plant, their length, breadth, and relation to their stems, as well as their green fodder production by crossing three pearl millet genotypes and ten Napier genotypes. All the characters exhibited a lot of variety. Pearl millet (IP 6426) and Napier (FD 439), which were both effective general combiners, were outperformed by IP 6426 X FD 469 as a hybrid for green fodder yield and leafiness. Although some rye species have had success with repeated phenotypic selection on spaced plants (Gates *et al.*, 1999); (Burton and Mullinix, 1998). Improved agronomic management techniques and plant breeding techniques, both traditional and innovative techniques, can produce the required forage qualities(Figure-1)(Pandey *et al.*, 2019).




Figure.1) Represents a blend of agronomic, conventional, and modern breeding methods to increase the nutrient content of fodder crops. (Pandey *et al.*, 2019)

Biotechnology tools for forage crop quality improvement

Turf and forage grasses are a broad collection of plants that exhibit clump-forming or sod-like growth patterns, annual or perennial life cycles, and cool- or warm-season growth preferences. In North America, breeding for better fodder and turf grasses began seriously in the 1930s and 1940s, particularly with the initiatives of L. C. Newell in Nebraska and G. W. Burton in Georgia. Numerous forage types of grass coexist with other plant species. Grasses differ substantially in terms of chromosome count, reproductive department, incompatibility relationships, and practice of pollination(Vogel *et al.*, 1991). Cultivars have been produced with verified improvements in attributes including fodder quality(Asay *et al.*, 1985). and seedling strength(Lowe and conger, 1979).

Comment [u11]: complete the theoretical source

The development of a callus culture technique the ability to rejuvenate plants cornerstone of genetic engineering in monoclonal grass species at the level of cells. To create a tissue culture system, the original explant must first be stimulated to generate calluses. Different explants, such as fescues have produced seeds and mature or undeveloped embryos, and ryegrass, have been used to start regenerable callus cultures in grasses (Dale, 1980; Torello *et al.*, 1984; Hanning and Conger, 1986). leaf bases in orchard grass(Vasil, 1981), and immature bud and shoot apices in pearl millet *Pennisetum Americanum*(Artunduaga *et al.*, 1988) and Scutch grass *Cynodon dactylon*(Zhang and Main, 2003). Advancements in genomics research have led to the development of innovative tools like functional molecular markers, and a deeper understanding of inheritance patterns that can significantly enhance the accuracy and efficacy of crop improvement in alfalfa. This progress in genomics research can pave the way for better crop management and increase agricultural productivity. Pearl millet uses DNA markers to generate genetic linkage maps. Important characteristics like disease adaptability, insect resistance, drought resistance, etc. have DNA markers associated with them(Boora *et al.*, 2009). Marker-assisted selection (MAS) is a novel method that effectively raises the quality of fodder. (Hash *et al.*, 2003) used Marker aided selection (MAS) and quantitative traits loci (QTL) mapping to increase stover yield, increase forage disease resistance, and figure out the nutritional value of different pearl millet residue portions for ruminants. Only two forage species, perennial ryegrass(Cogan *et al.*, 2006) and white clover, have seen SNP - single nucleotide polymorphism - markers described for them. Based on single base pair alterations in a DNA order, these biological indicators(Cogan *et al.*, 2007). Disease resistance and water-soluble carbohydrate content were discovered to be markers for traits in perennial ryegrass(Dracatos *et al.*, 2008);(Skot *et al.*, 2007). SNP markers are certain to be due to their enormous frequency and the swift technological advancements for high-throughput

Comment [u12]: you should mention all the characters if you don't need to delete them

generation and identification, they will be crucial in the advancement of marker-assisted fodder crop breeding in the future. Another biotechnological tool for forage crop improvement is RNA interference (RNAi). RNAi is a process that enables the downregulation of specific genes by inducing the degradation of the corresponding messenger RNA (mRNA). RNAi has been used to reduce the expression of lignin biosynthesis genes in forage crops, leading to increased digestibility and improved nutritional quality (Gallego & Giraldo *et al.*, 2016). Transgenic rye expressing the gene for rumen-stable amylase has been developed to increase the starch digestibility of rye silage (Selinger *et al.*, 1996).

NUTRITIONAL VALUE

Nutritional value in forage crops is not only beneficial to animals, but it also helps in overcoming pressure conditions during the grazing of animals. Every fodder crop has enough cellulose, quartz, and other phenolic substances to cut down on ruminant consumption (Van Soest, 2018). These substances could serve as a general defensive strategy for plants to grow or multiply when challenged with grazing pressure (McCarthy *et al.*, 1989).

One of a forage's key traits is its ability to nutritional value is digestibility. It serves as a measure of the ruminant's energy availability. It also has an impact on how soon forage particles exit the rumen since feed particles need to do so after being sufficiently broken down (Poppi *et al.*, 1985; Waghorn *et al.*, 1989). Legumes are a substantial source of protein and can make up for cereal deficiency in that regard (Gebrehiwot *et al.*, 1996). Companion planting which involves growing crop mixtures next to legumes can boost the quantity of forage protein in diets. The forage crops' ability to serve as a source of feed for animals depends on maintaining their quality. Research has focused on developing techniques of long-term fodder crop quality preservation, such as ensiling and drying (Baker *et al.*, 2019). Researchers have also looked at the nutritional potential of conserved fodder crops, for instance, using inoculants that contain bacteria that hasten fermentation (Thomas, 2019). When fed more sugar ryegrass to graze on, Charolais steers devoured 20% more feed and put on 25% more live weight (Marley *et al.*, 2017). Maize often gets far less attention than it merits despite having more forage quality characteristics, such as being highly palatable, having high nutritional contents, lasting a longer time in storage, and being easily digestible (Iqbal *et al.*, 2006). inferior-quality chemicals like hydrocyanic acid (HCN) and oxalate, it has superior quality compared to Because it doesn't contain any sorghum and pearl millet

(Gupta *et al.*, 2004). Crude protein (CP) concentration is the most important factor compared to other factors which influence a crop's quality for use to be fodder (Caballero *et al.*, 1995); (Asefa and Ledin, 2001). According to research, oats have a higher crude protein concentration in the first cut (12.10–15.63%) than in the second cut (9.63–13.57%) Table-2 (Poonia and Phogat, 2017).

Crop	Phenol (mg/g)	Protein (mg/g)	Phytic (mg/g)	Beta-glucan (mg/g)	Zinc (mg/100g)	Iron (mg/100g)
Oats (<i>Avena sativa</i>)	11.90-31.3	138.70-160.50	3.70-8.00	31.00-53.50	4.96-6.50	2.48-4.89

(Table.2) Nutrients and quality parameters available in oats.(Poonia and Phogat, 2017)

Forage maize has a crude protein content of 7.5-8.5%, a crude fiber content of 32-34%, and a typical fat content of 1-2.5%. Approximately 32–34% of the material is dry, 7-9% is ash, and 50–50% is a nitrogen-free extract (Iabal *et al.*, 2015).

Digestibility

Studying the improvement of fodder crop digestibility has also been a priority. A feed crop's ability to be more successfully used by animals is encouraged by its higher digestibility, which improves animal performance (Chaabane. *et al.*, 2020) reported that increasing the cell wall digestibility of fodder crop varieties and implementing relevant agronomic strategies have been effective in improving digestibility. Additionally, in vitro, the incubation of fodder in rumen fluid is a reliable method for predicting digestibility (Deinum. *et al.*, 1984).

The percent of each component that has vanished in the animal digestive tract serves as a standard measurement of the digestibility of all forage components, including dry matter, organic matter, and cell walls. Indigestibility can alternatively be thought of as gm.kg^{-1} of metabolic weight, although it is often expressed as kg DM per animal and per day (live weight 0.75) (Barrière *et al.*, 2003). According to (Barrière *et al.*, 2003), the digestibility of various constituents of forage (such as dry matter, organic matter, or cell wall) is commonly calculated as the proportion of each component that is lost in the animal's digestive tract. Indigestibility is typically measured in kilograms of dry matter per animal per day, but it can also be expressed as grams per kilogram of metabolic weight (live weight to the power of 0.75). It is crucial to enhance WSC in many fodder crops, especially grasses because protein digestion and absorption in cattle are strongly connected to power availability (ME) (McCarthy *et al.*, 1989); (Miller *et al.*, 2001)

Marley *et al.*, 2017 investigated how fertilization affects the nutritional content of grass-clover swards in one study. According to the results, adding more nitrogen (N) to the swards significantly boosted the amount of crude protein (CP) while simultaneously enhancing digestibility. The Italian ryegrass cultivar "*Tribune*," which, when given as silage, produces 6% more milk, was developed as a result of raising the plant's stem digestibility (Wilman and Ahmad, 1999). (Zhang *et al.*, 2020) conducted a second study to examine how growth regulators affect the quality of alfalfa. The scientists found that the addition of gibberellic acid (GA3) increased the height and green color of alfalfa plants, which resulted in a considerably larger content of total digestible nutrients (TDN) and crude protein (CP). Casler, 2001 reported that during the 20th century, advancements in breeding techniques resulted in an increase of 1.0, 14.7, and 6.5% per generation cycle for ryegrass characteristics related to digestibility, intake, and crude protein, respectively. In ruminants, some types of compounds, such as sugars and organic acids are entirely digested. Whereas lignin, cutin, silica, and tannins are nearly indigestible, proteins are fairly easily digested. Forage-maize digestibility can be significantly increased if this variation is employed in breeding programs. The digestibility of cellulose and hemicellulose varies depending on whether they are encrusted with lignin (Deinum *et al.*, 1986). Digestibility in forage crops can be calculated using several methods, including in vivo digestibility trials, in vitro techniques, and near-infrared reflectance spectroscopy (NIRS). Here are examples of how to calculate digestibility using two commonly used methods (Van Soest, P. J., 2018).

1. In vivo digestibility trials:

In vivo, digestibility trials involve feeding animals a known amount of forage, collecting fecal samples, and analyzing them for nutrient content. The difference between the nutrient intake and the fecal nutrient content is used to calculate the digestibility of the forage. One commonly used formula for digestibility is:

$$\text{Digestibility (\%)} = 100 - (\text{fecal nutrient content} / \text{forage nutrient intake} \times 100)$$

Comment [u13]: enter the source of the theory

2. In vitro techniques:

In vitro, techniques involve using laboratory equipment to simulate the digestive process and estimate nutrient digestibility. The most commonly used in vitro method is the two-stage in vitro digestibility (TIVD) method, which involves incubating the forage with rumen fluid and then with intestinal fluid. The difference between the nutrient content before and after

incubation is used to calculate digestibility. One commonly used formula for TIVD digestibility is:

$$\text{Digestibility (\%)} = \frac{(\text{nutrient content before incubation} - \text{nutrient content after incubation})}{\text{nutrient content before incubation}} \times 100$$

Comment [u14]: enter the source of the theory

Conclusion

In conclusion, the review targeted at enhancing the nutritional quality, and digestibility of forage crops has finally in substantial advancements in the field of forage crop quality enhancement. These initiatives have produced fodder crops that are more nutrient-dense, easily digested, and stable, all of which are advantageous to the livestock sector.

Comment [u15]: State the type of feed according to the description and theory

The use of biotic and abiotic factors, breeding methods, and biotechnological tools affects and improves the quality of feed crops. For livestock producers wanting to maximize the quality of their fodder crops to full fill the dietary needs of their animals, the knowledge from the research is invaluable.

References:

Abdel Magid, EA., Mustafa, MA., and Ayed, L. (1982). Effects of irrigation interval, urea, and gypsum on N, P and K uptake by forage sorghum on highly saline-sodic clay. *Experimental agriculture*.

Ali, M., and Rawat, CR. (1986, October). Production potential of different food fodder cropping systems under dryland conditions of Bundelkand. In *Proceedings of the National Seminar on Advances in Forage Agronomy and Future Strategy for Increasing Biomass Production*. Indian Grassland and Fodder Research Institute, Jhansi, Uttar Pradesh, India (pp. 6-7).

Alipatra, A., Kundu, CK., Mandal, MK., Banerjee, H., and Bandopadhyay, P. (2013). Yield and quality improvement in fodder oats (*Avena sativa L.*) through the split application of fertilizer and cutting management. *Journal of Crop and Weed*, 9(2), 193-195.

Allard, G., Nelson, C J., and Pallardy, SG. (1991). Shade effects on growth of tall fescue: I. Leaf anatomy and dry matter partitioning. *Crop Science*, 31(1), 163-167.

Altom, W., Rogers, JL., Raun, WR., Johnson, GV., and Taylor, SL. (1996). Long-Term Rye-Wheat-Ryegrass Forage Yields as Affected by Rate and Date of Applied Nitrogen. *Journal of Production Agriculture*, 9(4), 510-516.

- Asay, KH., Dewey, D R., Gomm, F B., Johnson, D A., and Carlson, J. R. (1985). Registration of 'Bozoisky Select' Russian wildrye. *Crop Science*, 25(3), 575-576.
- Assefa, G., and Ledin, I. (2001). Effect of variety, soil type and fertiliser on the establishment, growth, forage yield, quality and voluntary intake by cattle of oats and vetches cultivated in pure stands and mixtures. *Animal feed science and technology*, 92(1-2), 95-111.
- Badve, VC., Nisal, PR., Joshi, AL., and Rangnekar, D. V. (1993). Variation in quality of sorghum stover. Feeding of Ruminants on Fibrous Crop Residues: Aspects of Treatment, Feeding, Nutrient Evaluation.
- Balandier, P., Lacoïnte, A., Le Roux, X., Sinoquet, H., Cruiziat, P., and Le Dizès, S. (2000). SIMWAL: a structural-functional model simulating single walnut tree growth in response to climate and pruning. *Annals of Forest Science*, 57(5), 571-585.
- Barker, AR., Muck, RE., and Mertens, DR. (2019). Preservation and quality of forage crops. In *Forage Quality, Nutrition and Feeding* (pp. 563-582). John Wiley and Sons, Ltd.
- Barnes, RF., Miller, DA., Nelsen, CJ., and Heath, ME. (1995). Forages. v. 1. An introduction to grassland agriculture.
- Barrière, Y., Guillet, C., Goffner, D., and Pichon, M. (2003). Genetic variation and breeding strategies for improved cell wall digestibility in annual forage crops. A review. *Animal Research*, 52(3), 193-228.
- Bebawi, FF. (1988). Forage Sorghum Production on a Witchweed Infested Soil in Relation to Cutting Height and Nitrogen. *Agronomy Journal*, 80(3), 537-540.
- Binder, JM., Karsten, HD., Beegle, DB., and Dell, CJ. (2020). Manure injection and rye double cropping increased nutrient recovery and forage production. *Agronomy Journal*, 112(4), 2968–2977. <https://doi.org/10.1002/agj2.20181>
- Boora KS., P. Boora, Urvashi and Sonika 2009 : In: Forage symposium on "Emerging trends in Forage research and livestock production. Feb.16-17,2009 at CAZRI, RRS, Jaisalmer (Rajasthan) India,pp.17-28.
- Brown, PD., McKenzie, RIH., and Mikaelson, K. (1980). Agronomic, Genetic, and Cytologic Evaluation of a Vigorous New Semidwarf Oat 1. *Crop Science*, 20(3), 303-306.

Burton, GW., and Mullinix, BG. (1998). Yield distributions of spaced plants within Pensacola bahiagrass populations developed by recurrent restricted phenotypic selection. *Crop science*, 38(2), 333-336.

Buxton, DR., and Casler, M. D. (1993). Environmental and genetic effects on cell wall composition and digestibility. *Forage cell wall structure and digestibility*, 685-714.

Caballero, R., Goicoechea, EL., and Hernaiz, PJ. (1995). Forage yields and quality of common vetch and oat sown at varying seeding ratios and seeding rates of vetch. *Field crops research*, 41(2), 135-140.

Casler, MD. (2001). Breeding forage crops for increased nutritional value. *value*.

Chaabane, M., Bejaoui, S., Trabelsi, W., Telahigue, K., Chetoui, I., Chalhaf, M., ... and Soudani, N. (2020). The potential toxic effects of hexavalent chromium on oxidative stress biomarkers and fatty acids profile in soft tissues of *Venus verrucosa*. *Ecotoxicology and Environmental Safety*, 196, 110562.

Chand, S., Singhal, RK., and Govindasamy, P. (2022). Agronomical and breeding approaches to improve the nutritional status of forage crops for better livestock productivity. *Grass and Forage Science*, 77(1), 11-32.

Chaudhary, JD., Pavaya, RP., Malav, JK., Dipika, G., Chaudhary, N., Kuniya, NK., ... and Jat, J. R. (2018). Effect of nitrogen and potassium on yield, nutrient content and uptake by forage sorghum (*Sorghum bicolor* (L.) Moench) on loamy sand. *International Journal of Chemical Studies*, 6(2), 761-765.

Cogan, N. OI., Drayton, MC., Ponting, RC., Vecchies, AC., Bannan, R., Sawbridge, TI., ... and Forster, JW. (2007). Validation of in silico-predicted genic SNPs in white clover (*Trifolium repens* L.), an outbreeding allopolyploid species. *Molecular Genetics and Genomics*, 277(4), 413-425.

Cogan, NO., Ponting, RC., Vecchies, AC., Drayton, MC., George, J., Dracatos, PM., ... and Forster, JW. (2006). Gene-associated single nucleotide polymorphism discovery in perennial ryegrass (*Lolium perenne* L.). *Molecular Genetics and Genomics*, 276(2), 101-112.

Collard, BC., and Mackill, DJ. (2008). Marker-assisted selection: an approach for precision plant breeding in the twenty-first century. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1491), 557-572.

- Dahmardeh, M., Ghanbari, A., Syasar, B., and Ramroudi, M. (2009). Effect of Intercropping Maize (*Zea mays L.*) VVith Cow Pea (*J'igna unguiculata L.*) on Green Forage Yield and Quality Evaluation. *Asian journal of plant sciences*, 8(3), 235-239.
- Dale, PJ. (1980). Embryoids from cultured immature embryos of *Lolium multiflorum*. *Zeitschrift für Pflanzenphysiologie*, 100(1), 73-77.
- Devos, KM., and Gale, MD. (1997). Comparative genetics in the grasses. *Oryza: From Molecule to Plant*, 3-15.
- Dhaliwal, SS., Sandhu, AS., Shukla, AK., Sharma, V., Kumar, B., and Singh, R. (2020). Bio-fortification of oats fodder through zinc enrichment to reduce animal malnutrition. *J. Agric. Sci. Technol. A*, 10, 98-108.
- Divya, G., Vani, KP., Babu, PS., and Devi, KS. (2017). Yield attributes and yield of summer pearl millet as influenced by cultivars and integrated nutrient management. *International Journal of current microbiology and applied sciences*, 6(10), 1491-1495.
- Doyle, PT., Chanpongsang, S., Wales, WJ., and Pearce, GR. (1987). Variation in the nutritive value of wheat and rice straw. *Ruminant Feeding Systems Utilizing Fibrous Agricultural Residues. IDP, Australia*, 75-86.
- Dracatos, PM., Cogan, NOI., Dobrowolski, MP., Sawbridge, TI., Spangenberg, GC., Smith, KF., and Forster, JW. (2008). Discovery and genetic mapping of single nucleotide polymorphisms in candidate genes for pathogen defence response in perennial ryegrass (*Lolium perenne L.*). *Theoretical and Applied Genetics*, 117(2), 203-219.
- Dudley, JW. (1992). Ninety generations of selection for oil and protein in maize. *Maydica*, 37, 81-87.
- Eskandari, H., Ghanbari, A., and Javanmard, A. (2009). Intercropping of cereals and legumes for forage production. *Notulae Scientia Biologicae*, 1(1), 07-13.
- Gallego Giraldo, L., Shadle, G., Shen, H., Barros Rios, J., Fresquet Corrales, S., Wang, H., and Dixon, RA. (2016). Combining enhanced biomass density with reduced lignin level for improved forage quality. *Plant Biotechnology Journal*, 14(3), 895-904.
- Gates, RN., Hill, GM., and Burton, GW. (1999). Response of selected and unselected bahiagrass populations to defoliation. *Agronomy Journal*, 91(5), 787-795.

- Gebrehiwot, L., McGraw, RL., and Assefa, G. (1996). Forage yield and quality profile of three annual legumes in the tropical highlands of Ethiopia. *Tropical agriculture*.
- Ghanbari-Bonjar, A., and Lee, HC. (2003). Intercropped wheat (*Triticum aestivum L.*) and bean (*Vicia faba L.*) as a whole-crop forage: effect of harvest time on forage yield and quality. *Grass and forage science*, 58(1), 28-36.
- Gulia, SK., Wilson, JP., Carter, J., and Singh, BP. (2007). Progress in grain pearl millet research and market development. *Issues in new crops and new uses*, 196-203.
- Gupta, B. K., Bhardwaj, B. L., and Ahuja, A. K. (2004). Nutritional value of forage crops of Punjab. *Punjab Agricultural University Publication, Ludhiana*.
- Hack, CM., Porta, M., Schäufele, R., and Grimoldi, AA. (2019). Arbuscular mycorrhiza mediated effects on growth, mineral nutrition and biological nitrogen fixation of *Melilotus alba Med.* in a subtropical grassland soil. *Applied Soil Ecology*, 134, 38-44. <https://doi.org/10.1016/j.apsoil.2018.10.008>
- Hanning, GE., and Conger, BV. (1986). Factors influencing somatic embryogenesis from cultured leaf segments of *Dactylis glomerata*. *Journal of plant physiology*, 123(1), 23-29.
- Harika, AS., and Sharma, DD. (1994, February). Quality and yield differences in maize stover due to varieties and stage of harvesting. In *Variation in the Quantity and Quality of Fibrous Crop Residues. Proc. National Seminar held at the BAIF Development Research Foundation* (pp. 20-28).
- Harlan, JR. (1992). Origins and processes of evolution. *Grass Evolution*.
- Harrington, JB. 1952. Cereal breeding procedures. FAO Develop. Paper 28. FAO of UN, Rome.
- Hash, CT., Raj, AB., Lindup, S., Sharma, A., Beniwal, CR., Folkertsma, RT., ... and Blümmel, M. (2003). Opportunities for marker-assisted selection (MAS) to improve the feed quality of crop residues in pearl millet and sorghum. *Field Crops Research*, 84(1-2), 79-88.
- Hou, P., Liu, Y., Liu, W., Liu, G., Xie, R., Wang, K., ... and Li, S. (2020). How to increase maize production without extra nitrogen input. *Resources, Conservation and Recycling*, 160, 104913.

- Humphreys, M., Feuerstein, U., Vandewalle, M., and Baert, J. (2010). Ryegrasses. *Fodder crops and amenity grasses*, 211-260.
- Humphreys, MO. (1997, June). The contribution of conventional plant breeding to forage crop improvement. In *Proceedings of the 18th International Grassland Congress. Winnipeg and Saskatoon, Canada* (pp. 8-17).
- Iqbal, A., Ayub, M., Zaman, H., and Ahmad, R. (2006). Impact of nutrient management and legume association on agro-qualitative traits of maize forage. *Pakistan Journal of Botany*, 38(4), 1079.
- Iqbal, MA., Iqbal, A., Akbar, N., Khan, HZ., and Abbas, RN. (2015). A study on feed stuffs role in enhancing the productivity of milch animals in Pakistan-Existing scenario and future prospect. *Global Veterinaria*, 14(1), 23-33
- Ismaeil, SM., Khafagi, OA., Kishk, ET., and Sohsah, SM. (1993). Effect of some seed hardening treatments on germination, growth and yield of Sudan grass grown under saline conditions. *The Desert Institute Bulletin (Egypt)*.
- Jonker, A., Gruber, MY., McCaslin, M., Wang, Y., Coulman, B., McKinnon, JJ., ... and Yu, P. (2010). Nutrient composition and degradation profiles of anthocyanidin-accumulating L c-alfalfa populations. *Canadian journal of animal science*, 90(3), 401-412.
- Kaur, G., Asthir, B., and Bains, NS. (2018). Modulation of proline metabolism under drought and salt stress conditions in wheat seedlings.
- Kelley, TG., Rao, PP., and Purohit, ML. (1996). Adoption of improved cultivars of pearl millet in an arid environment: straw yield and quality considerations in western Rajasthan. *Experimental Agriculture*, 32(2), 161-171.
- Kumar, A., Arya, RK., Kumar, S., Kumar, D., Kumar, S., and Panchta, R. A. V. I. S. H. (2012). Advances in pearl millet fodder yield and quality improvement through breeding and management practices. *Forage Res*, 38(1), 1-14.
- Kumar, D., Singh, M., Kumar, S., Meena, RK., and Kumar, R. (2021). Fodder quality and nitrate estimation of oats grown under different nutrient management options. *Indian Journal of Dairy Science*, 74(4).
- Liu, SL., Yang, RJ., Ma, MD., Dan, F., Zhao, Y., Jiang, P., and Wang, MH. (2015). Effects of exogenous NO on the growth, mineral nutrient content, antioxidant system, and ATPase

activities of *Trifolium repens* L. plants under cadmium stress. *Acta Physiologiae Plantarum*, 37(1), 1721.

Lowe, KW., and Conger, BV. (1979). Root and Shoot Formation from Callus Cultures of Tall Fescue 1. *Crop Science*, 19(3), 397-400.

Marley, CL., Bruce, DB., Arthaud, J., and Teixeira da Silva, JA. (2017). Fertilizer effects on the quality of grass-clover swards: a review. *Agronomy*, 7(11), 133.

Marshall, AH., Collins, RP., Humphreys, MW., and Scullion, J. (2016). A new emphasis on root traits for perennial grass and legume varieties with environmental and ecological benefits. *Food and Energy Security*, 5(1), 26–39

McCarthy Jr, RD., Klusmeyer, TH., Vicini, JL., Clark, JH., and Nelson, DR. (1989). Effects of source of protein and carbohydrate on ruminal fermentation and passage of nutrients to the small intestine of lactating cows. *Journal of Dairy Science*, 72(8), 2002-2016.

Muir, JP., Pitman, WD., and Foster, JL. (2011). Sustainable, low input, warm season, grass–legume grassland mixtures: Mission (nearly) impossible? *Grass and Forage Science*, 66(3), 301-315.

Nawaz, F., Naeem, M., Ashraf, MY., Tahir, MN., Zulfiqar, B., Salahuddin, M., Shabbir, RN., and Aslam, M. (2016). Selenium supplementation affects physiological and biochemical processes to improve fodder yield and quality of maize (*Zea mays* L.) under water deficit conditions. *Frontiers in Plant Science*, 7, 1438.

Nelson, CJ., and Moser, LE. (1994). Plant factors affecting forage quality. *Forage quality, evaluation, and utilization*, 115-154.

Oram, RN., and Culvenor, RA. (1994). Phalaris improvement in Australia. *New Zealand Journal of Agricultural Research*, 37(3), 329-339.

Osman, KT., and Osman, KT. (2018). Poorly fertile soils. *Management of Soil Problems*, 219-254.

Pandey, AK., Pusuluri, M., and Bhat, BV. (2019). Down-regulation of CYP79A1 gene through antisense approach reduced the cyanogenic glycoside dhurrin in [*Sorghum bicolor* L.) Moench.] to improve fodder quality. *Frontiers in Nutrition*, 6, 122.

- Patel, KI., Ahlawat, RPS., and Trivedi, SJ. (1993). Effect of nitrogen and phosphorus on nitrogen uptake and protein percentage of forage sorghum. *Gujarat Agricultural University Research Journal (India)*.
- Patil, FB., Gadekar, DA., and Bhoite, AG. (1992). Response of forage sorghum varieties to seed rates and nitrogen. *J. Mah. Agric. Univ*, 17, 150-151.
- Poonia, A., Phogat, DS., Nagar, S., Sharma, P., and Kumar, V. (2022). Biochemical assessment of oat genotypes revealed variability in grain quality with nutrition and crop improvement implications. *Food Chemistry*, 377, 131982.
- Poonia, ATMAN., and Phogat, DS. (2017). Genetic divergence in fodder Oat (*Avena sativa* L.) for yield and quality traits. *Forage Research*, 43, 101-105.
- Poppi, DP., Hendricksen, RE., and Minson, DJ. (1985). The relative resistance to escape of leaf and stem particles from the rumen of cattle and sheep. *The Journal of Agricultural Science*, 105(1), 9-14.
- Ramanzin, M., Ørskov, ER., and Tuah, AK. (1986). Rumen degradation of straw 2. Botanical fractions of straw from two barley cultivars. *Animal Science*, 43(2), 271-278.
- Reddy, BVS., Reddy, PS., Bidinger, F., and Blümmel, M. (2003). Crop management factors influencing yield and quality of crop residues. *Field Crops Research*, 84(1-2), 57-77.
- Reid, RL., and Jung, GA. (1974). Effects of elements other than nitrogen on the nutritive value of forage. *Forage fertilization*, 395-435.
- Selinger, LB., Forsberg, CW., and Cheng, KJ. (1996). The rumen: a unique source of enzymes for enhancing livestock production. *Anaerobe*, 2(5), 263-284.
- Shand, WJ., Ørskov, ER., and Morrice, LF. (1988). Rumen degradation of straw 5. Botanical fractions and degradability of different varieties of oat and wheat straws. *Animal Science*, 47(3), 387-392.
- Sharma, A., and Phillips, CJ. (2019). Lameness in sheltered cows and its association with cow and shelter attributes. *Animals*, 9(6), 360.
- Sheaffer, CC., Miller, DW., and Marten, GC. (1990). Grass dominance and mixture yield and quality in perennial grass-alfalfa mixtures. *Journal of Production Agriculture*, 3(4), 480-485.

Skot, L., Humphreys, J., Humphreys, MO., Thorogood, D., Gallagher, J., Sanderson, R., Armstead, I.P. and Thomas, I.D. 2007. Association of candidate genes with flowering time and water-soluble carbohydrate content in *Lolium perenne* (L.). *Genetics* 177:535–547.

Subba Rao, A., Prabhu, UH., and Oosting, SJ. (1993). Genetic and managerial effects on variability of straw quality from ginermillet (*Eleusine coracana*).

Sukanya, DH., V. Ramamurthy and CR. Ramesh 2001: *Forage Res.* 27: 115-118.

Taylor, NL. (2008). A century of clover breeding developments in the United States. *Crop Science*, 48(1), 1-13.

Thomas, SL., and Thomas, UC. (2019). Innovative techniques in fodder production-a review. *Forage Res*, 44, 217-223.

Torello, WA., and Symington, AG. (1984). Regeneration from perennial ryegrass callus tissue. *HortScience*, 19(1), 56-57.

Van Soest, PJ. (2018). *Nutritional ecology of the ruminant*. Cornell university press.

Vasiljević, S., Milić, D., and Mikić, A. (2009). Chemical attributes and quality improvement of forage legumes. *Biotechnology in animal Husbandry*, 25(5-6-1), 493-504.

Veeramani, A., Palchamy, A., Ramasamy, S., and Rangaraju, G. (2000). Integrated weed management in soybean (*Glycine max*) under different moisture regimes and population densities. *Indian Journal of Agronomy*, 45(4), 740-745.

Vogel, KP., and Pedersen, JF. (1993). Breeding systems for cross-pollinated perennial grasses.

Vogel, KP., Haskins, FA., Gorz, HJ., Anderson, BA., and Ward, JK. (1991). Registration of Trailblazer switchgrass.

Waghorn, GC., Shelton, ID., and Thomas, VJ. (1989). Particle breakdown and rumen digestion of fresh ryegrass (*Lolium perenne* L.) and lucerne (*Medicago sativa* L.) fed to cows during a restricted feeding period. *British journal of nutrition*, 61(2), 409-423.

Wahab, K., and Singh, KN. (1983). Effect of irrigation on nutritional uptake and protein content of grain in two types of barley. *Indian journal of agronomy*.

Walker, B., and Weston, EJ. (1990). Pasture development in Queensland—a success story. *Tropical Grasslands*, 24(4), 257-268.

- Walli, TK., Ørskov, ER., and Bhargava, PK. (1988). Rumen degradation of straw 3. Botanical fractions of two rice straw varieties and effects of ammonia treatment. *Animal Science*, 46(3), 347-352.
- Wang, S., Chen, G., Yang, Y., Zeng, Z., Hu, Y., and Zang, H. (2021). Sowing ratio determines forage yields and economic benefits of oat and common vetch intercropping. *Agronomy Journal*, 113(3), 2607-2617.
- Wang, Y., Majak, W., and McAllister, TA. (2012). Frothy bloat in ruminants: cause, occurrence, and mitigation strategies. *Animal feed science and technology*, 172(1-2), 103-114.
- Woodfield, DR., and Brummer, EC. (2001). Integrating molecular techniques to maximise the genetic potential of forage legumes. In *Molecular Breeding of Forage Crops: Proceedings of the 2nd International Symposium, Molecular Breeding of Forage Crops, Lorne and Hamilton, Victoria, Australia, November 19–24, 2000* (pp. 51-65). Springer Netherlands.
- Yadav, RK., and Kumar, A. (1997). Feasibility of cultivating different forage crops on saline soil. *CROP RESEARCH-HISAR*, 13, 45-50.
- Zhang, S., Allen, HL., and Dougherty, PM. (1997). Shoot and foliage growth phenology of loblolly pine trees as affected by nitrogen fertilization. *Canadian Journal of Forest Research*, 27(9), 1420-1426.
- Zhang, Y., and Mian, JR. (2003). Functional genomics in forage and turf-present status and future prospects. *African Journal of Biotechnology*, 2(12), 521-527.
- Zhu, Y., Dong, K., Liu, Z., Wang, J., Wei, S., Shi, Y., and Zheng, Y. (2016). Ergot alkaloids in endophyte-infected grasses are toxic to livestock animals. *Toxins*, 8(5), 144.