

## Original Research Article

“Influence of Seaweed Sap and Zinc on growth and yield of Baby corn (*Zea mays* L.)”

Comment [MF1]: Influence of Seaweed Sap and Zinc on growth and yield of Baby corn (*Zea mays* L.)

### ABSTRACT

A field experiment was conducted during *Khari* 2022 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P) to determine the “Influence of seaweed sap and Zinc on growth and yield of Babycorn (*Zea maize* L.)” The result showed that treatment 9 [ZnSO<sub>4</sub> (20kg/ha) + Seaweed sap (15%)] recorded significantly higher plant height (176.75 cm), higher plant dry weight (g) (88.47), higher crop growth rate (88.27), higher relative growth rate (0.0910), maximum number of cobs/plant (2.82), maximum length of cob/plant (16.90 cm), maximum girth of cob (13.02 cm), maximum cob weight with husk (g) (47.57), maximum cob weight without husk (g) (11.57 g), maximum cob yield with husk (9.5t/ha), maximum green fodder yield (14.18t/ha) compare to other treatments. The maximum gross returns (99,900.00 INR/ha), maximum net return (62,464.00 INR/ha) and highest benefit cost ratio (1.68) was recorded in treatment 9 [ZnSO<sub>4</sub> (20kg/ha) + Seaweed sap (15%)] as compared to other treatments.

Comment [MF2]: Autumn

Comment [MF3]: Sam Higginbottom University of Agriculture, Technology and Sciences

Comment [MF4]: Influence of seaweed sap and Zinc on the growth and yield of Babycorn.

**Keywords:** Babycorn, Economics, Growth, Seaweed sap, Yield and Zinc.

## INTRODUCTION

Baby corn is a vegetable picked from regular maize or sweet corn plants when the ears are still premature and immediately after the emergence of white silk (2–3 cm) length. It is a short season maize variety that can be grown throughout the year. The cobs from a baby corn crop take about 60–65 days to mature, and the rest of the plant can be utilized as green fodder. The young ear is de-husked, eaten raw as a salad and used to make vegetables, pickles, and soup. The nutritional value of baby corn is similar to that of several vegetables, including cauliflower, cabbage, tomato, and radish. The cultivation of the crop is gaining interest in India with increasing production in the states such as Meghalaya, Haryana, Maharashtra, Karnataka, and Andhra Pradesh. Farmers are becoming more interested in baby production due to its low production costs, high domestic demand, promising market, potential value addition, support for the domestic economy and higher revenue **Sale et al. (2022)**

In India, which is the fifth-largest producer in the world and accounts for 3% of worldwide production, maize is grown over an area of roughly 9.18 million hectares, with a yield of 27.23 million tonnes and an average productivity of 2965 kg/ha. With a contribution of 14.87% (1.37 million tonnes) of the total Indian maize produced area, Madhya Pradesh leads the list. In India, Tamil Nadu had the highest output of about 6551 kg/ha while Karnataka produced corn at a rate of about 3.73 million tonnes, or 13.69% of the nation's total production. While Uttar Pradesh gives an area of approximately 0.73 million hectares with a 7.98% to the entire country of India, which has a production of approximately 1.53 million **(Agricultural Statistics at a Glance, 2019)**.

zinc is the most commonly deficient micronutrient in agricultural soils. Analysis of over 256,000 soil samples from all over India showed that about 50% of the soils were deficient in zinc and it causes leaf bronzing and poor tillering at the early growth stages, leading to delayed maturity and significant yield loss, its main cause of deficiency of plant available Zn in soil is the precipitation or adsorption of Zn with various soil components, depending on the pH and redox potential **(Zinzala and Narwade 2019)**. Zinc is an essential micro nutrient for crop plants. Zinc is essential for several biochemical processes in rice plant, such as cytochrome and nucleotide synthesis, auxin metabolism, chlorophyll production, enzyme activation and membrane integrity. Zinc enrichment leads to more root surface area and the ability to change chemistry and biology of rhizosphere by releasing phytosiderophores from roots which ultimately increases Zinc uptake by plants **(Suvarna et al. 2015)**. zinc is major component and activator of several enzymes involved in metabolic activities, its deficiency

Comment [MF5]: corn.Plants

Comment [MF6]: short-season

Comment [MF7]: Cobs

Comment [MF8]: The untutored ear is de-husked, eaten raw as a salad, and used to make vegetables, pickles, and soup.

Comment [MF9]: comparable

Comment [MF10]: being

Comment [MF11]: Zinc

Comment [MF12]: micronutrient of

Comment [MF13]: roots,

Comment [MF14]: Zinc is a major

continues to be one of the key factors in determining rice production in several parts of country (Muthukumararaja and Sriramachandrasekharan 2012). Foliar application of zinc usually applied under emergencies to save the crops to symptoms of Zinc appear, single foliar application may not be adequate to severe deficiency symptoms, foliar application of 0.5% aqueous solution of Zinc twice at 20 and 30 days after transplanting (Verma *et al.* 2019). Heavy application of inorganic fertilizer degrades the soil health and environment, under such situation there is no option left except to depend on the use of available organic and natural fertilizer. Seaweed sap can serve as an important source of organic liquid fertilizer as it contains abundant natural source of major and minor plant nutrients, amino acids, vitamins, as well as growth promoting substances like cytokinins, auxins, and abscisic acid and have been reported to stimulate the growth and yield of crops (Zodape 2009, Crouch and Van-Staden 1993).

*Ascophyllum nodosum* extracts are referred to as biostimulants in most scientific literature as they are typically used to stimulate plant growth. Even at extremely dilute (micromolar) concentrations, *A. nodosum* enhances plant growth, which may be attributed to the variety of plant hormones within this alga. Two of the most prominent hormones within *A. nodosum* are cytokinins and abscisic acid. Cytokinins are thought to improve nutrient uptake efficiency (N, P, K, etc.) by beneficially changing roots to be longer and thinner and can stimulate chlorophyll production. Abscisic acid has novel effects on plants when applied as a biostimulant and can reduce the risk of desiccation by closing leaf stomata. However, its functional effects in *A. nodosum* are unclear. Hormones are not the only beneficial substance in *A. nodosum* extracts. Other growth stimulators from this seaweed, such as various betaines, have been shown to combat common plant infections like bean rust on top of improving the amount of stored chlorophyll. Additionally, a large portion of *A. nodosum* is comprised of phlorotannin polymers which neutralized harmful yeast fungi in an experiment using phlorotannins from other brown algae (Dalzell *et al.* 2019).

Blanket application of nutrients may not be taken by plants properly but, foliar application through plant parts consumes directly by crop. Besides, it is economically cheaper than chemical fertilizer and eco-friendly source of nutrients. In India as a step towards the expansion of native source of natural manures, the seaweed liquid fertilizer application can be utilized as a booster for yield enhancement of crops. (Singhet *et al.* 2015). Bio-stimulant substances extracted from marine algae are used as fertilizer to increase the quality and yield

Comment [MF15]: the country

Comment [MF16]: zinc is usually applied in emergencies

Comment [MF17]: zinc

Comment [MF18]: .

Comment [MF19]: adequate for

Comment [MF20]: zinc

Comment [MF21]: . In such a situation,

Comment [MF22]: under such a situation,

Comment [MF23]: fertilizers.

Comment [MF24]: an abundant

Comment [MF25]: and abscisic acid, and

Comment [MF26]: directly

Comment [MF27]: an eco-friendly

Comment [MF28]: sources

of plants. Seaweed extract is natural organic fertilizers which is highly effective nutrient and promotes maximum yield, quick germination of seeds and ability of resistance of many crops (Dewivediet *al.*2014). The efficacy of the bio-stimulant over 27 different vegetable crops with the application of seaweed *K. alvarezii*. Application of 3–4 foliar applications based on the crop cycle improved the growth, yield, and quality of the produce. An increase of 11% to 52% in the yield levels were noticed for different vegetable crops Karthikeyan and Shanmugam (2016). Keeping in view the abovefact, the experiment was conducted to find out “Influence of Seaweed Sap and Zinc on growth and yield of Baby corn(*Zea maysL.*)”

Comment [MF29]: a natural

Comment [MF30]: fertilizer that is a highly effective

Comment [MF31]: was

Comment [MF32]: Influence of Seaweed Sap and Zinc on the growth and yield of Baby corn

## MATERIALS AND METHODS

The experiment was conducted during ~~Khari season of 2022 at Crop Research Farm, Department of Agronomy~~, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj (U.P). The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, slightly alkaline in soil reaction (pH 8), low level of organic carbon (0.28%), available N (219 Kg/ha), P (11.6 kg/ha) and K (217.2 kg/ha). The experiment was conducted in randomized block design consisting of 10 treatments with 3 different levels of Seaweed sap viz. 5%, 10%, 15% (15 & 30 DAS), three different levels of Zinc viz. 10, 15, 20 (basal application) with 3 replications and the treatments were allocated randomly in each replication. The treatment combinations are Treatment 1. ZnSO<sub>4</sub> 10kg/ha + Seaweed sap 5%, Treatment 2 (ZnSO<sub>4</sub> 10kg/ha + Seaweed sap 10%), Treatment 3 (ZnSO<sub>4</sub> 10kg/ha + Seaweed sap 15%), Treatment 4 (ZnSO<sub>4</sub> 15kg/ha + Seaweed sap 5%), Treatment 5 (ZnSO<sub>4</sub> 15kg/ha + Seaweed sap 10%), Treatment 6 (ZnSO<sub>4</sub> 15kg/ha + Seaweed sap 15%), Treatment 7 (ZnSO<sub>4</sub> 20kg/ha + Seaweed sap 5%), Treatment 8 (ZnSO<sub>4</sub> 20kg/ha + Seaweed sap 10%), Treatment 9 (ZnSO<sub>4</sub> 20kg/ha + Seaweed sap 15%), Treatment 10 (Control (90:60:40 NPK Kg/ha). Growth parameters, yield attributes and economics was recorded. Data recorded on different aspects of crop, such as, growth, yield attributes and yield were subjected to statistical analysis by analysis of variance method (Gomez and Gomez, 1976).

Comment [MF33]: the autumn of 2022 at a crop research farm, department of Agronomy,

Comment [MF34]: Make a table of soil characteristics

Comment [MF35]: Zinc sulfate

Comment [MF36]: Making a table is better for an observation of the ten treatments

Comment [MF37]: Statistical analysis What program was used for the analysis? variance method? least significant difference (LSD)?

## RESULT AND DISCUSSION

### Growth parameters

#### Plant height (cm)

Data revealed that significantly and higher plant height (176.75 cm) was recorded in the treatment 9 [ZnSO<sub>4</sub> (20kg/ha) + Seaweed sap (15%)] (Table 1). Significant and higher plant height was observed with application of Zn might be due to it plays a major role in metabolic

Comment [MF38]: a significantly higher in plant height

Comment [MF39]: treatment

Comment [MF40]: as the application

activity, physiological reaction and act as a catalyzing enzymes transformation of carbohydrates, chlorophyll and protein synthesis. Similar results were also reported by Ammisetty *et al.* (2021). Further, significant and higher plant height was recorded with application of Seaweed sap increases shoot characteristics due to the auxins content in the seaweed sap extracts which have an effective role in cell division and cell enlargement this may have leads to increase shoot growth, leaf area. Similar results were also reported by Bezawada *et al.* (2022).

#### Plant dry weight (g)

The results revealed that significantly higher plant dry weight (88.47g) was recorded in treatment 9 [ZnSO<sub>4</sub> (20kg/ha) + Seaweed sap (15%)]. Significant and higher plant dry weight was with the application of Zinc(20kg/ha) might be due to the involvement of zinc in auxin metabolism, which led to increase higher hormonal activity and growth performance at critical crop growth stages. Similar results were reported by Kumar *et al.* (2013). Further, higher plant dry weight was recorded with application of Seaweed sap (15%) may be due to increase of shoot characteristics due to the auxins content in the seaweed sap extracts which have an effective role in cell division and enlargement this leads to increase shoot growth, leaf area and plant dry weight. Similar results were also reported by Raviteja *et al.*(2022).

#### Crop growth rate (g/m<sup>2</sup>/day)

The data was found significantly higher crop growth rate (88.27) was recorded in treatment 9 [ZnSO<sub>4</sub> (20kg/ha) + Seaweed sap (15%)]. Significant and higher crop growth rate was recorded with the application of Zinc (20kg/ha) might be due to the involvement of zinc in auxin metabolism, which lead to higher hormonal activity and growth performance at critical crop growth stages, resulted higher crop growth rate. Similar results were also described by Kumar *et al.* (2014).

#### Relative growth rate (g/g/day)

Data found that significant and higher relative growth rate (0.0910g/g/day) was recorded in treatment 9 [ZnSO<sub>4</sub> (20kg/ha) + Seaweed sap (15%)]. Significant and higher relative growth rate was recorded with application of seaweed sap (15%) might be due to presence of amino acids like betainies and sterols which enhance the photosynthetic activity, N metabolism and protein synthesis which boost corn production and also availability growth regulators in extract especially Auxin and Cytokinin which are responsible for internodal elongation and cell enlargement and there by increases the relative growth rates. Similar findings also reported by Hegede *et al.* (2016).

Comment [MF41]: and physiological

Comment [MF42]: acts

Comment [MF43]: enzyme

Comment [MF44]: the application

Comment [MF45]: led to an increase

Comment [MF46]: and leaf area.

Comment [MF47]: a significantly

Comment [MF48]: delete

Comment [MF49]: increased

Comment [MF50]: the application

Comment [MF51]: an increase in

Comment [MF52]: , and plant

Comment [MF53]: A significant

Comment [MF54]: leads to

Comment [MF55]: resulting in higher crop growth rates.

Comment [MF56]: a significant

Comment [MF57]: A significant

Comment [MF58]: the application

Comment [MF59]: the presence

Comment [MF60]: thereby

Comment [MF61]: were also

**Table 1. Effect of Seaweed sap and Zinc levelson growth attributes of Baby Corn.**

S. No.	Treatments combinations	60 DAS		45-60 DAS	
		Plant height (cm)	Plant dry weight (g)	Crop growth rate (g/m <sup>2</sup> /day)	Relative growth rate (g/g/day)
1.	ZnSO <sub>4</sub> (10kg/ha) + Seaweed sap (5%)	164.02	73.02	68.38	0.0667
2.	ZnSO <sub>4</sub> (10kg/ha) + Seaweed sap (10%)	162.26	75.09	68.54	0.0639
3.	ZnSO <sub>4</sub> (10kg/ha) + Seaweed sap (15%)	165.02	72.65	62.40	0.0580
4.	ZnSO <sub>4</sub> (15kg/ha) + Seaweed sap (5%)	167.39	74.13	64.46	0.0590
5.	ZnSO <sub>4</sub> (15kg/ha) + Seaweed sap (10%)	166.78	76.11	66.72	0.0597
6.	ZnSO <sub>4</sub> (15kg/ha) + Seaweed sap (15%)	166.26	80.28	69.59	0.0587
7.	ZnSO <sub>4</sub> (20kg/ha) + Seaweed sap (5%)	169.23	83.31	71.50	0.0577
8.	ZnSO <sub>4</sub> (20kg/ha) + Seaweed sap (10%)	174.61	86.95	75.12	0.0583
9.	ZnSO <sub>4</sub> (20kg/ha) + Seaweed sap (15%)	176.75	88.42	97.53	0.0910
10.	Control (120:60:40) NPK kg/ha	162.77	70.79	60.11	0.0568
	F- test	S	S	S	S
	SEm(±)	2.23	0.68	1.40	0.0018
	CD (p = 0.05%)	6.64	2.02	4.17	0.00528

## Yield and Yield parameters

### Number of cobs/plant

Data revealed that significant and maximum number of cobs/ plant (2.82) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. significant and maximum number of cobs/plant was with the application of Zinc (20kg/ha) might be due to higher chlorophyll contents and this had apparently a positive effect of photosynthetic activity synthesis of metabolites and growth regulator substances, oxidation and metabolic activities and ultimately better growth and development of crop which leads to increase in yield attributes namely number of cob/plant of baby corn. Similar findings also reported by Meena *et al.* (2013). Further, significantly increase in number of cobs/plant was with the application of seaweed sap may be due to an effective fertilizer in increasing the growth, biochemical, yield characters of many crop plants and it also improves soil fertility and sustainable yield. Similar findings also reported by Raviteja *et al.* (2022).

Comment [MF62]: a significant

Comment [MF63]: effect on

Comment [MF64]: a significant increase in the number

Comment [MF65]: and yield characteristics

Comment [MF66]: were also

### Length of cob/plant (cm)

Results revealed that significant and maximum length of cob/plant (16.90 cm) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. significant and maximum length of cob/plant was with the application of Zinc (20kg/ha) might be due to continuous filling of grains due to sufficient photosynthesis might have resulted in increased length and size of the cob. Similar findings also reported by Sravan *et al.* (2014).

Comment [MF67]: a significant

Comment [MF68]: The significance

Comment [MF69]: delete

Comment [MF70]: the continuous

Comment [MF71]: were also

### Girth of cob/plant (cm)

Data found that significant and maximum girth of cob (13.02 cm) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. significant and maximum girth of cob was with the application of Zinc (20kg/ha) might be due to higher contents and this had apparently positive effect on photosynthetic activity, synthetic metabolites and growth elongating substances lead to better growth and development. Similar findings also reported by Chand *et al.* (2017).

Comment [MF72]: a significant

Comment [MF73]: , and growth

Comment [MF74]: were also

### Cob weight with husk (g/ha)

Results found that significant and maximum cob weight with husk of baby corn, (47.57 g) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. significant and maximum cob weight with husk was with the application of Zn (20kg/ha) might be due to its role in photosynthesis and metabolic process

augments the production of photosynthates and their translocation to different plant parts. Similar findings also reported by **Karrimiet al. (2017)**.

Comment [MF75]: were also

#### **Cob weight without husk (g/ha)**

Results showed that significant and maximum cob weight with husk of baby corn, (11.57 g) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. Significant and maximum cob weight without husk maximum was with the application of Zn (20kg/ha) might be due to vigorous root development, which promotes growth and development of plant leading to higher photosynthetic activity, which may have results in better development of yield attributes. Similar findings also reported by **Karrimiet al. (2018)**.

Comment [MF76]: the husk

Comment [MF77]: delete

Comment [MF78]: the plant

Comment [MF79]: resulted

Comment [MF80]: were also

Comment [MF81]: **Karrimiet al. (2018)**.

UNDER PEER REVIEW

**Table 2. Effect of Seaweed sap and Zinc levelson yieldattributes of Baby Corn.**

S. No.	Treatments	Number of cobs per plant	Length of cob (cm)	Girth of cob (cm)	Cob weight with husk (gm)	Cob weight without husk (gm)
1.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 5%	2.23	11.25	11.28	41.84	10.21
2.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 10%	2.34	11.74	11.59	43.45	10.68
3.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 15%	2.31	12.42	11.36	44.11	10.78
4.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 5%	2.35	12.01	11.43	43.77	10.79
5.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 10%	2.30	12.38	12.01	44.25	11.23
6.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 15%	2.36	12.43	12.32	45.57	11.56
7.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 5%	2.41	12.70	12.35	45.94	11.52
8.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 10%	2.71	12.72	12.64	47.51	11.76
9.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 15%	2.82	16.90	13.02	47.57	12.01
10.	Control	2.05	11.02	10.81	39.88	10.02
	F-test	S	S	S	S	S
	SEm (±)	0.15	0.34	0.25	0.457	0.18
	CD (p=0.05)	0.44	1.01626	0.74	1.36	0.54

### Cob yield with husk (t/ha)

Data found that significant and maximum cob yield with husk of baby corn, (9.5t/ha) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. Significant and maximum cob yield with husk was with the application of Zn (20kg/ha) might be due to beneficial effect on physiological process, plant metabolism and plant growth, which might have leads to higher cob yield with husk. Similar findings also reported by Chand *et al.* (2017). Further, significant and maximum cob yield with husk was with the application of seaweed sap may be due to acting as an effective fertilizer in increasing the biochemical for growth and yield character of plants, which increased cob yield with husk. Similar findings also reported by Raviteja *et al.* (2022).

Comment [MF82]: the husk

Comment [MF83]: the beneficial

Comment [MF84]: the physiological

Comment [MF85]: led

Comment [MF86]: were also

Comment [MF87]: the significant

Comment [MF88]: delete was

Comment [MF89]: were also

### Green fodder yield (t/ha)

Results revealed that significant and maximum cob weight with husk of baby corn, (14.18t/ha) was observed in treatment 9[(ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%)] which was superior over all other treatments. Significant and maximum green fodder yield was with the application of Zn (20kg/ha) might be due to enhanced translocation of photosynthesis with applied zinc, which resulted in higher production in respective levels of essential nutrients. Similar findings also reported by Chand *et al.* (2017). Further, significant and maximum green fodder was with the application of seaweed sap may be due to saps also enhance nutrient uptake by maize cob and fodder; hence the presence of some macro and micro-elements and plant growth regulators, especially cytokinins, IAA, GA, responsible for increase in yield of baby corn. Similar findings also reported by Pal *et al.* (2015).

Comment [MF90]: revealed

Comment [MF91]: the husk

Comment [MF92]: delete was

Comment [MF93]: were also

Comment [MF94]: delete was

Comment [MF95]: enhancing

Comment [MF96]: an increase

Comment [MF97]: were also

**Table 3. Effect of Seaweed sap and Zinc levelson yield of Baby Corn.**

<b>S. No.</b>	<b>Treatments</b>	<b>Cob yield with husk (t/ha)</b>	<b>Green fodder yield (t/ha)</b>
1.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 5%	7.57	22.37
2.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 10%	7.90	23.49
3.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 15%	8.22	23.22
4.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 5%	8.71	23.37
5.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 10%	8.64	23.64
6.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 15%	8.73	23.73
7.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 5%	9.00	24.00
8.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 10%	9.46	24.12
9.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 15%	9.95	24.61
10.	Control	7.05	20.76
	F-test	S	S
	SEm (±)	0.18	0.19
	CD (p=0.05)	0.53	0.58

### Economics analysis

The result showed that maximum gross returns (99,500.00 INR/ha), Maximum netreturn (62,464.00INR/ha) and highest benefit cost ratio (1.68) was recorded treatment-9 (ZnSO<sub>4</sub> 20kg/ha) + (Seaweed sap 15%). Higher B:C ratio was with application of Zn (20kg/ha) might be due to increase in yield attributes, higher gross return, net returns was found economically viable. Similar results also reported by Chand *et al.* (2022). Further, higher B:C ratio was with application of seaweed sap (15%) may be due to application as a foliar organic nutrient source for enhancing growth and yield of maize, which adds higher gross returns, net returns. Similar results also reported by Basavarajaet *al.* (2018).

Comment [MF98]: , and highest benefit-cost ratio

Comment [MF99]: were

Comment [MF100]: an increase

Comment [MF101]: , and net returns were found economically viable.

Comment [MF102]: were also

Comment [MF103]: B: C ratio with the application

Comment [MF104]: , the higher B: C ratio with the application

Comment [MF105]: its application

Comment [MF106]: the growth and yield of maize,

Comment [MF107]: , and net returns.

Comment [MF108]: were also

UNDER PEER REVIEW

**Table 4. Effect of Zinc and Seaweed sap on Economics of Baby corn.**

S. No.	Treatments	Total cost of cultivation (INR/ha)	Gross return (INR/ha)	Net return (INR/ha)	B:C ratio
1.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 5%	35,036	75700	40664	1.16
2.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 10%	35,536	79000	43464	1.22
3.	ZnSO <sub>4</sub> 10kg/ha + Seaweed sap 15%	36,036	82220	46184	1.28
4.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 5%	35,536	87100	51564	1.45
5.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 10%	36,036	86400	50364	1.39
6.	ZnSO <sub>4</sub> 15kg/ha + Seaweed sap 15%	36,536	87300	50764	1.38
7.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 5%	36,036	90000	53964	1.49
8.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 10%	36,536	94600	58064	1.58
9.	ZnSO <sub>4</sub> 20kg/ha + Seaweed sap 15%	37,036	99500	62464	1.68
10.	Control	33,536	70500	36964	1.10

## CONCLUSION:

Based on the above findings it is concluded that application of ZnSO<sub>4</sub> (20kg/ha) and Seaweed sap (15%) perform better growth parameters and yield attributes of baby corn and also proven profitable.

**Comment [MF109]:** Based on the above findings, it is concluded that the application of ZnSO<sub>4</sub> (20kg/ha) and Seaweed sap (15%), performs better growth parameters and yield attributes of baby corn and has also proven profitable.

## REFERENCES

1. Amutham, G. T., Karthikeyan, R., Thavaprakash, N., and Bharathi, C. (2019). Agronomic bio-fortification with zinc on growth and yield of baby corn under irrigated condition. *Journal of Pharmacognosy and Phytochemistry*, **8**(3):434-437.
2. Aher, A., and Umesha, C. (2023). Effect of Nano Urea and Zinc on Growth and Yield of Baby Corn (*Zeamays L.*) under Prayagraj Condition. *International Journal of Environment and Climate Change*, **13**(6):285-291.
3. Ammisetty Sai Sravan, Dhanush Reddy, Pavan Ganesh and Joy Dawson (2021). Effect of Potassium and Zinc on Growth and Yield of Baby Corn (*Zeamays L.*)
4. Boonchayauy, P., Cakmak, I., Rekasem, B. and ProU-thai, C. (2013). Effect of different foliar zinc application at different growth stages on seed zinc concentration and its impact on seedling vigour in rice. *Soil Science of plant Nutrition* **59**(2): 180-188.
5. Dalzell, C. G. (2019). The effects of a novel, slow-release application method for an Ascophyllum nodosum seaweed extract biofertilizer on maize (*Zeamays L.*) growth and nutrient accumulation.
6. Borase, C. L., Lomte, D. M., Thorat, S. D., & Dhonde, A. S. (2018). Response of Kharif maize (*Zeamays L.*) to micronutrients. *Journal of Pharmacognosy and Phytochemistry*, **7**(3): 482-484.
7. Chand, S. W., Susheela, R., Sreelatha, D., Shanti, M., and Soujanya, T. (2017). Quality studies and yield as influenced by zinc fertilization in baby corn (*Zeamays L.*). *Int. J. Curr. Microbiol. App. Sci.*, **6**(10): 2454-2460.
8. Channakeshava B.C., Ramaprasanna K.P., Ramachandrappa B.K. (2023). Effect of Plant Growth Regulators and Micronutrients on Growth Components and Seed Yield in African Tall Fodder Maize (*Zeamays L.*). *Agricultural Science Digest*. **27**(1).
9. Devi, H. S., and Ghosh, G. (2017). Effect of planting geometry, nitrogen levels and zinc application on growth and yield of hybrid maize (*Zeamays L.*). *Journal of Pharmacognosy and Phytochemistry*, **6**(4): 1067-1069.
10. Devi, N. L., and Mani, S. (2015). Effect of seaweed saps *Kappaphycus alvarezii* and *Gracilaria* on growth, yield and quality of rice. *Indian Journal of Science and Technology*, **8**(19): 1-6.
11. Gumpula, P.K., and Dawson, J. (2022). Effect of Crop Geometry and Seaweed (*Kappaphycus* and *Gracilaria*) Extract on Growth and Yield of Baby Corn (*Zeamays L.*). *International Journal of Plant & Soil Science*, **34**(21): 672-677.
12. Gomez, K.A and Gomez, A.A. (1976). Three more factor experiment in: Statistical procedure for agricultural Research 2nd edition pp:139-141.

13. Hekmat, A. W., Mohammadi, N. K., and Ghosh, G. (2019). Effect of NPK, biofertilizer and zinc foliar nutrition on growth and growth attributes of baby corn (*Zea mays L.*). *IJCS*, **7**(4): 2432-2436.
14. Jerusha, B. M., and Singh, S. (2022). Effect of Nitrogen Levels and Seaweed Extract (*Kappaphycus alvarezii*) on Growth and Yield of Summer Maize (*Zea mays L.*). *International Journal of Plant and Soil Science*, **34**(22): 1313-1321.
15. Krishna, I. V. S. R., and Mayur, B. M. (2022). Effect of spacing and seaweed (*Kappaphycus alvarezii*) extract on yield and economics of sweet corn (*Zea mays L.*).
16. Krishna, T. A., Maharajan, T., and Ceasar, S. A. (2023). The role of membrane transporters in the biofortification of zinc and iron in plants. *Biological Trace Element Research*, **201**(1): 464-478.
17. Keerthi, P., and Singh, S. (2022). Impact of Plant Geometry and Zinc on Growth and Yield Attributes of Sweet Corn (*Zea mays L.*). *International Journal of Plant and Soil Science*, **34**(16): 37-42.
18. Karrimi, A. S., Reddy, A. P. K., Babazoi, F., and Kohistani, T. (2018). Growth, yield and post-harvest soil available nutrients in sweet corn (*Zea mays L.*) as influenced by zinc and iron nutrition. *Journal of Pharmacognosy and Phytochemistry*, **7**(4): 2372-2374.
19. Kumar, R., and Dawson, J. (2019). Growth and yield of sweet corn (*Zea mays L. saccharata*) as influenced by zinc and boron. *Journal of Pharmacognosy and Phytochemistry*, **8**(3): 4332-4335.
20. Roy, S., Singh, V., Singh, S., and Singh, A. C. (2019). Effect of nitrogen and zinc levels on growth, yield and economics of baby corn (*Zea mays L.*). *Journal of Pharmacognosy and Phytochemistry*, **8**(5): 1577-1580.
21. Krishna, P. M., Singh, R., and Indu, T. (2022). Effect of phosphorus and zinc on yield and economics of baby corn (*Zea mays L.*).
22. Kwadzo, K. A., Narkhede, W. N., and Khazi, G. S. (2016). Effect of fertility levels on growth, yield and soil fertility status of maize (*Zea mays L.*) in vertisol of Maharashtra. *Journal of Applied and Natural Science*, **8**(4): 1779-1785.
23. Layek, J., Das, A., Ramkrushna, G. I., Ghosh, A., Panwar, A. S., Krishnappa, R., and Ngachan, S. V. (2016). Effect of seaweed sap on germination, growth and productivity of maize (*Zea mays L.*) in North Eastern Himalayas. *Indian Journal of Agronomy*, **61**(3): 354-359.
24. Mukesh T. Shah, Sudhakar T. Zodape, Doongar Ram Chaudhary, Karuppanan Eswaran & Jitendra Chikara (2013). Seaweed Sap As An Alternative Liquid Fertilizer For Yield And Quality Improvement Of Wheat. *Journal of Plant Nutrition*, **36**(2): 192-200.
25. Pal, A., Dwivedi, S. K., Maurya, P. K., and Kanwar, P. (2015). Effect of seaweed saps on growth, yield, nutrient uptake and economic improvement of maize (sweet corn). *Journal of Applied and Natural Science*, **7**(2): 970-975.
26. Paramesh, V., Dhar, S., Vyas, A. K., and Dass, A. (2014). Studies on impact of phosphorus riched compost, chemical fertilizer and method of zinc application on yield, uptake and quality of maize (*Zea mays L.*). *Indian Journal of Agronomy*, **59**(4): 613-618.
27. Raviteja, B., and Dawson, J. (2022). Effect of Seed Inoculation and Seaweed Extract on Growth and Yield of Baby-corn (*Zea mays L.*). *International Journal of Plant and Soil Science*, **34**(21): 619-625.

28. Rahman, M. T., Jahiruddin, M., Humauan, M. R., Alam, M. J., and Khan, A. A. (2008). Effect of sulphur and zinc on growth, yield and nutrient uptake of boro rice (cv. BRRIDHAN29). *J. Soil. Nature*, **2**(3): 10-15.
29. Rajesh, Sake & Singh, V. and Tiwari, Dhananjay & Pratap, P. and Reddy, Vasu. (2021). Influence of Zinc and Spacing on Growth and Yield of Baby Corn (*Zea mays L.*). *International Journal of Current Microbiology and Applied Sciences*. **10**. 434-438. 10.20546/ijemas.2021.1001.053.
30. Rahul, G. B. (2021). Effect of zinc and spacing on growth and yield of baby corn (*Zeamays L.*).
31. Rohini, G., Singh, V., George, S. G., and Singh, S. K. (2022). Influence of Spacing and Zinc Application on Growth and Productivity of Baby Corn (*Zea mays L.*). *International Journal of Plant & Soil Science*, **34**(17): 124-129.
32. Singh, S. K., Thakur, R., Singh, M. K., Singh, C. S., & Pal, S. K. (2015). Effect of fertilizer level and seaweed sap on productivity and profitability of rice (*Oryza sativa*). *Indian Journal of Agronomy*, **60**(3): 420-425.
33. Satapathy, B. S., Pun, K. B., Singh, T., and Rautaray, S. K. (2014). Effect of liquid seaweed sap on yield and economics of summer rice. *ORYZA-An International Journal on Rice*, **51**(2): 131-135.
34. Sharma, A. J., Singh, M. K., Kumar, S., and Shambhavi, S. (2019). Effect of plant geometry, fertility level and zinc level on kharif baby corn (*Zea mays L.*). *International Journal of Current Microbiology and Applied Sciences*, **8**(07): 1658-67.
35. Sasikanth, Mandadhi Tony, Vikram Singh, and Shruti G. George. (2021). "Effect of biofertilizers and levels of zinc on growth and yield of wheat (*Triticum aestivum L.*)."
36. Sharma, H., and Pyare, R. (2021). To assess the most adoptable hybrid variety of maize (*Zea mays L.*) for different doses of nitrogen management with seaweed extract (*SoliGroGr*). *Journal of Pharmacognosy and Phytochemistry*, **10**(1): 2761-2763.
37. Tharaka, M., Ravi Chandra, K., and Singh, V. (2021). Influence of nitrogen and zinc on growth and yield of baby corn (*Zea mays L.*). *International Journal of Plant and Soil Science*, **33**(23): 64-70.