

Physiological Parameters and Yield of Green Gram As Influenced By Weed Management Practices

ABSTRACT

A field experiment was conducted at Anbil Dharmalingam Agricultural College and Research Institute, Tiruchirappalli during *Summer*, 2021 to evaluate the weed management practices in green gram. The study comprised of eleven weed management practices with different combination *viz.*, pre emergence (PE) application of Diclosulam at 17.5 g/ha, Pendimethalin + Imazethapyr 1 kg/ha, post emergence (POE) application of Fluazifop-p-butyl + Fomesafen 313 g/ha, Acifluorfen sodium + Clodinofof propargyl 187.5 g/ha, PE Diclosulam 17.5 g/ha *fb* POE Fluazifop-p-butyl + Fomesafen 313 g/ha, PE Pendimethalin + Imazethapyr 1 kg/ha *fb* POE Fluazifop-p-butyl + Fomesafen 313 g/ha, PE Diclosulam 17.5 g/ha *fb* POE Acifluorfen sodium + Clodinofof propargyl 187.5 g/ha, PE Pendimethalin + Imazethapyr 1 kg/ha *fb* POE Acifluorfen sodium + Clodinofof propargyl 187.5 g/ha, PE Pendimethalin 1 kg/ha *fb* HW on 30 DAS, HW on 20 and 40 DAS and unweeded control (UWC) in green gram. The results revealed that the highest plant height, number of branches plant⁻¹, DMP, LAI, CGR, RGR and NAR were registered in PE Pendimethalin + Imazethapyr at 1 kg ha⁻¹ *fb* POE Fluazifop-p-butyl + Fomesafen at 313 g ha⁻¹. The grain and haulm yields were also higher in PE Pendimethalin + Imazethapyr at 1 kg ha⁻¹ *fb* POE Fluazifop-p-butyl + Fomesafen at 313 g ha⁻¹ and it was on par with PE Pendimethalin + Imazethapyr at 1 kg ha⁻¹ *fb* POE Acifluorfen sodium + Clodinofof propargyl at 187.5 g ha⁻¹. Hence, application of PE Pendimethalin + Imazethapyr at 1 kg ha⁻¹ *fb* POE Fluazifop-p-butyl + Fomesafen at 313 g ha⁻¹ was found to be the most viable option for getting higher growth and yield in green gram.

Keywords: Green gram; weed management; Growth; Physiological parameters; Yield.

INTRODUCTION

Pulses are rich source of dietary proteins and inseparable ingredients of vegetarian diet. India contributes about 25 per cent of global pulse production. India is the first largest producer of pulses in the world [1]. Green gram (*Vigna radiata L.*) is one of the most important pulse crop. Among the grain legumes, in India green gram stands third position after chickpea and pigeon pea [2]. In India, green gram is grown in an area of 4.75 m.ha with a production of 2.45 mt and an average productivity of 516 kg ha⁻¹. In Tamil Nadu, green gram is the major traditional pulse crop cultivated under 1.70 L.ha with production of 0.76 L.t and an average productivity of 451 kg ha⁻¹[3].

Weeds are the major yield limiting biotic constraint in green gram. The yield loss due to severe weed infestation varies from 30-80 per cent in green gram [4]. The control of weeds during critical period of crop weed competition is very important so as to avoid yield loss [5]. Improper weed control results in greater yield loss [6]. So there is a need to control weeds in order to reduce the yield loss in green gram. Therefore, effective weed management practices are more important for green gram cultivation.

Hand weeding is difficult due to non-availability of labour as well as high cost of weeding [5]. Under these circumstances, one of the best choice for controlling weeds is through application of herbicides at appropriate time. Herbicides control the weeds effectively and is economically feasible one [7]. Herbicides are controlling weeds in very broad spectrum with appropriate application [8]. Hence, the present investigation has been carried out to evaluate the different weed management practices on the performance of green gram under sodic soil.

MATERIALS AND METHODS

A field experiment was conducted at Anbil Dharmalingam Agricultural College and Research Institute, Tiruchirappalli, Tamil Nadu during *Summer*, 2021. The experimental field was located at 10° 45' N latitude, 78 36' E longitude and at an altitude of 85 m above MSL. The experimental soil was low in available nitrogen, medium in available phosphorus and high in available potassium.

The field experiment was laid out in randomized block design (RBD) with three replications and eleven treatments. The treatments comprised of pre emergence (PE) application of Diclosulam at 17.5 g/ha, Pendimethalin + Imazethapyr 1 kg/ha, post emergence (POE) application of Fluazifop-p-butyl + Fomesafen 313 g/ha, Acifluorfen sodium + Clodinofof propargyl 187.5 g/ha, PE Diclosulam 17.5 g/ha *fb* POE Fluazifop-p-butyl + Fomesafen 313 g/ha, PE Pendimethalin + Imazethapyr 1 kg/ha *fb* POE Fluazifop-p-butyl + Fomesafen 313 g/ha, PE Diclosulam 17.5 g/ha *fb* POE Acifluorfen sodium + Clodinofof propargyl 187.5 g/ha, PE Pendimethalin + Imazethapyr 1 kg/ha *fb* POE Acifluorfen sodium + Clodinofof propargyl 187.5 g/ha, PE Pendimethalin 1 kg/ha *fb* HW on 30 DAS, HW on 20 and 40 DAS and unweeded control (UWC). The variety used for the experiment was VBN (Gg) 4.

Observations recorded during the course of investigation were plant height (cm), number of branches plant⁻¹, dry matter production (kg/ha), leaf area index (LAI), crop growth rate (CGR) (g/m² /day), relative growth rate (RGR) (g/g/day) and net assimilation rate (NAR) (g/cm² /day) by adopting standard procedure. The grain and haulm yields were recorded and expressed in kg/ha. All the recorded data were analyzed statistically as per the method suggested by Gomez and Gomez [9].

RESULTS AND DISCUSSION

Growth parameters

Adoption of different weed management practices improved the growth parameters *viz.*, plant height, number of branches plant⁻¹ and dry matter production of green gram (Table 1). The tallest plant height (12.1, 24.5 and 46.8 cm), higher number of branches plant⁻¹ (4.3/plant) and dry matter production (1066, 1703 and 2700 kg/ha) at 30 DAS, 45 DAS and harvest stages were recorded in PE Pendimethalin + Imazethapyr 1 kg/ha *fb* POE Fluazifop-p-butyl + Fomesafen 313 g/ha and it was comparable with PE application of Pendimethalin + Imazethapyr at 1 kg ha⁻¹ *fb* POE Acifluorfen sodium + Clodinofof propargyl at 187.5 g ha⁻¹. This was due to better weed control by the herbicides and increased WCE which result in well-developed foliage canopy and better light interception on crops. This is in conformity with the findings of Ramesh and Rathika [5].

Physiological parameters

The leaf area index (LAI) of green gram promotes photosynthate translocation to meet sink requirements, making it directly tied to grain yield. The highest LAI (1.14, 3.13 and 2.82 at 30 DAS, 45 DAS and harvest stages) was registered with PE application of Pendimethalin + Imazethapyr at 1 kg ha⁻¹ fb POE Fluazifop-p-butyl + Fomesafen at 313 g ha⁻¹ and it was on par with PE Pendimethalin + Imazethapyr at 1 kg ha⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl at 187.5 g ha⁻¹ (Table 2). The effective control of weeds in these treatments might have favoured the crop growth through better availability of resources. This is in conformity with the findings of Shruthi and Salakinkop [10] and Shanmugapriya et al. [11]. UWC registered lower LAI due to severe weed infestation.

Among all the weed management practices, application of PE Pendimethalin + Imazethapyr at 1 kg ha⁻¹ fb POE Fluazifop-p-butyl + Fomesafen at 313 g ha⁻¹ was registered higher CGR (4.35 and 4.09 g m⁻² day⁻¹), RGR (0.075 and 0.019 g g⁻¹ day⁻¹) and NAR (0.275 and 0.140 mg cm⁻² day⁻¹) at 30-45 DAS and 45-harvest stage, respectively) and it was followed by PE application of Pendimethalin + Imazethapyr at 1 kg ha⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl at 187.5 g ha⁻¹ at 30-45 DAS and 45-harvest (Table 2). Sequential application of pre emergence followed by post emergence herbicides was effectively controlled the grasses, sedges and broad leaved weeds and thereby enhanced the CGR, RGR and NAR as reported by Srijani Maji et al. [12].

Table 1. Effect of weed management practices on plant height, no. of branches plant⁻¹ and dry matter production (DMP) in irrigated green gram

Treatments	Plant height (cm)			No. of branches plant ⁻¹	DMP (kg ha ⁻¹)		
	30 DAS	45 DAS	Harvest		30 DAS	45 DAS	Harvest
T ₁ - PE Diclosulam @ 17.5 g ha ⁻¹	8.8	16.2	29.7	2.16	809	1300	1887
T ₂ - PE Pendimethalin + Imazethapyr @ 1 kg ha ⁻¹	9.1	17.1	33.6	2.37	868	1406	2256
T ₃ - POE Fluazifop-p-butyl + Fomesafen @ 313 g ha ⁻¹	8.9	16.9	32.9	2.29	852	1388	2125
T ₄ - POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha ⁻¹	8.7	16.7	31.0	2.21	839	1362	1974
T ₅ - PE Diclosulam @ 17.5 g ha ⁻¹ fb POE Fluazifop-p-butyl + Fomesafen @ 313 g ha ⁻¹	10.8	19.3	38.5	2.64	947	1489	2378
T ₆ - PE Pendimethalin + Imazethapyr @ 1 kg ha ⁻¹ fb POE Fluazifop-p-butyl + Fomesafen @ 313 g ha ⁻¹	12.1	24.5	46.8	4.35	1066	1703	2700
T ₇ - PE Diclosulam @ 17.5 g ha ⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha ⁻¹	10.3	18.9	37.4	2.42	933	1472	2339
T ₈ - PE Pendimethalin + Imazethapyr @ 1 kg ha ⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha ⁻¹	11.9	23.8	44.2	3.73	1043	1677	2655
T ₉ - PE Pendimethalin @ 1 kg ha ⁻¹ fb HW on 30 DAS	10.9	21.3	42.5	2.97	965	1550	2453
T ₁₀ - HW at 20 and 40 DAS	11.0	22.2	43.3	3.32	1001	1631	2576
T ₁₁ - Unweeded Control (UWC)	7.9	15.7	28.0	1.45	722	988	1305
SEd	0.5	0.9	1.6	0.13	41	63	99
CD (P=0.05)	0.9	1.7	3.2	0.26	82	126	199

Table 2. Effect of weed management practices on leaf area index (LAI), crop growth rate (CGR), relative growth rate (RGR) and net assimilation rate (NAR) in irrigated green gram

Treatments	LAI			CGR (g m ⁻² day ⁻¹)		RGR (g g ⁻¹ day ⁻¹)		NAR (mg cm ⁻² day ⁻¹)	
	30 DAS	45 DAS	Harvest	30-45 DAS	45 - Harvest	30-45 DAS	45 - Harvest	30-45 DAS	45 - Harvest
T ₁ - PE Diclosulam @ 17.5 g ha ⁻¹	0.52	1.97	1.03	3.35	2.40	0.03	0.015	0.221	0.107
T ₂ - PE Pendimethalin + Imazethapyr @ 1 kg ha ⁻¹	0.71	2.39	1.78	3.67	3.48	0.03	0.018	0.232	0.136
T ₃ - POE Fluazifop-p-butyl + Fomesafen @ 313 g ha ⁻¹	0.68	2.12	1.71	3.66	3.02	0.03	0.017	0.232	0.125
T ₄ - POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha ⁻¹	0.66	2.00	1.62	3.57	2.51	0.03	0.015	0.225	0.108
T ₅ - PE Diclosulam @ 17.5 g ha ⁻¹ fb POE Fluazifop-p-butyl + Fomesafen @ 313 g ha ⁻¹	0.88	2.82	2.16	3.70	3.64	0.03	0.018	0.241	0.138
T ₆ - PE Pendimethalin + Imazethapyr @ 1 kg ha ⁻¹ fb POE Fluazifop-p-butyl + Fomesafen @ 313 g ha ⁻¹	1.14	3.13	2.82	4.35	4.09	0.07	0.019	0.275	0.140
T ₇ - PE Diclosulam @ 17.5 g ha ⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha ⁻¹	0.82	2.74	2.03	3.68	3.55	0.03	0.018	0.241	0.137
T ₈ - PE Pendimethalin + Imazethapyr @ 1 kg ha ⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha ⁻¹	1.11	3.03	2.70	4.33	4.01	0.03	0.019	0.272	0.139
T ₉ - PE Pendimethalin @ 1 kg ha ⁻¹ fb HW on 30 DAS	0.96	2.68	2.43	4.00	3.70	0.03	0.018	0.251	0.138
T ₁₀ - HW at 20 and 40 DAS	1.01	2.84	2.56	4.30	3.78	0.03	0.018	0.256	0.139
T ₁₁ - Unweeded Control (UWC)	0.27	1.34	0.86	1.81	1.28	0.02	0.011	0.217	0.118
SEd	0.04	0.11	0.09	0.16	0.14	0.00	0.0005	0.01	0.005
CD (P=0.05)	0.07	0.21	0.18	0.32	0.27	0.00	0.0010	0.02	0.011

Grain and haulm yield

Grain and haulm yields of green gram varied significantly with different weed management practices. PE application of Pendimethalin + Imazethapyr @ 1 kg ha⁻¹ fb POE Fluazifop-p-butyl + Fomesafen @ 313 g ha⁻¹ recorded significantly higher grain and haulm yields (708 kg ha⁻¹ and 1602 kg ha⁻¹) than other treatments. However, it was comparable with PE Pendimethalin + Imazethapyr @ 1 kg ha⁻¹ fb POE Acifluorfen sodium + Clodinofof propargyl @ 187.5 g ha⁻¹ with grain and haulm yields of 673 kg ha⁻¹ and 1531 kg ha⁻¹ (Fig. 1). This might be due to the cumulative effect of increased level of yield attributes which was due to lesser weed competition at critical stages, better uptake of nutrients and good crop stand. These results were in conformity with the findings of Ramesh and Rathika [13].

The UWC registered significantly lower grain and haulm yield (281 kg ha⁻¹ and 745 kg ha⁻¹). This could be due to severe weed infestation creates crop weed competition, results in crop plants unable to express their genetic potential. This corroborates with findings of Sudesh Kumar et al. [14].




Fig 1. Grain and haulm yields (kg ha⁻¹) of green gram as influenced by weed management practices

CONCLUSION

From this field experiment, it was concluded that PE application of Pendimethalin + Imazethapyr @ 1 kg ha⁻¹ fb POE Fluazifop-p-butyl + Fomesafen @ 313 g ha⁻¹ was found to be the best weed management practices to attain increased growth and yield of irrigated green gram under sodic soil.

REFERENCES

1. Rathika S, Udhaya A, Ramesh T, Shanmugapriya P. Weed management strategies in green gram: A review. *The Pharma Innovation Journal*. 2023;12(3): 5574-5580.
2. IIPR; 2021. Available from: <http://iipr.icar.gov.in>.
3. Indiastat; 2019. Available from: www.indiastat.com.
4. Algotar SG, Raj VC, Pate DD, Patel DK. Integrated weed management in green gram. *In: 25th Asian-Pacific Weed Science Society Conference on "Weed Science for Sustainable Agriculture"*, India, Hyderabad, 2015; October 13-16.
5. Ramesh T, Rathika S. Management of emerged weeds in irrigated blackgram (*Vigna mungo* L.) through post emergence herbicides. *Legume Research*. 2016;39(2):289- 292.
6. Ramesh T, Rathika S, Parthipan T, Ravi V. Productivity enhancement in black gram through refinement of nutrient management under rice fallow condition. *Legume Research*. 2016;39(1):106-109
7. Muoni, Tarirai, Leonard Rusinamhodzi, Christian Thierfelder. Weed control in conservation agriculture systems of Zimbabwe: identifying economical best strategies. *Crop Protection*. 2013;53:23-28.
8. Ramesh T, Rathika S. Weed management in rice fallow black gram through post-emergence herbicides. *Madras Agricultural Journal*. 2015;102(10-12):313-316.
9. Gomez, KA and Gomez, AA. *Statistical Procedure for Agricultural Research*. John Wiley and Sons, New York; 1984.
10. Shruthi GK, Salakinkop SR. Efficacy of sequential application of pre and post-emergence herbicides in *Kharif* green gram (*Vigna radiata* L.). *Karnataka Journal of Agricultural Sciences*. 2015;28(2):155-159.
11. Shanmugapriya P, Rathika S, Ramesh T, Janaki P. Response of transplanted finger millet to weed management practices under sodic soil condition. *Journal of Applied and Natural Science*. 2022;14(SI): 263 - 268.
12. Srijani Maji, Hasim Reja MD, Rajib Nath, Pintoo Bandopadhyay, Pushpendu Dutta. Herbicidal management in monsoon green gram (*Vigna radiata* (L.) Wilczek) and its effect on the following rapeseed (*Brassica campestris* L. var. Yellow Sarson) in the Indo-Gangetic plains of Eastern India. *Journal of the Saudi Society of Agricultural Sciences*. 2020;19(8):499-509.
13. Ramesh T, Rathika S. Management of *Trianthema portulacastrum* through herbicides in green gram. *Indian Journal of Weed Science*. 2020;52(3):286-289.
14. Sudesh Kumar, Gupta KC, Rani Saxena, Yadav M.R, Bhadhoria S.S. Efficacy of herbicides on weed management in green gram (*Vigna radiata* L.) in semiarid eastern plain zone of Rajasthan. *Annals of Plant and Soil Research*. 2019;21(1):14-18.