

Comparative Analysis of the Impact of Crude Oil and Kerosene on Soil Microbiota and the Bio-utilization Potentials of the Indigenous Microorganisms

ABSTRACT

This research is focused on evaluating the impact of crude oil and kerosene on soil microbiota. The enumeration of total heterotrophic bacterial count revealed that the soil impacted with crude oil and kerosene yielded low microbial counts ranged from 3.2×10^6 CFU/g, 3.2×10^5 to 2.6×10^3 CFU/g while that of total coliform count obtained from the samples ranges from 2.6×10^5 CFU/g to 2.2×10^3 CFU/g. and total fungal count ranged from 2.0×10^2 CFU/g to 1.4×10^2 CFU/g. The genera of bacteria isolated from soils impacted with crude oil and kerosene belong to the genera: *Pseudomonas* spp, *Bacillus* spp, *Staphylococcus* spp, *Streptococcus* spp, *Escherichia coli*, *Micrococcus* spp, *Klebsiella* spp, *Corynebacterium* and *Salmonella* species while *Aspergillus* spp, *Alternaria* spp, *Candida* spp, *Fusarium* spp, *Trichoderma* spp, *Mucor* spp, *Penicillium* spp, *Cephalosporium* spp, *Rhizopus* spp are the isolated fungi. The percentage occurrence revealed that *Pseudomonas* spp and *Escherichia coli* had the highest prevalence of 23.1%, followed by *Staphylococcus* spp (7.7%), *Salmonella* spp (7.7%), *Bacillus* spp (7.7%), *Micrococcus* spp (7.7%) etc. While the fungal isolates had the following prevalence; *Aspergillus* spp (12.5%), *Penicillium* spp (12.5%), *Candida* spp (12.5%), *Mucor* spp (12.5%), etc. The results obtained indicates that crude oil and kerosene have a negative impact on the microbiota of the soil. The combined effect of the compounds is more harmful than their individual effects. This result implies that farmlands impacted with these products would not be suitable for agricultural purposes since the nitrogen fixing bacteria and other soil enriching microorganisms must have been either killed or inactivated by the toxicity of compounds.

Key words: Crude oil and Kerosene, comparative analysis, bioutilization, prevalence, soil microbiota.

INTRODUCTION

Environmental pollution has been on the rise in the past few decades owing to increased human activities on energy reservoirs, unsafe agricultural practices and rapid industrialization [9]. Amongst the pollutants that are of environmental and public health concerns due to their toxicities are: heavy metals, nuclear wastes, pesticides, greenhouse gases, crude oil and refined petroleum products like kerosene. Environmental pollution associated with petroleum hydrocarbons is one of the world's most common environmental problems [3; 27]. Crude oil spillage is one of the most serious environmental problems currently facing the oil producing areas and occurs in large scale in some communities. The oil spillage could be attributed to different causes such as accidental spills, leakage, and vandalization of pipelines and corrosion of pipelines which allow the seepage of crude oil into the environment [26]. The effect of oil spillage on land has become a global issue as land play an important role in the sustenance of man [1]. When land is contaminated, the contaminants change the chemical and biological properties of the soil and are toxic to some soil microorganisms [10; 27; 24]. The chemical composition of crude oil and kerosene varies significantly and can have diverse effects on different organisms within the ecosystem and these differences are due to variation in concentration levels of the various constituents The contamination changes the physiochemical and biological properties of the soil because the oil may be toxic to some soil microorganisms and plants [14].

Contamination of soil by crude oil could lead to reduced microbial density and activities. Soil conditions of agricultural land, microorganisms as well as plants are damaged or altered by any contact with crude oil [19]. Excess oil in soil limits the availability of nitrogen [11]. Soils that are polluted with petroleum hydrocarbons (PHCs) and associated products are different from unpolluted soils due to changes in their biological as well as physicochemical properties. Petroleum hydrocarbon may interfere with the plant-fungus relationship by altering the soil environment so that movement of diffusible chemical signals such as auxins is prevented. Soil biological activity, including soil microbial biomass, is influenced by a range of physiochemical, environmental parameters and perturbations. Therefore, soil microbial activity may be used to assess disturbed soil [12]. Biologically and biochemically mediated processes in soils are of utmost importance to ecosystem functions. There is a huge diversity of organisms belonging to different taxonomic and physiologic groups that interact at different levels within the community in soil biota [13; 7; 8].

In this biota, soil microorganisms constitute a source and are the driving force behind many soil processes, including the transformation of organic matter, nutrient release, transformation of carbon, nitrogen, phosphorus and sulphur, degradation of xenobiotic compounds, the formation of soil physical structure and enhanced nutrient uptake by plants [13]. They degrade organic pollutants by using them as their carbon and energy source. And more than 200 species of bacteria, fungi, and even algae are capable of degrading hydrocarbons because of their ubiquitous nature. Onwurah et al. [20], reported that *Pseudomonas*, *Micrococcus* and *Bacillus* can metabolize the toxic components of crude oil, leading to degradation. Crude oil and refined petroleum products pollute our farmlands and aquatic environment thereby hindering the production of crops growth and aquatic lives and may negatively affect the activities of soil microorganisms responsible for the good crop yield and soil fertility. Crude oil and refined petroleum products may also affect the agricultural productivity of ecosystems by causing dislocation in the biogeochemical cycle where microorganisms participate. Due to the need for sufficient information in this area, this study was designed to analyze the comparative toxicity of crude oil and kerosene on soil microbiota.

MATERIALS AND METHODS

Sample collection

Soil samples were collected with the aid of a sterilized auger from Cross River University of Technology Campus, Calabar agricultural farmland. The Bonny light crude oil was collected from EPZ depot, Calabar, Cross River State, and kerosene was collected from Nigeria National Petroleum Corporation (NNPC) station, Calabar, Cross River State in a sterilized McCartney bottle. All the samples collected were taken to Microbiology laboratory for further investigation.

Microbiological Analysis

Enumeration of Total Heterotrophic Bacterial (THB) counts

The total heterotrophic bacterial loads of soil samples were determined using pour plate technique as described by Cheesbrough [4]. Serial dilution (10^1 - 10^{10}) was done, from the soil samples. Exactly one millimeter (1ml) was taken from each selected dilution (10^2 , 10^4 , 10^6) into sterile petri dishes molten nutrient agar at 37°C for 24 hours. Thereafter, plates with colony growth were counted and expressed in colony forming unit.

Determination Total Coliform counts

The total coliform bacterial loads of soil samples were determined using pour plate technique as described by Cheesbrough [4]. Serial dilution (10^1 - 10^{10}) was done, from the soil samples. Exactly one millimeter (1ml) was taken from each selected dilution (10^2 , 10^4 , 10^6) into sterile petri dishes with molten MacConkey agar at 37°C for 24 hours. Thereafter, plates with colony growth were counted and expressed in colony forming unit.

Screening for Hydrocarbon Utilizing Bacteria

The hydrocarbon utilizing potential of sample was carried out under aerobic conditions following the method adopted by Chikere and Ekwuabu [5]. A loop full of 24 hours old culture of each hydrocarbon utilizing bacteria were inoculated into BH (Bushnell-Hass) broth containing 1% (v/v) crude oil. Biodegradation was recorded with the discolouration of Dichlorophenolindophenol (DCPIP) oxidation reduction reagent after 14 days of incubation at 30°C.

Biochemical characterization and identification of bacterial isolates

Pure isolates from the corresponding agar slants were characterized and identified using morphological, biochemical and physiological characterization as described by Cheesbrough [4].

Determination total fungal count

The total fungal loads of soil samples were determined using spread plate technique. Serial dilution (10^1 - 10^{10}) was done, from the soil samples. Exactly one millimeter (1ml) was taken from each selected dilution (10^2 , 10^4 , 10^6) into sterile petri dishes with molten Potato dextrose agar at 37°C for 72 hours. Thereafter, plates with colony growth were counted and expressed in colony forming unit.

Identification of fungal isolates

The colonial morphologies of the fungal isolates on Sabouraud Dextrose Agar were observed for colour and type of growth following microscopic identification as described by Murray *et al.* [16]. This was done by preparing a wet mount using lactophenol cotton blue to observe the microscopic characteristics of the fungi such as type of hyphae (whether septate or non-septate). A drop of Lactophenol blue was placed on a clean microscope slide, with the aid of an inoculating needle, a small portion of growth midway between the colony center and edge was gently removed and placed on the dropped Lactophenol blue on the slide. With two sterile dissecting needles, the fungus was gently teased apart so that it is thinly spread out in the Lactophenol. After which a coverslip was placed on the edge of the Lactophenol and slowly lowered. Then placed under the microscope for examination. Both microscopic and macroscopic features of the fungal isolates were matched based on the mycological atlas for fungal identification [16].

Screening for Hydrocarbon Utilizing Fungi

In order to screen for hydrocarbon utilizing fungi, the fungal isolates were transferred into plates of Rose-Bengal Chloramphenicol (RBC) agar and Bushnell-Hass (mineral salt) agar supplemented with 0.05% (v/v) streptomycin. Sterile Whatman filter papers were soaked in medium containing crude oil and were aseptically placed into the lids of inoculated Bushnell-Haas agar plates; this technique is called the vapour phase transfer [6]. After the inoculation procedures, RBC agar plates and Bushnell-Haas agar were incubated at 30°C for 7 days and 14 days respectively and isolates counted and recorded.

RESULTS

The enumeration of the samples analyzed had different microbial loads. The total heterotrophic bacterial count of the samples from different locations revealed that the crude oil and kerosene has effect on the soil microbiota as the soil samples without crude oil and kerosene yielded high bacterial count (3.7×10^9 CFU/ml) when compared to soil samples impacted with crude oil and kerosene (2.6×10^3 CFU/ml) as shown in Table 1.

Table 1: Total heterotrophic bacterial (THB) count of soil samples impacted with Crude oil and kerosene respectively and soil samples without Crude oil and kerosene (control)

SAMPLE CODE	SAMPLES	THB COUNTS (CFU/ml)
SC	Soil + Crude oil	3.2×10^5
SK	Soil + Kerosine	3.0×10^6
SCK	Soil + crude oil+ Kerosine	2.6×10^3
SS	Soil (control)	3.7×10^9

The total coliform count of the samples showed that the soil samples impacted with crude oil and kerosene yielded low coliform count (2.4×10^3 CFU/ml) when compared to the control (3.2×10^7 CFU/ml) as displayed in Table 2.

Table 2: Total coliform (TC) count of soil samples impacted with Crude oil and kerosene respectively and soil samples without Crude oil and kerosene (control)

SAMPLE CODE	SAMPLES	TC COUNTS (CFU/ml)
SC	Soil + Crude oil	2.2×10^5
SK	Soil + Kerosine	2.6×10^5
SCK	Soil + crude oil+ Kerosine	2.4×10^3
SS	Soil (control)	3.2×10^7

The fugal screening also revealed that the Crude oil and kerosene had a negative impact on fugal loads as the soil without Crude oil and kerosene had higher fungal load (2.2×10^4 CFU/ml) compared to the soil samples impacted with Crude oil and kerosene (1.4×10^2 CFU/ml) as seen in Table 3.

Table 3. Total Fungal (TF) count of soil samples impacted with Crude oil and kerosene respectively and soil samples without Crude oil and kerosene (control)

SAMPLE CODE	SAMPLES	TF COUNT (CFU/ml)
-------------	---------	-------------------

SC	Soil + Crude oil	2.0×10^2
SK	Soil + Kerosine	1.8×10^2
SCK	Soil + crude oil+ Kerosine	1.4×10^2
SS	Soil (control)	2.2×10^4

Biochemical characterization and identification processes revealed as shown in Table 4 that *Salmonella* species, *Micrococcus* species, *Bacillus* species, *Staphylococcus* species, *Escherichia coli*, *Klebsiella* species etc. were the suspected bacterial isolates observed in the samples.

Table 4: Biochemical characterization and identification of bacterial isolates

Morphological Characteristics	Cell shape	Suspected organisms														
		Gram reaction	Oxidase	Catalase	Citrate	Indole	Methyl red	Voges Proskauer	Glucose	Lactose	Sucrose	TSI slant	TSI butt	Gas	H ₂ S	
Circular, pinkish	Cocci in clusters	-	+	+	-	-	+	-	-	-	+	Y	Y	+	-	<i>Staphylococcus</i> species
Creamy, circular, rough	Long rods	+	+	+	-	-	-	-	-	-	-	R	Y	+	-	<i>Bacillus</i> species
Circular, transparent, Opaque	Short rods	-	+	+	-	+	+	+	-	-	+	R	Y	+	+	<i>Corynebacterium</i> species
Spreading, colorless, rough	Rods in chains	-	-	+	+	-	-	-	+	-	+	Y	Y	+	-	<i>Pseudomonas</i> species
Colorless, rod like	Tiny rods	-	+	+	+	-	-	-	-	+	+	Y	Y	+	-	<i>Klebsiella</i> species
Creamy, long, smooth	Rods in single	-	-	+	-	+	+	-	-	+	-	Y	Y	-	-	<i>Escherichia coli</i>
Circular, creamy, curve-shaped	Cocci	+	+	+	-	+	-	-	-	+	-	R	Y	-	-	<i>Micrococcus</i> species
Convex, curve-shaped	Rods	-	+	+	+	-	+	+	+	+	-	Y	R	-	+	<i>Salmonella</i> species
Creamy, long, smooth	Rods	-	+	+	-	+	+	-	-	+	-	Y	Y	-	-	<i>Escherichia coli</i>
Circular, raised, rough	Cocci	-	-	+	-	+	+	-	-	+	-	Y	Y	+	-	<i>Streptococcus</i> species
Spreading, rough, colorless	Rods in chains	-	+	+	+	-	-	-	+	-	+	Y	Y	+	-	<i>Pseudomonas</i> species

Creamy, long, smooth	Rods	-	+	-	+	+	+	-	-	+	-	Y	Y	-	-	<i>Escherichia coli</i>
Spreading, rough, colorless	Rods in chains	-	+	+	+	-	+	-	-	-	+	Y	Y	+	-	<i>Pseudomonas species</i>

The degradative potentials of hydrocarbons by bacterial isolates are presented in Table 5.

Table 5 Hydrocarbon Utilization Bacteria

S/N	ORGANISMS	DEDRADATIVE SCREENING
1	<i>Pseudomonas spp</i>	+
2	<i>Streptococcus spp</i>	-
3	<i>Salmonella spp</i>	-
4	<i>Bacillus spp</i>	-
5	<i>Escherichia coli</i>	+
6	<i>Micrococcus spp</i>	-
7	<i>Klebsiella spp</i>	+
8	<i>Staphylococcus spp</i>	-
9	<i>Corynebacterium spp</i>	-

The prevalence of the bacterial isolates revealed that *Pseudomonas species* and *Escherichia coli* had the highest number of occurrences followed by *Streptococcus specie*, *Bacillus specie* etc. as displayed in Table 6.

Table 6: The prevalence of bacterial isolates obtained from this study

ORGANISMS	FREQUENCY	PERCENTAGE (%)
<i>Pseudomonas spp</i>	3	23.1
<i>Streptococcus spp</i>	1	7.7
<i>Salmonella spp</i>	1	7.7
<i>Bacillus spp</i>	1	7.7
<i>Escherichia coli</i>	3	23.1
<i>Micrococcus spp</i>	1	7.7
<i>Klebsiella spp</i>	1	7.7

<i>Staphylococcus</i> spp	1	7.7
<i>Corynebacterium</i> spp	1	7.7
Total	13	100

The total fungal screening (Table 7) revealed that *Aspergillus* species, *Mucor* species, *Penicillium* species etc. were the fungal isolate obtained.

Table 7: Characterization and identification of fungal isolates

MORPHOLOGICAL CHARACTERISTICS	MICROSCOPIC EXAMINATION	SUSPECTED ORGANISMS
Yellow-green, blue-green, grey-green, filamentous growths that turn black sporulation	Long septate hyphae with swollen conidiopore	<i>Aspergillus</i> spp.
Green with raised rough surface colonies	Septate and branch brush like conidial conidiophore	<i>Penicillium</i> spp
White wooly growth that turns darker as it sporulates	Non septate hyphae with straight Sporangiophores spherical spores	<i>Mucor</i> spp.
Olivaceous-black, greewish colour on plate	Multicelled, matalae phialide	with <i>Alternaria</i> spp
Yellowish green on plate	Branch phialides chlamydosphores	with <i>Trichoderma</i> spp.
Whitish on petri dish	Multicelled-metalae pseudohyphae form	with <i>Candida</i> spp.
Grey colour on plate	Conidia bearing phialide	<i>Cephalosporium</i> spp.
Shiny velvet black fluffy growth	Curve septate hyphae conidia	with <i>Curvularia</i> spp.
Whitish felt mycelium	Branched conidiophores, smooth and rough conidia in pairs and chain	<i>Fusarium</i> spp.

The degradative potentials of hydrocarbon utilizing fungi isolates are presented in Table 8.

Table 8 Hydrocarbon Utilization Fungi

S/N	ORGANISMS	DEGRADATIVE SCREENING
1	<i>Aspergillus</i> spp	+
2	<i>Penicillium</i> spp	+
3	<i>Mucor</i> spp	+
4	<i>Aiternaria</i> spp	-
5	<i>Trichodermma</i> spp	-
6	<i>Candida</i> spp	+
7	<i>Cephalosporium</i> spp	+
8	<i>Curvularia</i> spp	-
9	<i>Fusarium</i> spp	+

The prevalence of fungal isolates showed that the fungal species present in this study, *Aspergillus* species, *Penicillium* species, *Mucor* species, *Candida* specie etc as displayed in Table 9.

Table 9: Percentage of occurrence of fungal isolate.

ORGANISMS	FREQUENCY	PERCENTAGE (%)
<i>Aspergillus</i> spp	1	12.5
<i>Penicillium</i> spp	1	12.5
<i>Mucor</i> spp	1	12.5
<i>Aiternaria</i> spp	1	12.5
<i>Trichodermma</i> spp	1	12.5
<i>Candida</i> spp	1	12.5
<i>Cephalosporium</i> spp	1	12.5
<i>Curvularia</i> spp	1	12.5
Total	8	100

DISCUSSION

The results obtained showed that soil samples impacted with crude oil and kerosene yielded low microbial population and diversity when compared to soil samples without kerosene and crude oil. The low microbial population obtained from soil impacted with crude oil and kerosene could be traced to the toxicity of crude oil and kerosene. This is in agreement with the reports of Mona *et al.* [15], who stated that excessive levels of heavy metals (hydrocarbons) can be damaging to organisms. The result corroborates the findings of Akubuenyi [2] which reported that the total heterotrophic bacterial and fungal counts of soil samples impacted with engine oil decreased when compared to unimpacted soil. Also, the reports of Shabir *et al.* [23], clearly stated that spills of crude oil and other hydrocarbons can lead to a significant decline in quality of soil and make it unfit for use. In a related study, Nyoyoko *et al.* [18] reported that higher concentration of crude oil has an adverse effect on fungi diversity while enhancing the population of fewer fungi.

The ability of some microorganisms to survive the impact of these hydrocarbons (Crude oil and Kerosene) could be traced to the fact that those microbes developed the physiological ability to use petroleum products as a source of carbon and energy. This corroborates the reports of Vinothini *et al.* [25], who opined that refined petroleum supply only carbon and energy source to resident microbes while crude oil supplies in addition to carbon and energy, mineral nutrients such as nitrogen, sulphur. These nutrients stimulate the growth of the hydrocarbon utilizing microorganisms, thereby enhancing their potentials for application in bioremediation. Raghad *et al.* [21] while working on the effect of petroleum hydrocarbons contamination on soil microorganisms and biodeterioration deposited that soil contamination with hydrocarbons significantly increased the count hydrocarbon utilizing bacteria.

The biochemical and identification processes revealed that the probable bacterial organisms present in the samples. They were found to include the following genera: *Pseudomonas species*, *Bacillus species*, *Staphylococcus species*, *Streptococcus species*, *Escherichia coli*, *Micrococcus species*, *Klebsiella species* and *Corynebacterium and Salmonella species*. While *Aspergillus species*, *Alternaria species*, *Candida species*, *Trichoderma species*, *Fusarium species*, *Mucor species*, *Penicillium species*, *Cephalosporium species* *Curvularia species* were the genera of fungi present in the sample. This result is in accordance with the reports of previous studies [17], were a total of twelve fungal isolates belonging to the genera; *Aspergillus species*, *Alternaria species*, *Candidia species*, *Fusarium species*, *Trichoderma species*, *Mucor species*, *Penicillium species*, *Cephalosporium species*, *Rhizopus species*, *Rhodoturula species* *Curvularia species*, *Cladosporium specie* and nine bacterial isolates belonging to the genera *Pseudomonas species*, *Bacillus species*, *Staphylococcus species*, *Streptococcus species*, *Escherichia coli*, *Micrococcus species*, *Klebsiella species* and *Corynebacterium and Salmonella species*, were isolated in the Niger Delta region. Akubuenyi [2] also obtained similar microorganisms from the study of the impact of engine oil on soil microbial community around mechanic workshops. Nyoyoko *et al* [18] and Sarmadi *et al* [22] in separate studies isolated and identified similar microorganisms,

Petroleum contamination of soil is particularly a serious problem because of the impact it has on soil functioning, and on the whole ecosystem. Agricultural soils, which are continually exploited to produce food and fodder, are particularly sensitive to contamination, as agricultural soils generally display poor resilience that is, they are incapable of recovering from any type of aggression, and contamination. This is in agreement with the position of Sarmadi *et al.* [22] that the kerosene contamination of soil leads to reduction of optimum moisture content of the soil. Research has shown that kerosene contamination increases cohesion, decreased internal strength of soils, and caused reduction in the permeability of soil [22]. The effect of crude oil and kerosene brought about alterations to soil functioning which reduces soil fertility and its consequent low crop yield.

CONCLUSION

Crude oil and kerosene contain substances that impact negatively on the growth of microbiota of soil origin. The combined effect of the compounds is more harmful than their individual effects. Because soil microorganisms are major contributors of soil fertility, then farmlands impacted with these products would not be suitable for agricultural purposes since the nitrogen fixing bacteria and other soil enriching microorganisms must have been either killed or inactivated by the toxicity of compounds. Hence, crude oil and kerosene should not be emptied into farmlands without proper treatment.

REFERENCES

1. Abii TA, Nwosu PC. The effect of oil-spillage on the soil of Eleme in Rivers State of the Niger Delta area of Nigeria. *Research Journal of Environmental Sciences*. 2009; 3(3): 316-320.
2. Akubuenyi, F.C. Determination of the influence of used engine oil on soil microbial community around mechanic workshops. *International Journal of Biotech Trends and Technology*. 2019; 9(4): 52-58
3. Benal T, Shivani K, Pagre RL, Chitnis S. Study of prevailing of deuteromycetous fungi on the petro-polluted soil, *International Research Journal of Biological Sciences*. 2014; 3: (11) 28-31.
4. Cheesbrough M. *District Laboratory Practice in Tropical Countries, Part 2. Cambridge University Press, United Kingdom*. 2006; 62-70.
5. Chikere C, Ekwuabu C. Culture-Dependent Characterization of Hydrocarbon Utilizing Bacteria in Selected Crude Oil-Impacted Sites in Bodo, Ogoni land, Nigeria. *African Journal of Environmental Science and Technology*. 2014; 8, 401-406.
6. Chikere CB, Azubuike CC. Catechol-2,3-dioxygenase screening in putative hydrocarbon utilizing bacteria. *Int. Res. J. Microbiol.* 2013;4 (1): 1-6.
7. Dombrowski N, Donaho JA, Gutierrez T, Seitz KW, Teske AP, Baker BJ. Reconstructing metabolic pathways of hydrocarbon-degrading bacteria from the Deep-water Horizon oil spill. *Nature Microbiology*. 2016; 1:16057.
8. Dvorak P, Nikel P, Damborský J, de Lorenzo V. Bioremediation 3.0: engineering pollutant-removing bacteria in the times of systemic biology. *Biotechnol Advances*. 2017; 35, 845–866.
9. Hadia-e-Fatima Ahmed A. Heavy Metal Pollution- A mini view. *Journal of Bacteriology and Mycology*. 2018; 6(3):179-181.
10. Hentati OR, Lachhab R, Ayadi M, Ksibi M. Toxicity assessment for petroleum-contaminated soil using terrestrial invertebrates and plant bioassays, *Environmental Monitoring and Assessment*, 2013; 185: 2989-2998.
11. John RC, Akpan MM, Essien JP, Ikpe DL. Variation in Rhizosphere Microbiological Properties of Tropical Legumes Grown on Oil Contaminated Wetland Utsoil. *Nigerian Journal of Microbiology*. 2010; 24(1):2081-2087.
12. Labud V, Garcia C, Hernander T. Effect of hydrocarbon pollution on the microbial properties of sandy and a clay soil. *Chemosphere*. 2007; 66:1863–1871.
13. Lopes AR, Faria C, Prieto-Fernández A, Trasar-Cepeda C, Manaia CM, Nunes OC. Comparative study of the microbial diversity of bulk paddy soil of two rice fields Subjected to organic and conventional farming. *Soil Biology and Biochemistry*. 2012; 43 (1):115-125.

14. Minai-Tehrani D, Herfatmanesh A. Biodegradation of aliphatic and aromatic fractions of heavy crude oil-contaminated soil: A pilot study. *Bioremediation Journal*. 2007; 11(2): 71-87.
15. Mona SZ, Nadia E, Hamman AM, Shalaby SI. Aquatic bioremediation of metals. *Life Science Journal*. 2014; 11 (4):66-72.
16. Murray PR, Baron EJ, Jorgensen JH, Pfaller M A, Tenover FC, Tenover FC. (ed.), *Manual of clinical microbiology*, 8th edition. ASM Press, Washington, DC; 2003.
17. Nseabasi NO, Antai SP. Toxic effect of kerosene contamination on the survival of bacteria and fungi species in soil from Niger delta. *International Research Journal of Microbiology*. 2012; 3(12): 382-387.
18. Nyooyoko VA, Anyanwu C, Christopher MA. Evaluation of the effect of crude oil and kerosene on soil microbial population. *International Journal of Applied and Natural Sciences*. 2022; 3: 117-138.
19. Onuoha SC, Victor O, Uraku AJ, Uchedu DO. Biodegradation potentials of hydrogen degraders from waste-lubricating Oil-spilled soils in Ebonyi State, Nigeria. *International Journal of Agriculture and Biology*. 2012; 13(4):586-590.
20. Onwurah INE, Ogugua VN, Onyike NB, Ochonogor AE, Otitoju OF. Crude oil spills in the environment, effects and some innovative clean-up biotechnologies. *Int. J. Environ. Res.*, 2007; 1(4) 307-320.
21. Raghad NH, Mehjin AA. Effect of petroleum hydrocarbons contamination on soil microorganisms and biodegradation. *Raf. J. Sci.* 2019; 28(1); 13-22.
22. Sarmadi MS, Zohrevand P, Rezaee M. Effect of kerosene contamination on the physical and mechanical properties of sandy soil. *Inno. Infrast. Solut.* 2019; 4:7
23. Shabir G, Afzal FM, Anwar R, Tahseen ZK. Biodegradation of kerosene in soil by a mixed bacterial culture under different nutrient conditions. *International journal of biodeterioration biodegradation*. 2008; 61:161.
24. Udeani CKT, Obroh AA, Okwuosa NC, Achukwu UP, Azubike N. Isolation of bacteria from mechanic workshops soil environment contaminated with used engine oil. *African Journal of Biotechnology*. 2009; 8(22): 6301-6303.
25. Vinothini C, Sudhakar S, Ravikumar R. Biodegradation of petroleum and crude oil by *Pseudomonas putida* and *Bacillus cereus*. *International journal of current microbiology and applied sciences*. 2015; 41: 318-329.
26. Wang C, Liu X, Guo J, Lv Y, Li Y. Biodegradation of marine oil spill residues using aboriginal bacterial consortium based on Penglai 19-3 oil spill accident. *China. Ecotoxicology and Environmental Safety*. 2018; 159.
27. Xu X, Liu W, Tian S, Wang W, Qi Q, Jiang P, Yu H. Petroleum Hydrocarbon-Degrading Bacteria for the Remediation of Oil Pollution under Aerobic Conditions: A Perspective Analysis. *Frontiers in Microbiology*. 2018; 9:145-156.