

Original Research Article

Antimicrobial Activity of Lactic Acid Bacteria Against Food-Borne Pathogens from Selected Fruits Sold In Major Markets, Ibadan, Nigeria.

Abstract

This study was conducted to determine the effect of antimicrobial compounds produced by Lactic acid bacteria against food-borne pathogens from selected fruits sold in major markets in Ibadan. The Lactobacillus species isolated from healthy fruits includes *L. casei*, *L. brevis*, *L. desidosus*, *L. jenseni*, *Lactiplantibacillus spp.* and *Fructilactobacillus spp.* while *Aeromonashydrophylia*, *Enterobacteraerogene*, *Eschericia coli*, *Salmonella typhii*, *Shigelladysentriae*, *Pseudomonas fluorescens*, *Bacillismegaterium*, *Candida valida*, *Saccharomyces cerevisiae*, *Rhizopusstolonifer* were isolated from spoilt fruits. Gram negative isolates were 100% resistant to Cefuroxime, Amoxycillin/Clauvulanate and Ampicillin while majority of the isolates were highly sensitive to Ofloxacin while Gram positive isolates were 100% resistant to Cloxicillin and highly sensitive to Ofloxacin and Gentamycin. *Lactiplantibacillus* spp had highest amount of Lactic acid (5.6 g/l) while *L. casei* had the lowest yield (3.6 g/l) at 48 hours. *L. casei* had highest amount of hydrogen peroxide (0.00036 g/l) while *Lactiplantibacillus spp.* and *Fructilactobacillus spp.* had lowest yield (0.00021 g/l) at 48 hours. The highest amount of diacetyl (3.01 g/l) was produced by *L. jenseni* while the lowest amount was observed in *L. brevis* (0.43 g/l) at 48 hours. The maximum inhibitory activity was observed in *L. brevis* against *B. megaterium* with diameter of 18mm zone of inhibition while the minimum activity by *L. desidosus*, *L. jenseni*, *Fructilactobacillus spp.* was observed against *B. subtilis* and *Lactiplantibacillus* spp against *K. pneumoniae* with diameter 8mm zone of inhibition. Antibiotic-resistant pathogenic bacteria and spoilage fungi found in this study could be a major threat to public health.

Keywords: Antibiotic-resistant bacteria, foodborne pathogens, Lactobacillus species

Introduction

Lactic acid bacteria (LAB) belong to groups of Gram-positive bacteria, with a common characteristic that produces lactic acid as the main product during fermentation of carbohydrates. They are anaerobic or facultative aerobic cocci or rod-shaped cells, non-sporulating, and acid-tolerant organisms [1].

Fresh fruits and vegetables are major component of human's diet because of their nutritional and medicinal properties and low energy content [2]. Several diseases such as Heart disease, colon cancer, obesity, and diabetes etc, can be reduced with a high intake of fruits and vegetables. Raw fruits and vegetables can be contaminated with pathogenic organisms of human health significance, and can occur directly or indirectly through animals or insects, soil, water, dirty equipment, and human handling etc. Fruits have low pH, high moisture content and nutrient composition, as a result of these are very susceptible to attack by pathogenic fungi, also causing rots which makes them unfit for consumption by producing mycotoxins [3, 4].

LAB produce an array of antimicrobial compounds, such as organic acids (lactic, citric, acetic, fumaric, and malic acid), hydrogen peroxide, CO₂, diacetyl, ethanol, reuterin, acetaldehyde, acetoin, ammonia, bacteriocins, bacteriocin-like inhibitory substances (BLIS), and other vital metabolites, which possess strong inhibitory effects on the growth and toxin production of many microorganisms [5, 6]. Furthermore, the antimicrobial effect of lactic acid bacteria is as a result of competition with pathogenic microorganisms for available nutrients [7].

Comment [RA1]: Verbose, should be re-phrased. Suggestion: "Antimicrobial effects of Lactic Acid Bacteria on Food-Borne Pathogens isolated from some fruits sold in Ibadan, Nigeria"

Comment [RA2]: remove

Comment [RA3]: what proportion or percentage of the isolates?

Comment [RA4]: remove and replace with a full stop.

Comment [RA5]: This conclusion does not project the significance of study i.e possible antimicrobial or preservative effects of the substances produced by these Lactic acid bacteria on pathogenic bacteria and spoilage fungi.

Comment [RA6]: Keywords are weak.

Comment [RA7]: Remove italics

Comment [RA8]: Repetition, remove

Comment [RA9]: Reframe

There is high prevalence of food borne pathogens on fruits due to contamination coupled with high antibiotic resistance of pathogens leading to food-borne illness which are hazardous to human health. Lactic acid bacteria is well known to produce antimicrobial compounds which inhibit microorganisms and also useful in preventing diseases. In this study, exploring antimicrobial potentials of LAB from sources such as apple, pineapple and soursop is necessary. Therefore, this study was conducted to determine the antibiotics resistance pattern of pathogens isolated from unhealthy fruits; quantification and antagonistic activity of antimicrobial compounds produced by isolated Lactic acid bacteria against isolated pathogens.

Comment [RA10]: Re-write this portion

2. Materials and methods

Comment [RA11]: Should not contain bullets. Remove the bullets

2.1 Sample Collection

The fruits samples such as apple, pineapple and soursop used for this study were randomly purchased from different locations in Bodija and Oje market in Ibadan, Oyo State, Nigeria. The healthy fruits samples were collected into labeled sterile containers without washing and transported to the laboratory for microbial analysis. The unhealthy (spoilt) fruits were obtained from the healthy fruits when spoilage occurs.

Comment [RA12]: Mention in specific terms the types and numbers of fruits used in the study.

Comment [RA13]: For how long were the fruits kept before spoilage and under what conditions were they kept?

2.2 Sample Preparation

Samples were prepared according to the method of Akoachere [8] with slight modifications. Twenty-five grams of each sample was weighed and homogenized by blending in 225 mL of sterile distilled water. One millilitre of the homogenate was introduced into 9 mL of the distilled water in a test tube, labelled 1:10 (10^{-1}) dilution and serially diluted to five other test tubes labeled 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} and 10^{-6} ; the procedure was repeated for each sample and the blender was carefully cleaned and disinfected in between samples to prevent any cross contamination.

Isolation and Characterization of probable lactic acid bacteria and food-borne pathogens were done accordingly [9, 10].

Comment [RA14]: Vague and scanty.. What criteria were used for the other organisms isolated.

2.3 Antibiotic Susceptibility Test

Antibiotic Susceptibility Test of the samples was analyzed in accordance with Clinical and Laboratory Standards Institute Standard guidelines [11].

Comment [RA15]: mention antibiotics used. Moreover, is this part a comparative study on antimicrobial activity or does it serve as control for the study?

2.4 Quantification of antimicrobial compounds produced by lactic acid bacteria

The test organism were grown in MRS broth for 48 hours and centrifuged at 4000 x g for 15 minutes.

2.5 Lactic acid, Hydrogen Peroxide and Diacetyl

Lactic acid, Hydrogen Peroxide and Diacetyl of the samples were determined following analytical procedures [12].

2.6 Screening for antagonistic activity of cell-free supernatant

Lactic acid bacteria were inoculated into de Man Rogosa Sharpe (MRS) broth and incubated at 30°C for 48 hours in a candle jar. After 48 hours of incubation, cell-free supernatants were obtained by centrifuging at 4000 rpm for 15 minutes at 4°C. The antagonistic activity of the cell-free supernatant was determined by agar well diffusion assay (AWDA) against foodborne pathogens isolated from spoilt fruits. Mueller Hinton plates and Potato Dextrose Agar plates were inoculated with bacterial and fungal isolates respectively using sterile swab. Well of 7mm were cut into the agar plate and 100µl of cell-free supernatant was dispensed into each well and incubated aerobically for 24 hours and 3 days

Comment [RA16]: Remove comma

Comment [RA17]: The antagonistic effects of the ce-free supernatant on bacteria isolated from spoilt fruits was determine using the agar-well diffusion assay (AWDA)

Comment [RA18]: Mueller Hinton agar plates

Comment [RA19]: Remove comma

for Bacterial and fungal isolates respectively at 30°C. The antimicrobial activity was determined by measuring the diameters of zone of inhibition around the wells [13].

Comment [RA20]: Inconsistency of words for inhibitory effects .

3. Results and discussion

The antibiotic resistance patterns of Gram negative bacteria isolated from spoiled fruits are shown in Table 1. All the isolates were 100% resistant to Cefuroxime (CRX), Amoxycillin/Clavulanate (AUG) and Ampicillin (AMP) while majority of the isolates were highly sensitive to Of This study was conducted to determine the effect of antimicrobial compounds produced by Lactic acid bacteria against food-borne pathogens from selected fruits sold in major markets in Ibadan ofloxacin (OFL) followed by Gentamycin (GEN) and Ciprofloxacin (CPR). All the isolates are resistant to three to four antibiotics as *Pseudomonas putida* isolated from apple was sensitive to Nitrofurantoin (NIT) with 33mm zone of inhibition. While the antibiotics resistance pattern of Gram positive bacteria isolated from spoiled fruits are shown in Table II. All the isolates were 100% resistant to Cloxacillin (CXC) and highly sensitive to Ofloxacin (OFL) and gentamycin. The isolates are resistant to at least one antibiotic. *Bacillus coagulans* had the highest sensitive of 27mm to Ceftriaxone (CTR).

Comment [RA21]: This result is inconsistent with the title. Seems like a deviation from the aim of the study. "This study was conducted to determine the effect of antimicrobial compounds produced by Lactic acid bacteria against food-borne pathogens from selected fruits sold in major markets in Ibadan"

The pathogens isolated from spoiled fruits in this study varied in their sensitivity to different antibiotics. The reason for antibiotic resistance could be due to chromosomal or plasmid resistance of the microorganism cell constituents [14]. The high resistance to ampicillin (100%) recorded in this study is similar to that obtained by Issa [15] and Yakubu [16] where 100% resistance to ampicillin were obtained from isolates in milk and processed meat respectively and Endang [17] who recorded a similar high resistance (87.0%) in isolates obtained from salted fish. The low resistance of isolates to Ofloxacin, Ciprofloxacin and Gentamicin in this study is in agreement with the study reported by Yakubu [18] and Rahimi [19] where over 80.0% of isolates from various sources were found to be susceptible to each of these antimicrobial agents. Recently, antibiotic-resistant microbial species increases including moulds and yeasts, they are becoming resistant also to preservatives such as sorbate and benzoate as well as chemical detergents [20].

The antimicrobial compounds produced by Lactic Acid Bacteria were quantified using standard methods. *Lactiplantibacillus spp.* produced the highest amount of Lactic acid (5.6 g/l) at 48 hours while *Lacticaseibacillus casei* had the lowest yield (3.6 g/l) at 48 hours; *Lacticaseibacillus casei* produced the highest amount of hydrogen peroxide (0.00036 g/l) while *Lactiplantibacillus spp.* and *Fructilactobacillus spp.* produced the lowest yield (0.00021 g/l) at 48 hour. The highest amount of diacetyl (3.01 g/l) was produced by *Lactobacillus. jenseni* while the lowest amount was observed in *Levilactobacillus brevis* (0.43 g/l) at 48 hours as shown in Table 3.

The antagonistic activity of antimicrobial compounds produced by Lactic acid bacteria against pathogenic organisms isolated from spoiled fruits are shown in Table 4. Among the pathogenic organisms examined, the maximum inhibitory activity was observed in *Levilactobacillus brevis* against *B. megaterium* with 18mm zone of inhibition while the minimum inhibitory activity by *Lactobacillus desidosus*, *Lactobacillus jenseni*, *Fructilactobacillus spp.* was observed against *B. subtilis*, and *Lactiplantibacillus spp.* against *K. pneumoniae* with 8mm zone of inhibition.

Most of the antimicrobial compounds produced by Lactic acid bacteria isolated from healthy fruits shows antagonistic activity against Gram positive, Gram negative bacteria and fungi with *L. brevis* showing the highest zone of inhibition of (18mm) against *B. megaterium* while the lowest zone of inhibition (8mm) was observed in *L. desidosus*, *L. jenseni*, *Fructilactobacillus spp.* against *B. subtilis*,

Table 1: Antibiotic resistance pattern of Gram-negative bacteria isolated from spoiled fruits

Isolate (Location)	CAZ	CRX	GEN	CPR	OFL	AUG	NIT	AMP
ANTIBIOTIC/ZONE OF INHIBITION (MM)								
OJE MARKET								
<i>Klebsiellapneumoniae</i> (P)	21 (S)	0 (R)	21 (S)	26 (S)	23 (S)	0 (R)	0 (R)	0 (R)

Comment [RA22]: Clumsy and poorly designed.

Comment [RA23]: How did you interpret the zone of inhibition to be sensitive or resistant?

<i>Aeromonashydrophylia</i> (P)	0 (R)	0 (R)	0 (R)	19 (I)	18 (S)	0 (R)	12 (R)	0 (R)
<i>Enterobacteraerogenes</i> (S)	18 (I)	0 (R)	0 (R)	19 (I)	23 (S)	0 (R)	24 (S)	0 (R)
<i>Aeromonashydrophylia</i> (A)	21 (S)	0 (R)	0 (R)	27 (S)	27 (S)	0 (R)	0 (R)	0 (R)
<i>Pseudomonas putida</i> (A)	29 (S)	0 (R)	21 (S)	26 (S)	29 (S)	0 (R)	33 (S)	0 (R)
<i>Eschericia coli</i> (A)	30 (S)	0 (R)	20 (S)	24 (S)	27 (S)	0 (R)	0 (R)	0 (R)
<i>Salmonella typhi</i> (A)	27 (S)	0 (R)	23 (S)	24 (S)	24 (S)	0 (R)	0 (R)	0 (R)
<i>Salmonella typhi</i> (P)	0 (R)	0 (R)	15 (S)	18 (I)	20 (S)	0 (R)	0 (R)	0 (R)
<i>Shigelladysentriae</i> (S)	23 (S)	11 (R)	19 (S)	13 (R)	20(S)	0(R)	0(R)	0(R)
<i>Shigelladysentriae</i> (P)	22 (S)	11 (R)	18 (S)	20 (I)	20(S)	11(R)	0(R)	0(R)
BODIJA MARKET								
<i>Enterobacteraerogenes</i> (S)	0 (R)	0 (R)	0 (R)	25 (S)	14 (I)	0 (R)	0 (R)	0 (R)
<i>Pseudomonas cepacia</i> (S)	28 (S)	0 (R)	21 (S)	28 (S)	28 (S)	0 (R)	20 (S)	0 (R)
<i>Enterobacteraerogenes</i> (P)	0 (R)	0 (R)	20 (S)	19 (I)	20 (S)	0 (R)	0 (R)	0 (R)
<i>Pseudomonas fluorescens</i> (S)	0 (R)	0 (R)	25 (S)	19 (I)	25 (S)	0 (R)	15 (I)	0 (R)
<i>Aeromonashydrophylia</i> (S)	0 (R)	0 (R)	21 (S)	14 (R)	18 (S)	0 (R)	0 (R)	0 (R)
<i>Pseudomonas cepacia</i> (A)	24 (S)	0 (R)	21 (S)	23 (S)	23 (S)	0 (R)	0 (R)	0 (R)

CAZ: Ceftazidime; CRX: Cefuroxime; GEN: Gentamicin; CPR: Ciprofloxacin; OFL: Ofloxacin; AUG: Amoxicillin/Clavulanate; NIT: Nitrofurantoin; AMP: Ampicillin; P: Pineapple; A : Apple ; S : Soursop ; R: Resistance; S: Sensitive; I: Intermediate

and *Lactiplantibacillus spp.* against *K. pneumoniae*. The Gram negative organisms namely *Pseudomonas cepacia*, *Pseudomonas fluorescens*, *Aeromonashydrophylia*, *Eschericia coli* showed slight or no inhibition compared to Gram positive organisms. This is in accordance with earlier reports which revealed that different strains of LAB possess more active inhibition against Gram positive organisms compared with Gram negative organisms [21]. This may be due to the presence of an outer protective membrane in Gram negative organisms, which covers the cytoplasmic membrane and peptidoglycan layer of the cells. It is responsible for preventing molecules such as antibiotics [22].

Table 2: Antibiotic resistance pattern of Gram-positive bacteria isolated from unhealthy fruits

Isolate (Location)	CAZ	CRX	GEN	CTR	ERY	CXC	OFL	AUG
ANTIBIOTIC/ZONE OF INHIBITION (MM)								
OJE MARKET								
<i>Bacillus coagulans</i> (S)	23 (S)	11 (R)	21 (S)	27 (S)	0 (R)	0 (R)	20 (S)	15 (I)
<i>Bacillus pulmilus</i> (A)	18 (I)	0 (R)	17 (S)	13 (I)	0 (R)	0 (R)	26 (S)	0 (R)
<i>Bacillus polymyxa</i> (P)	19 (I)	17 (I)	17 (S)	22 (I)	0 (R)	0 (R)	23 (S)	16 (I)
BODIJA MARKET								

<i>Bacillus licheniformis</i> (S)	21 (S)	0 (R)	17 (S)	24 (S)	0 (R)	0 (R)	21 (S)	0 (R)
<i>Bacillus megaterum</i> (P)	11 (R)	9 (R)	13 (I)	17 (R)	17 (I)	0 (R)	12 (I)	15 (I)
<i>Bacillus subtilis</i> (A)	20 (I)	16 (I)	17 (S)	13 (R)	15 (I)	0 (R)	21 (S)	19 (S)

CAZ: Ceftazidime; **CRX:** Cefuroxime; **GEN:** Gentamicin; **CTR:** Ceftriaxone; **ERY:** Erythromycin; **CXC:** Cloxicillin; **OFL:** Ofloxacin; **AUG:** Amoxicillin/ClavulanateP : Pineapple; A : Apple ; S : Soursop ; R: Resistance; S: Sensitive; I: Intermediate

Lactic acid bacteria are widely known to have antimicrobial compounds which are active against closely related bacteria and other pathogens [23]. It was reported by Ogunbanwo [24] that selected *Lactobacillus* strains (*L. plantarum*F1 and *L. brevis*OG1) produced bacteriocin, which showed inhibitory activity against Gram positive and Gram-negative strains (*E. coli*, *E. faecalis*, *B. cereus*, *S. aureus*, *Sh. dysentery*, *Sh. flexneri* and *Listeria monocytogenes*). Similar trend of result was reported by Navarro [25] reported similar result of antimicrobial activity of *L. plantarum* against both Gram-negative and Gram-positive bacteria with inhibition zone between 9-22mm. It was also reported that there was poor antimicrobial activity of cell free supernatant of *L. brevis* against Gram positive and Gram negative strains with diameter of inhibition zone between 7-15mm [26]. Galalet *et al.*, 2012 also

Table 3: Quantity of antimicrobial compounds produced by Lactic Acid Bacteria

Isolates	Antimicrobial compounds (g/l)		
	Lactic acid	Hydrogen peroxide	Diacetyl
<i>Lactocaseibacillus casei</i>	3.6	0.00036	1.72
<i>Levilactobacillus brevis</i>	5.4	0.00029	0.43
<i>Lactobacillus jensenii</i>	5.3	0.00032	3.01
<i>Lactobacillus desidosus</i>	5.3	0.00021	2.88
<i>Lactiplantibacillus spp.</i>	5.6	0.00021	0.77
<i>Fructilactobacillus spp.</i>	4.6	0.00032	0.86

Table 4: Antagonistic activity of antimicrobial compounds produced by Lactic acid bacteria against pathogenic organisms isolated from unhealthy fruits

Organisms	<i>L. brevis</i>	<i>L. casei</i>	<i>L. desidosus</i>	<i>L. jenseni</i>	<i>Lactiplantibacillus spp.</i>	<i>Fructilactobacillus spp.</i>
<i>Gram -positive bacteria</i> (Zone of inhibition in mm)						
<i>B. coagulans</i>	-	13	9	-	-	-
<i>B. pulmilus</i>	-	-	-	-	-	-
<i>B. megaterium</i>	18	-	11	14	10	13
<i>Bacillus subtilis</i>	-	-	8	8	12	8
<i>B. licheniformis</i>	15	-	12	14	14	9
<i>Gram- negative bacteria</i>						
<i>P. cepacia</i>	-	-	-	12	-	-
<i>P. putida</i>	-	-	-	-	13	9
<i>P. fluorescens</i>	-	-	-	-	-	10
<i>A. hydrophylia</i>	-	-	-	-	-	-
<i>E. aerogenes</i>	-	10	-	9	-	-
<i>E. coli</i>	-	-	-	-	-	-
<i>K. pneumoniae</i>	-	-	13	-	8	-
<i>S. dysentriae</i>	-	-	13	14	10	13
<i>S. typhi</i>	10	-	-	-	12	-
<i>Fungi (Yeast and moulds)</i>						
<i>C. valida</i>	10	14	11	11	12	10
<i>S. cerevisiae</i>	-	12	10	11	10	13
<i>R. stolonifer</i>	11	10	-	-	14	-

4. Conclusion

The use of antimicrobial compounds produced by Lactic acid bacteria could help to curtail multiple drugs resistance. Fruit contamination with antibiotic-resistant bacteria when disposed carelessly into the environment especially water bodies, could lead to water pollution which can be a major threat to public health

Comment [RA24]: Vague. Substantiate with a better write-up.

Comment [RA25]: How?

Comment [RA26]: Poor conclusion.

REFERENCES

- [1] Shoukat S. Potential anti-carcinogenic effect of probiotic and lactic acid bacteria in detoxification of benz[a]pyrene: A review. *Trends Food Sci. Technol.*, 2020; 99, 450–459.
- [2] Charlton K, Kowal P, Soriano MM, Williams S, Banks E, Vo, K. & Byles J. et al. Fruit and vegetable intake and body mass index in a large sample of middle-aged Australian men and women. *Nutrients*, 2014, 6: 2305–2319.
- [3] Berger CN, Sodha SV, Shaw RK, Griffin PM, Pink D. Hand P. & Frankel G. et al. Fresh fruit and vegetables as vehicles for the transmission of human pathogens. *Environ. Microbiol.*, 2020; 12, 2385–2397.
- [4] More AS, Ranadheera CS, Fang Z, Warner R. & Ajlouni S. et al. Biomarkers associated with quality and safety of fresh-cut produce. *Food Biosci.*, 2020; 34, 100524.
- [5] Zehra SA, Javed S, Nadeem SG. & Hakim ST. Lactic acid bacteria from fresh fruits and vegetables as biocontrol agent of foodborne bacterial pathogens. *RADS J. Biol. Res. Appl. Sci.* 2014; 5, 36–45.
- [6] Bartkiene E, Lele V, Ruzauskas M, Domig KJ, Starkute V, Zavistanaviciute P, Bartkevics V, Pugajeva I, Klupsaite D. & Juodeikien G. et al. Lactic acid bacteria isolation from spontaneous sourdough and their characterization including antimicrobial and antifungal properties evaluation. *Microorganisms*, 2020; 8: 64.
- [7] Siroli L, Patrignani F, Serrazanetti DI, Gardini F. & Lanciotti R. Innovative strategies based on the use of bio-control agents to improve the safety, shelf life and quality of minimally processed fruits and vegetables. *Trends Food Sci. Technol.*, 2015; 46: 302–310.
- [8] Akoachere JFTK, Bughe RN, Oben BO, Ndip LM. & Ndip RN. Phenotypic characterization of human pathogenic bacteria in fish from the coastal waters of South West Cameroon: Public health implications. *Reviews in Environmental Health* 2009; 24:147–155.
- [9] Cheesebrough M. *District Laboratory Practices in Tropical Countries*. Cambridge University Press, Edinburgh, UK. 2006; pp. 382–407.
- [10] Zheng JS, Wittouck E, Salvetti CMAP, Franz HMB, Harris P, Mattarelli PWO’Toole, B. Pot P, Vandamme J, Walter K, Watanabe S, Wuyts GE, Felis MG, Gänzle and Lebeer S. et al. A taxonomic note on the genus *Lactobacillus*: Description of 23 novel genera, emended description of the genus *Lactobacillus* Beijerinck 1901, and union of *Lactobacillaceae* and *Leuconostocaceae*. *Int. J. Syst. Evol. Microbiol.* 2020; 70:2782–2858. <https://doi.org/10.1099/ijsem.0.004107>.
- [11] Clinical and Laboratory Standards Institute (CLSI). Performance Standard for Antimicrobial Susceptibility Testing. 2020; 30th ed. CLSI supplement M100, Wayne, PA.
- [12] AOAC (2000). Official Methods of Analysis. Association of Official Analytical Chemists. 16th edn. Washington D.C. ISBN 2-93 558442-0.

- [13] Ogunbanwo ST, Ilesanmi F, and Molokwu AJ. Thermal stability of lactic acid bacteria metabolites and its application in preservation of tomato paste. *Malaysian Journal of Microbiology*, 2014;10(1): 15-23.
- [14] Pooma V. &Kaur R. "Prevalence and growth of pathogens on salad vegetables, fruits and sprouts". *Int. J. Hyg. Environ. Health*, 2001; 203: 205-213.
- [15] Issa ZM, Mustakim M, Mohamed SAS, Muda NM, Yen LH.&Radu S. Antibigram profiles of *Listeria monocytogenes* isolated from foods. *International Conference on Biotechnology and Food Science*, 2011; *IPCBEET*: 133-137.
- [16] Yakubu Y, Salihu MD, Faleke OO, Abubakar MB, Junaidu AU, Magaji AA, Gulumbe ML. &Aliyu RM. Prevalence and antibiotic susceptibility of *Listeria monocytogenes* in raw milk from cattle herds within SokotoMetropolist, Nigeria. *J. Vet Sci.*2012b;**10**:13-17.
- [17] Endang P, Radu S, Zaiton H, Rusul G. Antimicrobial drug resistance and resistance factor transfer among *Listeria* species. *Asian Fishery Science*, 1998;11: 261-270.
- [18] Yakubu Y, Salihu MD, Faleke OO, Abubakar MB, Junaidu AU, Magaji AA, Gulumbe ML &Aliyu RM. Disinfectant effect of methylated Ethanol against *Listeria species*. *Veterinary World*.2012a ;**5**: 91-93.
- [19] Rahimi E, Momtaz H, Sharifzadeh A, Behzzadnia A, Astari MS, Zandi S., Riahi M, Momeni M . Prevalence and antimicrobial resistance of *Listeria* species isolated from traditional dairy products in ChaharMahal and Bakhtiyari, Iran, Bulg. *J. Vet. Med.* 2012;15: 115-122.
- [20] Schnurer J. & Magnusson J. Antifungal lactic acid bacteria as biopreservatives. *Trends Food Sci Tech*, 2005;16: 70-80.
- [21] Savino F, Cordisco L, Tarasco V, Locatelli E, DiGioia D, Roberto OR&Matteuzzi, D. Antagonistic Effect of *Lactobacillus* strains against Gas producing Coliforms Isolated from Colicky Infants. *Biomed Central Microbiology*, 2011; 11: 157.
- [22] Dundar H. Characterization and Purification of a bacteriocin produced by *Leuconostocmesenteroides* subsp. *cremoris*. Ph.D.thesis; 2006 Department of Biotechnology, Faculty of Medicine, Hacettepe University, Turkey.
- [23] Flythe MD. and JB. Russell,. The effect of Ph and a bacteriocin (bovicinHC5) on *Clostridium sporogenes* MD1, a bacterium that has the ability to degrade amino acids in ensiled plant materials. *FEMS Microbiol. Ecol.*, 2004; 47: 215-222.
- [24] Ogunbanwo ST, Sanni AI, Onilude AA. Characterization of bacteriocin produced by *Lactobacillus plantarum*F1 and *Lactobacillus brevis*OG1. *African Journal of Biotechnology*, 2003; 2(8): 219-227.
- [25] Navarro LM, Zarazaga J. Saenz F. Ruizrharea and C. Torres. Bacteriocin production by lactic acid bacteria isolated from Rioja red wines. *J. Applied Microbiol.*,2000;**88**: 44-51.
- [26] Galal AA, Azzeddine K, Khadya K, Reda C. and Zakaria M. Screening of Lactic acid bacteria isolated from dried fruits and study of their Antimicrobial activity. *Middle East Journal of Scientific Research*,2012; 11(2): 209-215