

Isolation and screening of agriculturally important bacteria (PGPR) from organic sources of nutrient (Panchgavya, Jeevamrit and FYM) for future use

ABSTRACT

Agriculture is an essential component to maintain the human health but due to sudden increase in population the demand has increased and to meet the requirement the use of chemical fertilizer has also increase. The continuous use of chemical fertilizer and pesticides degrade the quality and fertility of soil and also reduce the yield of various crops. The organic sources such as panchgavya, jeevamrit and FYM consists load of beneficial microorganisms that promote the plant growth, yield and promote natural resources. The present study was carried out by isolating the bacteria from organic sources (panchgavya, jeevamrit and FYM) because minute amount of bacterial culture is sufficient to prepare large one according to need and can be stored for longer time as compare to organic input. The samples of panchgavya, jeevamrit and FYM were collected from different location of Himachal Pradesh. A total of 32 bacterial isolates from panchgavya, 43 from jeevamrit and 59 from FYM were isolated and screened for multiple plant growth promoting traits. Only 4 bacterial isolates from panchgavya, 2 from jeevamrit and 2 from FYM were preferred on the basis of maximum plant growth promoting traits like P-solubilization, Nitrogen fixer, Siderophore production, HCN production, IAA production and Antifungal activity. From the selected bacterial isolates 67.78% were P-solubilizers, 60.37% were Nitrogen fixer, 62.37% were Siderophore producer, 34.29% HCN producer, 45.88% IAA producer and 78.22% show antagonism against *Pythiumgraminicolum* and 74.81% show antagonism against *Collectotrichumcapsici*. Hence these bacterial isolates have potential to act as biofertilizer for environment friendly sustainable agriculture systems and alternative to harmful chemicals.

Keywords: Organic sources of nutrient, microorganism, biocontrol, PGPR, Plant growth, biofertilizers

Introduction

The agriculture and food security mainly depends upon the fertility status of soil to sustain the life in all forms. The microorganisms present in the soil play an important role in plant

growth promotion and health (Wang *et al.*, 2017). The continues uses of chemical fertilizers and pesticides in the agricultural fields leads to degradation of soil quality and fertility, thus agricultural land with fertile soil is decreasing day by day (Bhardwaj *et al.*, 2018). Nowadays isolation of microorganism from various traditional organic inputs like panchagavya, jeevamrutha and FYM plays important role in the expansion of sustainable agriculture and benefit the society by socially, commercially and environmentally. The organic inputs which are used in organic farming system are the fermented product of liquid formulations obtained from cow and used for plant growth promotion (Ramesh *et al.*, 2005). These organic inputs, like panchgavya, jeevamrit and FYM are used as widely conventional way to protect plants by acting as biocontrol agent and soil microorganisms from harmful chemical fertilizers and pesticides (Gunwantrao and Patil, 2015). Panchagavya is the mixture of five ingredients (dung, urine, milk curd and ghee) obtained from cow. Jeevamrit is also prepared from cow dung, urine, gram flour, organic soil and jaggery by fermentation. Panchagavya and jeevamrit are the richest sources of beneficial microorganisms which sustain, stimulate the plant growth and help in getting better vegetative growth and also good quality yield (Devakumaret *al.* 2011). The interest of scientist has been increased toward isolation of microorganism from panchagavya, jeevamrit and FYM for the enhanced growth of various crops in organic agriculture from last few years. Devakumaret *al.*, (2008) and Srinivaset *al.*, (2010) have reported the survival of many beneficial microorganisms viz., nitrogen fixers, phosphorus solubilizers, actinomycetes and fungi in panchgavya, jeevamrutha and FYM. These organic inputs contain beneficial microorganisms, predominately lactic acid bacteria, *Bacillus* and *Pseudomonas*, certain fungi such as *Aspergillus* and Yeast (*Saccharomyces cerevisiae*). Hence the microorganism present in the organic sources of nutrient such as panchgavya, jeevamrit and FYM have potential as biofertilizer can be used for enhanced growth of various crops and maintains the fertility of soil by natural means.

Materials and Method:

Collection of samples

The samples of panchgavya, jeevamrit and FYM were collected from Hamirpur, Solan and Una district of Himachal Pradesh. From each site two subsites were selected for sample collection. The samples were collected in plastic bags and placed in the Microbiology laboratory, Deptt. of Basic Science, College of Forestry, Dr YS Parmar University of

Horticulture and Forestry, Nauni-Solan. Panchgavya and Jeevamrit was also prepared in laboratory mixing the fresh cow urine, cow dung, cow milk, cow curd and cow ghee from local cow farm and placed in sterile containers and stored in refrigerator for further uses.

Method of Preparation of Panchgavya:

2.5 kg of fresh cow dung was taken and mixed thoroughly with 250 g of cow's ghee and kept in a mud pot. The mixture was kept for 2 days and mixed twice a day. On the 4th day, the cow's urine, milk curd and Jaggery was added. This solution was kept for another 9 days, shaken twice a day for about 20 minutes to provide aerobic fermentation which facilitates microbial growth and activity. On the 10th day the panchgavya solution was ready and was covered with a muslin or fine cloth to prevent the contamination caused by insect and flies.

Method of Preparation of Jeevamrit:

5 kg of fresh cow dung was mixed with 5 ltr of cow urine, 1 kg of jaggery, 0.5 kg of gram flour and 0.5 Kg of organic soil in pot. The mixture was kept for 7 days and mixed twice a day with wooden stick for about 20 minutes to provide aerobic fermentation which facilitates microbial growth and activity. On the 8th day the jeevamrit solution was ready and covered with a muslin or fine cloth to prevent the contamination caused by flies.

Isolation of microorganisms from Panchgavya, jeevamrit and FYM

Isolation was carried out by serial dilution method in which 1ml or 1g of panchgavya, jeevamrit and FYM sample were inoculated in test tube containing 9 ml of distill water. The samples were diluted up to 10⁻⁸ concentration. 0.1 ml of diluted sample was taken and spread on prepreud nutrient agar plate. After spreading the plates were incubated at 37C temperature for 24 hours in the incubator for the growth of colonies. The isolated colonies that showed diverse morphological variation was selected and inoculated separately in 5 ml of nutrient broth and incubated at 37°C for 24 hours. After incubation, it was again plated in nutrient agar plates to check the purity of isolates. After checking the purity of isolates, they were streaked on the slants and were stored in refrigerator for further use.

Bacterial isolates were screened out for the production of hydrogen cyanide (HCN) as per method described by Bakker and Schippers (1987). Bacterial cultures were streaked on prepreud plates of King's B medium amended with 1.4 g l⁻¹ glycine. Whatman No.1 filter paper strips were soaked in 0.5 per cent picric acid in 2 per cent sodium carbonate and were placed in

Comment [A1]: Where did you inspire? In the literature? Please indicate an author

the lid of each petriplates and plates were sealed with parafilm and were incubated at $35 \pm 2^{\circ}\text{C}$ for 1-4 days. Uninoculated control was kept for comparison of results. Plates were observed for change in color of filter paper from yellow to orangebrown.

Indole acetic acid (IAA) production

Bacterial cultures were grown in modified Luria Bertani broth amended with (5 mM L-tryptophan, 0.065% sodium dodecyl sulphate and 1% glycerol) 72 h at 35°C under shaking conditions. The cultures were centrifuged at 15,000 rpm for 20 minutes and supernatant were collected and stored at 4°C . The method described by Gordon and Palleg (1957) was used to determine the IAA equivalents; 3 ml of supernatant was pipette into test tube and 2 ml Salkowskireagent (2 ml 0.5 M $\text{FeCl}_3 + 98 \text{ ml } 35\% \text{ HClO}_4$) was added to it. The tubes containing the mixture were left for 30 minutes (in dark) for colour development. Intensity of colour was measured spectrophotometrically at 535 nm. Similarly, colour was also developed in standard solution of IAA ($10\text{-}100 \mu\text{g mL}^{-1}$) and a standard curve was established by measuring the intensity of thiscolour.

Antagonistic activity of bacterial isolates against test fungus

Agar streak plate method was used to test the efficacy of bacterial isolates against the test fungus. A loop full of 48 h old culture of each isolate were streaked a little below the centre of the prepared MEA petri plate and incubated at $30 \pm 2^{\circ}\text{C}$ for 24 h to check contamination. Mycelial disc of 5 days old culture of the test fungal pathogen (*Fusariumoxysporum* and *Rhizoctoniasolani*) was placed separately on one side of the streak in each plate.

$$I = \frac{C-T}{C} \times 100$$

Where, I = Per cent growth inhibition
 C = Growth of fungus in control
 T = Growth of fungus in treatment

STATISTICAL ANALYSIS

The data collected on various parameters under laboratory condition were subjected to statistical analysis as per methods outlined by Gomez and Gomez (1984). The CD at 5 per cent level was used for testing the significant differences among the treated means.

Results

Isolation of bacteria from Panchgavya, jeevamrit and FYM

A total of 27 bacterial isolates (14 from FYM, 5 from panchgavya and 8 from jeevamrit) were selected from different organic sources panchgavya on nutrient agar medium from different locations of Himachal Pradesh (Table 1 and Plate1).

Table 1: Isolation of bacteria from Panchgavya, jeevamrit and FYM

Organic source (OS)	Locations		
	Microbial count 10 ⁸ cfu/g/ml on NA media		
	Hamirpur	Una	Solan
FYM	92.44	107.46	78.26
Panchgavya	59.66	97.79	79.36
Jeevamrit	88.80	89.94	84.21
CD(0.05)	3.63		

Screening of bacterial isolates for multifarious plant growth promoting traits

On the basis of multifarious plant growth promoting traits i.e. P-solubilization, growth on nitrogen free medium, siderophore production, auxin, HCN production and antagonism against major fungal pathogens i.e. *Pythiumaphanidermatum* (causal organism of Rhizome rot) and

Comment [A2]: Where did you inspire? In the literature? Please indicate an author

Collectotrichumcapsici(causal organism of leaf spot) only best 3 bacterial isolate (1 from panchgavya, 1 from jeevamrit and 1 from FYM) were selected and again purified on the NA medium and preserved for further use (Plate 2-5).

Morphological Characterization of Selected bacterial isolates

On the basis of morphology and gram reaction out of selected 3 bacterial isolates, 2 isolates were gram positive rod and 1 bacterial isolate was gram negative rods (Table 2).

Table 2: Morphological characterization of selected bacterial isolates

Bacterial Isolates	Form	Elevation	Margin	Surface	Gram's Reaction	Shape
PGPR1	Irregular	Flat	Undulate	Rough	+	Rods
PGPR2	Circular	Flat	Lobate	Rough	-	Rods
PGPR3	Irregular	Raised	Curled	Smooth	+	Rods

Physiological Characterization of Selected isolates

The physiological characterization of selected bacterial isolates revealed that all the isolates were able to grown on a broad temperature range of 25 - 45°C but optimum temperature for the growth was found to be 35°C on the basis of maximum OD at 540 nm. The selected isolates were able to grow on the different pH range of 3.0 - 11.0 and with the maximum growth at the nearly neutral 7.0 pH. Similarly different incubation periods (24 - 96 h) were tested for the selected bacteria and found maximum growth at 24 - 48 hrs of incubation on the basis of OD at 540 nm (Figure 1-3).

Biochemical Characterization of Selected Isolates

All the selected three bacterial isolates were positive for Methyl red test, starch hydrolysis, Gelatin hydrolysis, Catalase test, Casein hydrolysis and fermentation of glucose. However, all the isolates were negative for Indole test, Citrate utilization and hydrogen sulphide (H₂S) production (Table-3).

Table 3: Biochemical characterization of selected bacterial isolates

Test	Bacterial Isolates		
	PGPR1	PGPR2	PGPR3
Indole test	-	-	-
Methyl red test	+	+	+
Citrate utilization test	+	-	-
Starch hydrolysis	+	+	+
Gelatin hydrolysis	+	+	+
Hydrogen sulphide production	+	-	-
Catalase test	+	+	+
Fermentation of Glucose	+	+	+
VogesProskauer test	+	+	+
Casein hydrolysis	+	+	+

Discussion

Isolation and screening of microorganisms from panchgavya, jeevamrit and FYM are new or less exploited source and can be used as alternative way to produce efficient biofertilizer for sustainable agriculture. During the study panchgavya and jeevamrit were prepared by mixing the five different products of cow like cow dung, cow urine, cow milk, cow ghee, cow curd, gram flour, organic soil and jaggery in the laboratory condition. Similar findings were recorded with (Raghavendra *et al.*, 2014) and (Devkumari *et al.*, 2014) who also prepared the panchgavya and jeevamrit from different by product of cow. Since the organic inputs such as panchgavya, jeevamrit and FYM are an enriched source of the microbial diversity, sufficient amount of microorganisms were isolated from the panchgavya, jeevamrit and FYM by serial dilution method on nutrient agar medium. The results of present study are in agreement with those of (Sharma and Singh, 2015) who isolated the maximum number of bacterial population was ranged from 60.5×10^4 to 175×10^4 cfu/ml and minimum concentration was exhibited in dilution 10^{-6} which ranged from 23.5×10^{-6} to 80.5×10^{-6} . The selected isolates were gram positive or negative rods, grow best at a temperature of $35 \pm 2^\circ\text{C}$, 24–48 hrs of incubation and at neutral pH on the basis of morphological, physiological and biochemical characterization. Similar results were observed with (Sharma & Singh, 2015 and Teo & Teoh, 2011) who studied a mixture of both gram positive and negative bacteria and rods and some bacilli species from panchgavya, jeevamrit and FYM. On the basis of morphological, physiological and biochemical characterization by Bergey's Manual of systematic bacteriology the selected bacterial isolates were tentatively identified as *Bacillus* sp. and *Pseudomonas* sp.

Conclusion

It is concluded from the present investigation that preparation of panchgavya, jeevamrit was very easy and cheap process. Being a richest source of microbial diversity, isolation

Comment [A3]: Must be documented

and characterization of bacterial isolates from these sources are new and less exploited. The bacterial isolates were isolated from panchgavya, jeevamrit and FYM by standard method and were further screened for various plant growth promoting activities. Only three bacterial isolates were selected on the basis of maximum plant growth promoting traits and tentatively identified as *Bacillus* sp. and *Pseudomonas* sp. and preserved for further use in net house and field trial on various crops. The selected bacterial isolates can be used as biofertilizer for enhanced production of crop and to maintain the fertility of soil by natural means. The use of bacterial isolates may minimize the use of chemical fertilizer in near future.

References

- Bakker A W and Schippers B. 1987. Microbial cyanide production in the rhizosphere to potato yield reduction and *Pseudomonas* sp. mediated plant growth stimulation. *Soil Biology and Biochemistry* 19:451-57
- Bhardwaj S, Kaushal R, Kaushal M and Bhardwaj K K. 2018. Integrated nutrient management for improved cauliflower yield and soil health. *International Journal of Vegetable Science* 24:29-42.
- Devakumar N, Rao G G E, Shubha S, Imrankhan, Nagaraj and Gowda S B. 2008. Activates of organic farming research centre. Navile, Shimoga, University of Agriculture Sciences Bangalore, 12 p.
- Devakumar N, Rao G G E and Shuba S. 2011. Evaluation of locally available media for the growth and development of nitrogen fixing micro-organisms. Proceedings of the 3rd scientific conference of ISOFAR Organic are life- knowledge for tomorrow, held on 28th September-01 October 2011, Korea. PP 504-509.
- Devakumar N, Shubha S, Gouder SB and Rao GGE. 2014. Microbial analytical studies of traditional organic preparations beejamrutha and jeevamrutha. In: Building Organic Bridges (Rahmann G and Aksoy U eds). Proceedings of the 4th ISOFAR Scientific Conference, Istanbul, Turkey. pp. 639-642.
- Edi-Premono, Moawad M A and Vleck P L G. 1996. Effect of phosphate solubilizing *Pseudomonas putida* on the growth of maize and its survival in the rhizosphere. *Indonesian Journal of Crop Science* 11: 13-23.
- Gomez K A and Gomez AA. 1984. Statistical procedures for agricultural research (2 ed.). John Wiley and sons, New York, 680p.
- Gordon S A and Palleg L G. 1957. Quantitative measurement of IAA. *Plant Physiology* 10:37- 38.

Gunwantrao B B and Patil C R. 2015. Bacterial succession during panchagavya making as revealed by DGGE analysis. *Journal of Pure and Applied Microbiology*. 9(4): 3005-3013

Pikovskaya R I. 1948. Mobilization of phosphorus in soil in connection with the vital activity of some microbial species. *Mikrobiologiya* 17:362-370.

Raghavendra K V, Gowthami R, Shashank R, Harish K S. 2014. Panchagavya in Organic Crop Production. *Popular Kheti*: 2(2) ISSN: 2321-0001.

Ramesh P, Singh M and Subbarao A. 2005. Organic farming: its relevance to the Indian context. *Current Science* 88:561-569.

Schwyn B and Neilands JB. 1987. Universal chemical assay for the detection and determination of siderophore. *Analytical Biochemistry* 28:751-759.


Sharma B and Singh B. 2015. Isolation and characterization of bacteria from cow dung of desi cow breed on different morpho-biochemical parameters in Dehradun, Uttarakhand, India. *International Journal of Advances in Pharmacy, Biology and Chemistry* 4:276-81.

Sreenivasa M N, Nagaraj M Naik and Bhat S N. 2010. Beejamruth: A source for beneficial bacteria. *Karnataka Journal of Agriculture Science* 17(3):72-77.

Teo K C and Teoh S M. 2011. Preliminary biological screening of microbes isolated from cow dung in Kampar. *African Journal of Biotechnology* 10(9): 1640-1645.

Vincent J M. 1947. Distortion of fungal hyphae in the presence of certain inhibitors. *Nature* 150: 850-67.

Wang R, Zhang H, Sun L, Qi G, Chen S and Zhao X. 2017. Microbial community composition is related to soil biological and chemical properties and bacterial wilt outbreak. *Scientific Reports* 7:343-353



[Blank label box]




[Blank label box]



[Blank label box]

[Blank label box]



[Blank label box]

[Blank label box]




[Blank label box]







[Blank label box]


[Blank label box]




[Blank label box]

[Blank label box]






Effect of different temperature on the growth of selected bacterial isolates



Effect of different pH on the growth of bacterial isolates



Comment [A4]: Must be performed

Effect of incubation period on the growth of selected bacterial isolates

UNDER PEER REVIEW