

Brief note on hooded ~~Hooded~~ grasshopper *Teratodes monticollis* (Gray, 1832) ~~morphometric~~ and intraspecific colour variation from Bonai forest division, Odisha, India

ABSTRACT

Hooded grasshopper belongs to order Orthoptera and family Acrididae and are native to India and Sri Lanka. Hooded grasshopper ~~*Teratodes monticollis*~~ was described as *Gryllus monticollis* by English Zoologist George Robert Gray in 1832. The distribution of this species of hooded grasshopper has been recorded from different parts of Burma, Ceylon and India. The contemporary reports about records of this species from various states of India is very limited. From our current study area of Bonai forest division we observed *Teratodes monticollis* (Gray, 1832) and its intraspecific colour variation from Sundargarh district of Odisha, India.

Keywords: ~~Distribution~~, Phytophagous; ~~morphology-Component~~; Intraspecific.

INTRODUCTION

Across the globe, grasshoppers are among the most economically important insects and are an important component of grassland and forest ecosystems. Grasshoppers are one of the earliest living groups of munching phytophagous insects, dating back to the early Triassic period, around 250 million years ago. In the past decades, grasshoppers were treated as a nuisance and believed to be the reason of ~~draught-drought~~ and famine because of their overpopulation and phytophagic nature, especially crops and plants. Due to lack of studies, less data are available in the literatures and previously more importance ~~waswere~~ given to larger majestic ~~animals, animals~~; there ~~waswere~~ not much ideas ~~we-have~~ about the importance of insects in our ecosystem. For the past few years, studies ~~has-have~~ been done on different insects including grasshoppers, and their importance to balance our ecosystem as a potent pollinator. Due to urbanization, we are losing forest lands, agricultural lands along with floral and faunal species, among them grasshoppers are also very crucial for our environment. The genus *Teratodes* (Brunner von Wattenwyl, 1893) ~~coming-comes~~ under the family Acrididae, and this genus is comprised of three species; *brachypterus* (Carl, 1916), *monticollis* (Gray, 1832) and *folius* (Herbst, 1803).

MATERIALS AND METHODS

The investigated specimen in our study was observed with intraspecific colour variation and identified in the Bonai forest division (21.75° N latitude and 84.97° E longitude) of Sundargarh district, Odisha, India. The studied specimen was spotted during a preliminary survey in the year 2022. This targeted specimen was investigated and closely observed in its natural habitat for 24 days (31st May 2022 to 23rd June 2022), during morning hours (7:00 am to 12:00 pm) and evening hours (4:30 pm to 7:00 pm). The main aim of our observation was to document its morphological features, habits, and habitat. Identification of *T. monticollis* (Gray, 1832) was confirmed after going through literatures to observe the characteristics to identify the specimen and to be confirmed about its distribution and the morphological traits and figures mentioned in Kirby [1] and Willems [2].

Comment [h1]: Avoid the use of 1st person pronoun

Comment [h2]: Please Include the major finding of the study, this is very general.

Comment [h3]: Avoid the use of 1st person pronoun

Comment [h4]: Avoid the use of 1st person pronoun

RESULT AND DISCUSSIONS

During the present investigation we observed *T. monticollis* (Gray, 1832) in Odisha, India, and its's intraspecific colour variation i.e., green and brown coloured, inside the deep Sal Forest areas, small shrub and open dry areas of Bonai forest division, Sundargarh.

Comment [h5]: Avoid the use of 1st person pronoun

MATERIAL EXAMINED

1 ♂ (green) and 1 ♀ (brown) Bonai (), Odisha State, India on small plant and tree- Jungle, 31.05.2022, Coll. Pradhan. I., Sanjeet Kumar, 1 ♀ (green) Bonai (), Odisha State, India on forest floor -Jungle, 23.06.2022, Coll. Sanjeet Kumar, Pradhan. I (Figs. 1 & 2).




Figure: 1 *Teratodes monticollis* (Gray, 1832) in Bonai Forest Division, Odisha, India **a.** lateral view of Male specimen **b.** lateral view of female specimen **c.** anterior view of female specimen **d.** anterior view of male specimen.

Table 1: Morphometric data of *T. monticollis* (in mm)

Comment [h6]: Table 1. Is not cited in the main text. Please cite it.

Morphometric analysis of <i>T. monticollis</i> (in mm) ()			
SI No.	Body part	Male (mm)	Female (mm)
1.	Length of body	30.3	51.6
2.	Length of head	6.2	13.7
3.	Length of antenna	8.9	16.8
4.	Hight of pronotum	11.4	23.6
5.	Length of pronotum	15.8	32
6.	Length of wing	22.9	26.7
7.	Length of fore limb	10.5	28.9
8.	Length of mid limb	10.4	28.8
9.	Length of hind limb	33.2	55.3
10.	Length of femur (hind limb)	11.1	22.3
11.	Length of tibia (hind limb)	13.9	25.4




Figure 2: a & b *Teratodes monticollis* (Gray, 1832) female specimens of green and brown variant.

DIAGNOSTIC FEATURES

This species has a broad head, medium-sized body, a pronotum (much compressed forming a crest, the front arched above its head at a point, the middle portion forming a higher crest, denticulate and covering head anteriorly, half of the abdominal length posteriorly and never cross transverse sulci, the frontal ridge is sulcate, short prosternal process, the mesosternal interspace is open, straight and pointed apex.), lateral carinae absent, fastigium of vertex is rounded, one pair of filiform antennae which are shorter than the combined length of pronotum and head, antenna are wide apart from each other, out of three pair of limbs, fore limbs, and mid limbs are shorter and hind limbs are longer. Hind femurs have small pointy spines on the upper carinae and on the hind tibia, 8-10 comparatively longer and strong spines are present dorsally. Tegmina is Tegmen is opaque, shorter or longer than the abdomen and the wings are hyaline. The green variant specimen is uniformly green and the brown variant one is uniformly brown. Antennae are reddish at-in the middle, yellowish at tips in both the variants while pale green at the basal part of the green variant and pale brown at the

Comment [h7]: It refers to?

basal part of brown variant. The line of the whole crest of the pronotum is yellow in both males and females of the green variant, and the ~~whole-entire~~ crest of the pronotum of the brown variant is brown. Two arches of each knee, the upper carinae of the femur of hind limbs, and the lower carinae and spines of the hind tibia are reddish and yellowish in colour in the green variant and brownish and reddish in the brown variant. The Tegmina of the green variant is evenly green with pale and yellow spots whereas brown variant has evenly the brown tegmina with pale and brown spots and tegmina of both the-variants are reticulated and thick. Wings are pale greenish in the green variant and pale brown in the brown variant. Around the crest of the pronotum and out upper carina of the femurs pale and yellow spots are observed in the green variant and pale and brown in the brown variant. The ~~Abdomen~~ abdomen area of female specimen is raised upward in both the-variants which differentiate them from male specimens and wings of the male specimen are longer than the female specimen in comparison to their body size.

Comment [h8]: Reference is required

HABITAT

Teratodes monticollis can be found near or on shrubs and trees, and dry open habitats.

DISTRIBUTION

T. monticollis is native to Burma, Ceylon and India. In India this hooded grasshopper has been reported from; Jharkhand, Bihar, Chhattisgarh, Bengaluru, Rajasthan, Karnataka, Kerala, Tamil Nadu, Telangana, Andhra Pradesh, Maharashtra, Madhya Pradesh, Uttar Pradesh, Gujarat, and Odisha.

CONCLUSIONS

Teratodes monticollis (Gray, 1832) and its intraspecific colour variation ~~is~~are formally new to the Bonai forest division. This study provides the first ever photographic evidence to confirm the occurrence, habit, and habitat study of *T. monticollis* from the Bonai Forest Division, Odisha, India. The green variant of *T. monticollis* has been observed and studied ~~from-in~~ India's various states, Jharkhand, Bihar, Chhattisgarh, Bengaluru, Rajasthan, Karnataka, Kerala, Tamil Nadu, Telangana, Andhra Pradesh, Maharashtra, Madhya Pradesh, Uttar Pradesh, Gujarat in addition to Odisha., But, the brown variant is reported for the first time from India in the State of Odisha. *T. monticollis* with ~~its-its~~ two variants are very rare and less in numbers than other grasshoppers. ~~We~~ observed it from deep Sal Forest areas of Bonai forest division. It is a major defoliator of sandalwood, and teak and also feeds on *Calotropis procera* and *Ziziphus nummularis*. There is very less research ~~that~~ has been done about the behavioural, genetic, and ecological aspects of this hooded grasshopper, so there is more need ~~of-for~~ studies.

Comment [h9]: Avoid the use of 1st person pronoun

REFERENCES

1. Kirby WF. Fauna of British India including Ceylon and Burma. Orthoptera (Acrididae). 1914;234-235.
2. Willemsse C. Tijdschrift voor Entomologie. 1930; 73:209.

Comment [h10]: The reference style should be the same, be consistent

3. Modi, A., & Raval, K. (n.d.). *Uttar Pradesh journal of zoology new report on the occurrence of the hooded grasshopper-Teratodes monticollis (Gray 1832) (Insecta: Orthoptera: Acrididae) and the observation of the mating behaviour from Gujarat, India.* <https://www.researchgate.net/publication/354895936>

Comment [h11]: It is not cited in the main body

4. Gogoi, K., Das, R., & Yonzon, R. (2012). *Intraspecific Colour Variation in Orchid Species of Assam, India* (Vol. 13, Issue 3).

Comment [h12]: It is not cited in the main body

5. Gupta SK. On a collection of Orthoptera (Insecta) fauna from Gurughasidas National Park Koriya and Surguja districts Chhattisgarh India. *International Journal of Global Science Research*. 2015;2(3):187-198.

Comment [h13]: It is not cited in the main body

6. *Teratodes monticollis* (G.R. Gray, 1832) in GBIF Secretariat (2021), GBIF Backbone Taxonomy. Checklist dataset 16. Available: <https://doi.org/10.15468/39omei> accessed via GBIF.org on 2021-08-20.

7.V, M. S., & N, N. K. (n.d.). *Studies on grasshoppers (orthoptera) in Tilari forest, Chandgad, Kolhapur district of Maharashtra (India).* [Http://www.recentscientific.com](http://www.recentscientific.com)

UNDER PEER REVIEW