

# Impact of different irrigation techniques on wheat crop micro-climate

## ABSTRACT

Aim of this experiment is to study the impact of irrigation techniques on microclimate of wheat crop in terms of temperature and humidity. Randomized block design is used on the wheat crop (*Triticum aestivum* L.) variety Karan Narendra (DBW 222) at research farm of ICAR-CSSRI, Karnal during the Rabi 2021-22. In conventional practice treatment, higher temperature fluctuation was observed and was 0.5-1°C cooler as compared to other plots. Drip irrigation treatment showed most stable crop microclimate in terms of canopy temperature and humidity dynamics. A sharp rise in temperature at maturity cause dryness in crop and leads to lower productivity. Except at physiological maturity, canopy was cooler than the ambient air, while relative humidity was higher both during morning and evening hour.

Keywords: *Wheat, canopy, temperature, relative humidity*

## 1. INTRODUCTION

Wheat (*Triticum aestivum* L.) crop occupies an important position in Indian agricultural sector and overall financial system. It is the most important staple food of about two billion people (36% of the world population). In India, its cultivation is spread between 10°N to 37°N and cultivated under a wide range of agro-climatic conditions from Kashmir and another mountainous region to moist soils in the deltaic coastal areas and arid soils in Rajasthan (Singh and Supriya, 2017). It is the second staple food crop of India, cultivated in about 30.60 million hectares with a production of 102.28 million tonnes and productivity of 32.24 quintals per hectare. Climate variables viz. temperature, light, humidity, and rainfall in optimum conditions, exploit the full genetic potential of a particular variety by providing a suitable environment for crops and accounts for more than 20 percent of the variation in wheat yield (Dahiya *et al.*, 2020). Canopy temperatures provide a more realistic measure of microclimatic conditions and real conditions of crop stress exhibited in the field. Working on north China plain, Liu and Kang, (2006) also reported that microclimatic changes in winter wheat crops occurred due to different irrigation practices at north China plain. Canopy temperature depression had an important role to explore the physiological basis of the grain yield of wheat (Bahar *et al.*, 2008). An increase in mean aerial temperature by 1°C resulted in a reduction of the vegetative period by about 3 days and reduction in grain yield of about 7 q ha<sup>-1</sup>, reproductive period by 4.6 days (Kalra *et al.*, 2008), whereas temperature above 18.2 to 18.8°C cause a drastic reduction in grain yield by 23.2 to 42.2 per cent over the highest yield that achieved at the mean temperature of 18.2°C during reproductive phase of the crop (Solanki *et al.*, 2017). Keeping the foremost mentioned points in view, microclimate of wheat crop was studied under different irrigation techniques and tillage practices.

## 2. METHODOLOGY

### 2.1 Experimental site

The investigation was conducted during Rabi 2021-22 at the Research Farm of ICAR-Central Soil Salinity Research Institute, Karnal, situated at 29°43' N latitude, 76°58' E longitude and altitude of 245 meters.

**Comment [H1]:** Please mention the lowest & highest canopy temp & RH observed during the study period.

**Comment [H2]:** It looks very general statement. Kindly mention the impact of crop micro climate on crop yield under selected treatments or irrigation methods.

**Comment [H3]:** Quote the reference.

**Comment [H4]:** It looks sentence is incomplete. Please check it.

**Comment [H5]:** Quote the reference please.

## 2.2 Climatic conditions of experiment site

The climate of Karnal is a typical semi-arid climate having hot and dry summer with desiccating winds. Maximum temperature is around 48°C during the summer months of May to June and temperature below freezing point accompanied by frost in the winter months of December and January are common in Karnal. Fig.1. depicts the weather conditions prevailed during the season at experimental area.




Fig.1. Mean weekly values of weather parameters during Rabi (2021-22)

Temperature (°C), relative humidity (%), bright sunshine hours (hrs), wind speed (kmph) (left margin); rainfall (mm), evaporation (mm) (right margin)

Comment [H6]: Sunshine hours data are depicted in right margin in fig. Kindly check it. Change to left margin.

## 2.3 Experimental layout

The experiment was conducted in randomized block design (RBD) with five replications. Treatments consist of conventionally tilled wheat with 6.5 cm surface irrigation provided at critical phenological stages (CRI, tillering, jointing, dough, and flowering) without straw mulch i.e farmers practice (T1); zero-tilled wheat with straw mulch and drip irrigation plot (T2); zero-tilled wheat with straw mulch and surface irrigation at critical phenological stages plot (T3); zero-tilled wheat with straw mulch and mini-sprinkler irrigation plot (T4). Full residue (1/3) of rice crop on surface was left over on surface as mulch. Karan Narendra (DBW 222) wheat variety was sown at recommended inter-row spacing of 22.5 cm. Total amount of irrigation applied in T1 is 26 cm, in T2 is 19.8 cm, T3 is 26 cm and T4 is 20.10 cm with rainfall of 12.42 cm.

Comment [H7]: Mention the date of sowing.

## 2.4 Observation recorded

Micro-meteorological observations (canopy temperature and relative humidity) were recorded at 20 cm height from ground at 0700 (morning) and 1400 hours (afternoon) daily by a digital thermo-hygrometer (HTC-2). It was installed in the middle of every treatment to ensure the prevention of any outer disturbance.

Comment [H8]: Is it standard timings to take micro climate observation? Or why have taken only these timings? As per reviewer knowledge standard timings are 0800 & 1400 hrs.


## 3. RESULTS AND DISCUSSION

### 3.1 Influence of irrigation practices on canopy maximum temperature

Canopy maximum temperature at 1400 hours (afternoon) among various treatments during the season is depicted in Fig.2 along with maximum temperature of ambient air

Comment [H9]: What was the maximum temperature of ambient air?


temperature. Highest canopy maximum was recorded in zero tillage mini-sprinkler irrigation treatment (T4) (47.16°C) followed by zero tillage surface irrigation treatment (T3) (46.26°C) and zero tillage drip irrigation treatment plot (T2) (43.07°C), whereas, lowest temperature was recorded in conventional tillage surface irrigation treatment plot (T1) (39.98°C) because of high transpiration cooling and heavy irrigation.



**Fig.2. Canopy Maximum Temperature (1400 hours) during Rabi (2021-22)**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen temperature

Canopy air temperature (Fig.3) of all the treatments was higher than the ambient air temperature during winter period (tillering and booting stage), at par during anthesis and higher during warm period (late February to April) at milking and physiological maturity stages due to heat wave. It was observed that maximum canopy temperature in afternoon was lower than the ambient air temperature up to milking stage and thereafter it was higher. When canopy temperatures are high, too much energy is lost through the process of transpiration by the plants, and the reduced residual energy results in poorer grain formation and lower yields.




**Fig.3. Canopy maximum temperature at various growth stages of wheat**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen temperature

### 3.2 Influence of irrigation practices on canopy minimum temperature


Canopy minimum temperature for morning among various treatments during the season is depicted in Fig.4 along with minimum temperature of air ambient minimum temperature. In morning, during winter period (post-joining to anthesis), minimum canopy temperature was higher than the ambient minimum air temperature (Fig.4). As the crop matured ambient air minimum temperature at morning was higher than the canopy minimum temperature. During jointing to anthesis stage lowest mean minimum morning temperature was recorded in conventional tillage T1 (4.1°C), while during anthesis to crop maturity period it was recorded in zero tillage mini-sprinkler irrigation treatment (T4) (15.43°C). Presence of mulch material over T2, T3 and T4 helped to trap more heat in canopy and buffered the canopy conditions which prevented fall in minimum temperature during winter period of crop growth cycle.

Comment [H10]: Mention the correct unit.



**Fig.4. Canopy Minimum Temperature (0700 hours) during Rabi (2021-22)**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen temperature



**Fig. 5. Canopy minimum temperature at various growth stages of wheat**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen temperature


Comparatively higher canopy temperature was recorded in T4 as it drains out the surplus water quickly because of higher infiltration rate ( $1.35 \text{ cm hr}^{-1}$ ), too much energy and water losses because of evapotranspiration and canopy interception, followed by T3, T2 and least in T1 because of high transpiration cooling and heavy irrigation. Due to more canopy cover and good crop stand, T1 was able to maintain a lower canopy temperature was also observed by Dhaliwal *et al.*, 2019 in his work. Presence of mulch over T2, T3 and T4 trapped more heat in canopy and raised the temperature during winter period. Terminal heat stress was more pronounced in T4 and T3 because of higher solar radiation interception due to low canopy covers, mulch, higher soil temperature and moisture due to sensible heat transfer from soil to crop canopy via canopy interception and conduction, which in later crop growth stages increased crop canopy temperature. The effect of mulch was observed up to 100-110 days after sowing. A sharp rise in canopy temperature of T4 accompanied by hot winds during the grain growth period caused forced maturity of the crop resulting in the production of immature and shriveled grains and reduced production. In T1, higher temperature fluctuation was observed and was  $0.5\text{-}1^\circ\text{C}$  cooler as compared to other plots.

Comment [H11]: Mention the place of study.

### 3.3 Influence of irrigation practices on canopy relative humidity

Crop canopy was warmer and retains higher relative humidity (>20-30%) than ambient air. Relative humidity at 0700 hours (RH-I) was higher in T1 (99.7%) due to which leaf temperature of crop in T1 was lowest ( $24.9^\circ\text{C}$ ) (Fig.6). Relative humidity at 1400 hours (RH-II) was higher than air relative humidity throughout the crop growth period (Fig.7). Mulching aids in maintaining a good amount of humidity near soil surface. In T4, RH falls sharply because of high canopy temperature ( $44\text{-}46^\circ\text{C}$ ), heat wave incidence, lower crop coefficient ( $K_c$ ) (0.40) and amount of irrigation.


Comment [H12]: Low amount or high amount?



**Fig.6. Canopy Relative Humidity at 0700 hours**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen temperature

**Comment [H13]:** Is it temperature or RH? Please check it



**Fig.7. Canopy Relative Humidity at 1400 hours**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen temperature


**Comment [H14]:** Please check it. Whether it is temp or RH.

Highest mean canopy relative humidity (RH-I) during morning hours was recorded at booting stage (99%), while lowest was observed at harvest (40%) depicted in Fig.8. From jointing to milking stage there was a little difference in RH-I among the different treatments. In these growth stages RH-I was higher in conventionally surface irrigation method treatment and least in sprinkler irrigation system treatment (T4) (54%). Canopy relative humidity for afternoon (RH-II) among various treatments is depicted in Fig.9 along with relative humidity of air at same time. Maximum mean canopy relative humidity during afternoon hours was recorded during heading stage because of high rainfall, and was observed highest in T2 (98%) followed by T4 (98%), T3 (98%) and T1 (94%), while lowest was recorded during

**Comment [H15]:** There is no changes among three treatment. Please check it.


harvest in T1 (13%), followed by T4 (31%), T3 (25%) and T2 (34%). It was observed that in mini-sprinkler irrigation treatment relative humidity fall sharply in later stages of the crop development.

**Comment [H16]:** Not in lowest order. Please correct it.



**Fig.8. Canopy Relative Humidity at 0700hrs of growth stages of wheat crop**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen RH



**Fig.9. Canopy Relative Humidity at 1400hrs of growth stages of wheat crop**

Where, SMW- Standard meteorological weeks, T1- conventional irrigation plot, T2- drip irrigation plot, T3- surface irrigation plot with straw mulch, T4- mini-sprinkler irrigation plot, SVS- Stevenson's screen RH

### 3.4 Influence of irrigation practices on Stomata conductance, Transpiration and Photosynthetic rate

At maximum vegetative growth, transpiration and photosynthetic rates depicted in Table1 were significantly higher in T1 due to abundance water availability and lower relative humidity of canopy, while lesser stomata conductance was responsible for minimum water losses, which results in lavish growth of crop under the plot.

**Table1. Influence of different irrigation practices on various physiological parameters at maximum vegetative growth in wheat**

Treatments	Transpiration rate (mmol m <sup>-2</sup> s <sup>-1</sup> )	Photosynthetic rate (μmol m <sup>-2</sup> s <sup>-1</sup> )	Stomata conductance (mol m <sup>-2</sup> s <sup>-1</sup> )
T1	16.23 <sup>A</sup>	0.33 <sup>A</sup>	1.11 <sup>C</sup>
T2	13.58 <sup>B</sup>	0.26 <sup>B</sup>	1.20 <sup>B</sup>
T3	12.99 <sup>B</sup>	0.21 <sup>C</sup>	1.30 <sup>A</sup>
T4	13.27 <sup>B</sup>	0.22 <sup>C</sup>	1.28 <sup>A</sup>
General Mean	14.02	0.26	1.22
p-Value	0.0167	<.0001	0.0007
CV (%)	6.56	4.67	2.42
SE(d)	0.751	0.01	0.024
LSD at 5%	1.8383	0.0238	0.0592

Experimental area faced high temperature (around >5°C) than normal during grain filling stage. Heat wave of 7 days stroked the crop which desiccated the crop and soil, reduced grain moisture and yield reduction of around 20-25%. Grain yield of T1 (47.35 qha<sup>-1</sup>) was higher followed by T2 (45.92 qha<sup>-1</sup>), T4 (43.66 qha<sup>-1</sup>) and T3 (41.67 qha<sup>-1</sup>), whereas, the harvest index significantly varied under different treatments. T2 had significantly higher harvest index (38.30%) as compared to T3 (37.50%), T4 (37.46%) and T1 had lowest (36.34%).

#### 4. CONCLUSION

Field microclimate was affected by different types of irrigation methods system not only during the period of irrigation but also during all irrigation intervals. Drip irrigation plot (T2) showed most stable crop microclimate in terms of soil and canopy temperature dynamics. A strong association of canopy temperature and humidity is seen with the method of irrigation applied. Deficit irrigation application in T4 with higher evapotranspiration losses and infiltration rates in zero tilled plots (T4 and T3) faced heat stress at maturity. T4 recorded to have highest canopy temperature at 1400 hours because of deficit irrigation and higher solar radiation interception in canopy, while T1 has lowest canopy temperature because of heavy irrigations and dense canopy cover. Crop canopy was warmer and retains higher relative humidity (>20-30%) than ambient air. Mulches helped in warming the air near soil surface, resulting in higher (2-5 °C) canopy temperature.

**Comment [H17]:** This statement indicates that, farmers practice T1 has more grain yield than other treatments. That means farmers practice is better than other treatments? Kindly justify how?

**Comment [H18]:** How canopy temperature & RH effected on grain yield? Write a conclusion on effect of crop micro climate on grain yield.

#### REFERENCES

Bahar B, Yildirim M, Barutcular C, and Ibrahim GENC. Effect of canopy temperature depression on grain yield and yield components in bread and durum wheat. Notulae Botanicae Horti Agrobotanici Cluj-Napoca.2008;36(1): 34-37.

- Dahiya P, Mishra EP, Mehra B, Khichar ML and Niwas R, 2020. Impact of Climatic Variability on Wheat Yield in Indian Subcontinent. *Journal of Agrometeorology*. 2020; Volume22(1): 141-148
- Dhaliwal LK, Buttar GS, Kingra PK, Singh S and Kaur S. Effect of mulching, row direction and spacing on microclimate and wheat yield at Ludhiana. *Journal of Agrometeorology*, 2019;21(1): 42-45.
- Kalra N, Chakraborty D, Sharma A, Rai HK, Jolly M, Chander S, Kumar PR, Bhadraray S, Barman D, Mittal RB and Lal M. Effect of increasing temperature on yield of some winter crops in northwest India. *Current science* 2008; 82-88.
- Liu HJ and Kang Y. Effect of sprinkler irrigation on microclimate in the winter wheat field in the North China Plain. *Agricultural Water Management*. 2006;84(1-2): 3-19.
- Singh M and Supriya K. Growth Rate and Trend Analysis of Wheat Crop in Uttar Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*, 2017;6(7): 2295-2301.
- Solanki NS, Samota SD, Chouhan BS and Nai G. Agrometeorological indices, heat use efficiency and productivity of wheat (*Triticum aestivum*) as influenced by dates of sowing and irrigation. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(3): 176-180.

UNDER PEER REVIEW