

EVALUATION OF SEED PRIMING ON GROWTH AND YIELD OF CHICKPEA (*Cicer arietinum* L.)

ABSTRACT

The present experiment entitled “Evaluation of seed priming on growth and yield of chickpea (*Cicer arietinum* L.)” was conducted during Rabi 2019 at the experimental field of Pulses Research Unit, Dr. PDKV, Akola to assess the suitable priming treatment for chickpea crop (variety: JAKI-9218). The experiment was laid out in randomized block design in three replications. The morpho-physiological traits viz. germination %, plant height, number branches, dry matter content, and number of days required to 50% flowering were taken. The morpho-physiological i.e. plant height, number of branches, dry matter content, germination % showed a significant increase in chickpea crop when seed priming was carried out with potassium nitrate for 4 hours. The number of days required to 50% flowering were recorded least in seed priming GA3 for 4 hours. Protein content was also recorded significantly higher in seed priming with potassium nitrate for 4 hours. Yield and yield attributes were also significantly increased in seed priming with potassium nitrate for 4 hours. It is concluded that seed priming with potassium nitrate for 4 hours recorded higher morpho-physiological characters over priming treatment and control.

Keywords: Seed priming, Germination, Growth parameters. GA3, Potassium nitrate

INTRODUCTION

There are about 60 domesticated grain legume species in the world (Hedley, 2001). The chickpea is fifth most important legume in the world, on the basis of total production after soybean, groundnut, beans and peas (Muzquiz and Wood, 2007). Chickpeas also known as garbanzo beans, grams or Bengal gram, Egyptian pea, chana, and chole. In India it is largest growing pulse crop. The botanical name of chickpea is *Cicer arietinum* L. The genus *Cicer* belongs to the family Leguminosae, sub family fabaceae of the tribe Viciae. The somatic chromosome number of chickpea is $2n=16$. The genus *Cicer* contains about 40 species out of which 31 are perennial and 9 are annual. Chickpea probably originated from south east Turkey. Four centers of diversity were identified in the Mediterranean, central Asia, the East and India, as well as a secondary centers of origin in Ethiopia Harris *et al.* (2012).

Seed priming is a process of controlled hydration of seeds to a level that permits pre-germination metabolic activity to proceed, but prevents actual emergence of the radicle. Benefits of seed priming includes, faster speed of emergence, enable seed to germinate and emerge even under adverse agro-climatic conditions, improve uniformity to optimize harvesting efficiency, increases shelf life of seeds, improves the resistance towards water and temperature stress, increase vigour for fast and strong plant development and increases yield potential.

Seed priming technique has been practiced in many countries including Pakistan, China, and Australia and more than thousand trials has been conducted to evaluate the performance of priming in variety of crops. Fifty three farmers tested maize seed priming in kharif season in 1996 in tribal areas of Rajasthan, Gujarat and Madhya Pradesh; India (Harris et al. 1999). Almost all farmers thought that primed crops grew more vigorously, flowered and matured earlier and produced bigger cobs and higher yield. Independent measurements on a subset of 35 trials showed a mean increase in cob weight of 6% (Harris et al. 2001).

Seed priming have various techniques for improving the performance of the growth, emergence and yield of the crop. There are some techniques which are used i.e. hydro-priming, halo-priming, osmo-priming and hormonal priming. [Please write down the goals of your experiment!!!](#)

MATERIALS AND METHODS

The details of materials used and methods employed during the experimental period have been presented in this chapter.

Design and treatments of the experiment:-The experiment was conducted in randomized block design.

Treatments were as follows:

T1:Control

T2:Hydro priming for 4 hours

T3:Priming with 0.5% urea for 4 hours

T4:Priming with 1% urea for 4 hours

T5:Priming with 0.5% Ammonium sulphate for 4 hours

T6:Priming with 1% Ammonium sulphate for 4 hours

T7:Priming with 0.5% KNO₃ for 4 hours

T8:Priming with 1% KNO₃ for 4 hours

T9:Priming with 50 ppm GA₃ for 4 hours

T10: Priming with 100 ppm GA3 for 4 hours

T11: Priming with 50 ppm chlormequat chloride for 4 hours

T12: Priming with 100 ppm chlormequat chloride for 4 hours

Comment [MDE1]: If all seed priming applications were 4 hours, so you don't need to write down "for 4 hours" in every single treatment. Just do it once at the beginning of the treatments. What control treatment actually? How many replications were used in the experiment? How many seeds were used for each experimental unit? What is the plant spacing? What statistics analysis used?

Field emergence and germination percentage

Field emergence was counted on the experimental plot. Emergence count was under taken on 8th day after sowing and was counted and noted for each plot and replications.

Growth Parameters

Plant height (cm)

Observations on plant height were recorded (cm) at 30, 60, and 90DAS. The height of the three observational plants from each treatment and replication was recorded with scale from the base of a plant to top most developing node. From these plants of each treatment, mean height was calculated and recorded.

Number of branches

The number of branches counted at 30, 60, 90 DAS and at harvest.

Dry matter production studies

For dry matter study, single plant from each treatment and replication was uprooted periodically at flowering, pod formation and at harvest. The plant samples were washed with tap water carefully in order to remove soil and dust particles adhered to it. The samples were allowed to dry at room temperature separately for 48 hrs. The plant samples were then placed in the big size brown paper perforated bags. After drying the sample on open air basis, the plant sample was finally dried in hot air oven at 70^oC up to achieving the constant weight. The average total dry matter was recorded (g/plant) genotype and replication wise.

Number of days required to 50% flowering

The day on which 50% plants from the plot found bloomed was noted and recorded. Thus, total number of days required for flowering of 50% plants, from date of sowing were counted and expressed as days to 50% flowering.

Data analysis

The analysis of variance was performed to get the significance of differences between the treatments for all the characters as per the methodology suggested by Panse and Sukhatme (1967). The 'F' test

whenever revealed significant, the critical differences were worked out at 5% level of significance for comparisons.

RESULTS AND DISCUSSION

Effect of seed priming on seed germination and germination percentage of chickpea.

The seed germination is important factor is directly affect the yield of the crop. The seed germination of chickpea was influenced significantly. The significantly higher seed germination (251 per plot) of chickpea was recorded in seed priming with 0.5 % potassium nitrate for 4 hours. The lowest seed germination of chickpea (198 per plot) was recorded in control (no seed priming). The seed germination is higher in seed priming as compared to control. Also the germination percentage of chickpea crop was influenced significantly. Significantly higher germination percentage (89.6%) was recorded in seed priming with 0.5 % potassium nitrate for 4 hours. The lowest germination percentage (70.7%) were recorded in control (no seed priming).

Formatted: Highlight

The higher germination per plot and germination percentage in seed priming with 0.5 % potassium nitrate for 4 hours might be due to stimulated hypocotyl growth, increased cell elongation resulting in faster emergence and positive effect on germination. Similar observations were recorded by Choudhary *et al.* (2008) in chickpea, Golenzani *et al.* (2008), Patilet *al* in chickpea and Selvarani *et al.* (2011) in onion.

Effect of seed priming on plant height of chickpea

The plant height (cm) was recorded 30, 60, 90 DAS and data is presented in table 2. Data pertaining to plant height of chickpea revealed that seed priming with 1.0 % potassium nitrate for 4 hours showed significantly higher (23.7 cm) plant height at 30 DAS, (35.8cm) at 60DAS and (39.7cm) at 90DAS. The lowest plant height was recorded in control (no seed priming) i.e. 17.1cm, 27.2cm, 34.4cm at 30, 60, 90DAS respectively. Plant height is important morphological parameter exhibiting direct relationship with grain yield. It is a visible measure of plant growth and is a function of leaf emergence and inter-nodal elongation. Since leaves and branches born on stem, leaf area development and biomass production shows close relationship with plant height. The effect of seed priming on plant height showed rapid increase in plant height over no seed priming (control). The enhancement of chickpea plant primed with 1% potassium nitrate might be due to increased cell division and seedling roots. Similar observations were recorded by Hamidi and Anosheh (2013) and Patilet *al.*(2018)in chickpea and Ahmadanadet *al.* (2012) in soybean.

Effect of seed priming on number of branches of chickpea

The number of branches was recorded at 30, 60, 90 DAS and at harvest of chickpea. The data is presented in table 3. The data recorded number of branches significantly higher in seed priming with 1.0% potassium nitrate ~~for 4 hours~~ (5.7) at 30DAS, (11.7) at 60DAS, (14.3) at 90DAS and (14.3) at harvest. The lowest number of branches recorded in control (no seed priming) and seed priming with normal water ~~for 4 hours~~. The treatment seed priming with 1% potassium nitrate showed significantly higher number of branches by 38.8% over control. Similar increases in number of branches of potassium primed were observed by Harris *et al.* (1999) in chickpea.

Effect of seed priming on dry matter content of chickpea

The dry matter content of chickpea was recorded at 30, 60 and 90 DAS. The data pertaining to dry matter content of chickpea reported in table 4. The dry matter content of chickpea was significantly increased due to seed priming treatment for ~~4 hours~~ over control (no seed priming). Significantly higher dry matter content of chickpea (3.2gm) at 30 DAS, (14.9gm) at 60DAS, (33.7gm) at 90DAS was recorded in seed priming with 1.0% potassium nitrate ~~for 4 hours~~. ~~Lowest~~ ~~The lowest~~ dry matter content of chickpea (2.0gm) at 30DAS, (12.1gm) at 60DAS, (27.3gm) at 90 DAS was recorded in control (no seed priming).

Plant dry matter content is the consequence of plant physiological and biological activity. Result of plant dry matter content of different seed priming treatments were found significant at 30, 60 and 90DAS. The highest dry matter content was recorded in seed priming with potassium nitrate for 4 hours. Similar observations were recorded by Ahmadavanti *et al.* (2012) in soybean and Patilet *et al.* (2018) in chickpea.

Effect of seed priming on number of days required for 50% flowering

The number of days required to attain 50% flowering was reduced with priming treatments. Seed priming with 100ppm GA₃ ~~for 4 hours~~ induced flowering (45 days) early by 5.6 days compared to ~~the~~ control (50.6 days). The early flowering may be due to higher endogenous ~~level~~ ~~levels~~ of GA₃, early completion of vegetative growth and better nourishment of plants. Similar observations in advancement of flowering were reported by Khairulet *et al.* (2015) in chickpea and Beediet *et al.* (2017) in chickpea.

CONCLUSIONS

Different seed priming treatments significantly improved morpho-physiological parameter. Amongst the eleven different seed priming treatments, seed priming with 0.5% potassium nitrate ~~for 4 hours~~ recorded significantly highest field emergence, germination percentage, and seed priming with 1% potassium nitrate ~~for 4 hours~~ was recorded significantly highest plant height, no. of branches, dry matter content.

REFERENCE

- Ahmadand, G., Soleimani, F., Saadation, B. and Pouya, M. 2012. Effect of seed priming on germination and emergence traits of soybean cultivars under salinity stress. *Int. Res J Appl. Basic Sci.* 3:234-41.
- Beedi S., Macha S., Gowda B.,Kurnalika V., Savitha AS 2017. Effect of seed priming on growth and seed yield of kabuli chickpea cv., MNK-1 (*Cicerarietinum L.*). *Int. J. of chemical studies.* 5(6):541-544.
- Choudhary, V. K., Caurasia, A. K., Gupta, A., Naseem, M. and Maiti, R. K. 2008. Effect of priming on germination and vigour of chickpea seeds. *Int. J. Agric. Environ and Biotech.* 1(4):196-198.
- Golezani, K. G., P. S. Mosaddegh and M. Valizadeh, 2008. Effect of hydropriming duration and limited irrigation on field performance of chickpea. *Res. J. of seed Sci.*1(1):34-40.
- Golenzani, G. K., Hosseinzadeh, M. A., Zehtab, S. S. and Tourchi, M. 2012. Improving field performance of aged chickpea seeds by hydro-priming under water stress. *Int. J. Plant Animal Environ. Sci.*, 2:168-176.
- Harris, D. A., Joshi, P., Khan, P., Gothkarand Sodhi, P. S. 1999. On farm seed priming in semi-arid agriculture: Development and evaluation in maize, rice and chickpea in India using participatory methods. *Exp. Agri.*, 35:15-29.
- Harris, D., B. S. Raghuwanshi, J. S. Gangwar, S. C. Singh, K. Joshi, A. Rashid and P. A. Hollington, 2001. Evaluation by farmers of on-farm seed priming in wheat in India, Nepal and Pakistan. *Exp. Agri.* 37:403-415.
- Hamidi R., Anosheh H., 2013. Comparison effect of different seed priming methods on sunflower germination and seedling growth. *J. Agro. Plant. Prod. Vol.*, 4(6):1247-50.
- KhairulMazed H. E. M., IsratJahanIrin, Mohammad KhairulKabirMollah, Syed Tarik Mahabub, Rashed Reza. 2015. Effect of sowing time and seed treatment on growth, yield and quality seed production of Chickpea (Bari Chola -6). *Int. J. of Res. and Review. Vol.* 2:398-405.
- Panse, V.G. and P.T. Sukhatme, 1967. Statistical method for Agricultural work, ICAR, New Delhi. *Indian Tab* 3 (4) : 186-188
- Patil K., Ravat A. L., Trivedi V., Hirpara A., Sasidharan N, 2018. Effect of seed priming on seed quality enhancement in chickpea (*Cicerarietinum L.*)*Int. J. of chemical studies.* 6(4):1064-1069.
- Selvarani, K. and Umarani, R. 2011. Evaluation of seed priming methods to improve seed vigour of onion (*Allium cepa* cv. Aggregatum) and carrot (*Daucuscarota*). *J. Agri. Tech.*, 7(3):875-867.

Table 1. Effect of seed priming on germination of chickpea

Treatments	Germination per plot	Germination (%)
T1: Control (no seed priming)	198	70.71
T2: Seed priming with normal water for 4 hrs	233	83.33
T3: Seed priming with 0.5% urea for 4 hrs	239	85.48
T4: Seed priming with 1.0% urea for 4 hrs	241	86.07
T5: Seed priming with 0.5% Ammonium sulphate for 4 hrs	244	87.02
T6: Seed priming with 1.0% Ammonium sulphate for 4 hrs	242	86.55
T7: Seed priming with 0.5% potassium nitrate (13:00:45) for 4 hrs	251	89.64
T8: Seed priming with 1.0% potassium nitrate (13:00:45) for 4 hrs	249	88.81
T9: Seed priming with GA3 50 ppm for 4 hrs	239	85.24
T10 Seed priming with GA3 100 ppm for 4 hrs	244	87.14
T11: Seed priming with Chlormequate chloride 50 ppm for 4 hrs	241	86.07
T12: Seed priming with Chlormequate chloride 100 ppm for 4 hrs	240	85.60
S.E. (m)+	8.46	3.02
CD@5%	24.8	8.86

Table 2. Effect of seed priming on plant height (cm) 30, 60, 90 DAS

Treatments	30 DAS	60 DAS	90 DAS
T1: Control (no seed priming)	17.1	27.2	34.4
T2: Seed priming with normal water for 4 hrs	18.8	31.7	37.3
T3: Seed priming with 0.5% urea for 4 hrs	19.4	35.3	38.4
T4: Seed priming with 1.0% urea for 4 hrs	19.3	34.2	38.5
T5: Seed priming with 0.5% Ammonium sulphate for 4 hrs	19.0	34.7	37.5
T6: Seed priming with 1.0% Ammonium sulphate for 4 hrs	19.3	33.1	38.0
T7: Seed priming with 0.5% potassium nitrate (13:00:45) for 4 hrs	21.1	35.4	39.7
T8: Seed priming with 1.0% potassium nitrate (13:00:45) for 4 hrs	23.7	35.8	39.8
T9: Seed priming with GA3 50 ppm for 4 hrs	19.2	32.3	38.2
T10 Seed priming with GA3 100 ppm for 4 hrs	20.3	35.0	38.6
T11: Seed priming with Chlormequate chloride 50 ppm for 4 hrs	18.9	29.7	37.5

Comment [MDE2]: All seed priming applications were 4 hours, so you don't have to write it on every single treatment. Just write down once in methodology.

T12: Seed priming with Chloromequate chloride 100 ppm for 4 hrs	18.2	29.8	36.6
S.E. (m)+	1.125	1.8	0.803
CD@5%	3.3	5.51	2.36

Table 3. Effect of seed priming on number of branches

Treatments	30 DAS	60 DAS	90 DAS	At harvest
T1: Control (no seed priming)	2.3	6.3	10.0	10.3
T2: Seed priming with normal water for 4 hrs	2.3	7.3	12.0	12.0
T3: Seed priming with 0.5% urea for 4 hrs	2.7	7.7	12.7	12.7
T4: Seed priming with 1.0% urea for 4 hrs	3.0	7.0	12.3	12.7
T5: Seed priming with 0.5% Ammonium sulphate for 4 hrs	3.3	8.0	12.0	12.7
T6: Seed priming with 1.0% Ammonium sulphate for 4 hrs	3.7	8.3	12.7	12.7
T7: Seed priming with 0.5% potassium nitrate (13:00:45) for 4 hrs	4.3	9.3	14.0	14.0
T8: Seed priming with 1.0% potassium nitrate (13:00:45) for 4 hrs	5.7	11.7	14.3	14.3
T9: Seed priming with GA3 50 ppm for 4 hrs	3.7	8.3	13.7	13.7
T10 Seed priming with GA3 100 ppm for 4 hrs	3.7	8.3	12.7	13.0
T11: Seed priming with Chloromequate chloride 50 ppm for 4 hrs	5.3	10.0	14.0	14.0
T12: Seed priming with Chloromequate chloride 100 ppm for 4 hrs	3.7	9.3	13.0	13.0
S.E. (m)+	0.7	1.05	1.44	0.5
CD@5%	2.14	3.08	4.25	1.6

Comment [MDE3]: Which treatments give significant result on the parameters?

Table 4. Effect of seed priming on dry matter content (g) at 30, 60, 90 DAS

Treatment	30 DAS	60 DAS	90 DAS
T1: Control (no seed priming)	2.0	12.1	27.3
T2: Seed priming with normal water for 4 hrs	2.5	12.8	27.4
T3: Seed priming with 0.5% urea for 4 hrs	2.8	12.8	28.2
T4: Seed priming with 1.0% urea for 4 hrs	2.6	13.5	30.8
T5: Seed priming with 0.5% Ammonium sulphate for 4 hrs	2.7	13.6	30.9
T6: Seed priming with 1.0% Ammonium sulphate for 4 hrs	2.6	13.8	31.5
T7: Seed priming with 0.5% potassium nitrate (13:00:45) for 4 hrs	3.0	14.5	32.5
T8: Seed priming with 1.0% potassium nitrate (13:00:45) for 4 hrs	3.2	14.9	33.7
T9: Seed priming with GA3 50 ppm for 4 hrs	2.8	13.3	29.6
T10 Seed priming with GA3 100 ppm for 4 hrs	2.6	13.5	30.3
T11: Seed priming with Chloromequate chloride 50 ppm for 4 hrs	2.8	12.9	28.8

T12: Seed priming with Chlormequate chloride 100 ppm for 4 hrs	3.0	14.3	31.8
S.E. (m)+	0.201	0.545	1.244
CD@5%	0.59	1.6	3.65

Table 5 .Effect of seed priming on number of days required for 50% flowering

Treatment	No of days required for 50% flowering
T1: Control (no seed priming)	50.6
T2: Seed priming with normal water for 4 hrs	49.7
T3: Seed priming with 0.5% urea for 4 hrs	49.0
T4: Seed priming with 1.0% urea for 4 hrs	50.0
T5: Seed priming with 0.5% Ammonium sulphate for 4 hrs	50.3
T6: Seed priming with 1.0% Ammonium sulphate for 4 hrs	45.3
T7: Seed priming with 0.5% potassium nitrate (13:00:45) for 4 hrs	46.0
T8: Seed priming with 1.0% potassium nitrate (13:00:45) for 4 hrs	45.7
T9: Seed priming with GA3 50 ppm for 4 hrs	45.0
T10 Seed priming with GA3 100 ppm for 4 hrs	45.0
T11: Seed priming with Chlormequate chloride 50 ppm for 4 hrs	50.0
T12: Seed priming with Chlormequate chloride 100 ppm for 4 hrs	50.3
S.E. (m)+	1.642
CD@5%	4.82

Snap 1: Evaluation of seed priming on Chickpea

Comment [MDE4]: What do you want to show of the pictures? What the difference between the picture?
Write more information of on the title of the picture!

UNDER PEER REVIEW

