

STUDY OF DIFFERENT RHIZOME SIZE AND TREATMENT ON TURMERIC (*Curcuma longa* L.)

ABSTRACT

The experiment was conducted at the Department of Seed Technology, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushingar during *Kharif* 2021-22. The experiment was laid out in Randomized Complete Block Design (Factorial concept) with three replications of GNT 2 variety comprising seven treatments and four rhizome sizes. The rhizome R₄ (mother rhizome) and treatment T₄ (Chiller treatment 4°C for 2 hrs.) revealed that all traits were reported. In these investigation mother rhizome given significantly maximum result for germination per cent (80.48%), plant height (64.52 cm), leaf length (36.96 cm), leaf width (13.50 cm), tillers plant⁻¹ (3.21), mother rhizome plant⁻¹ (2.32), primary finger plant⁻¹ (7.33), secondary finger plant⁻¹ (7.94), rhizome length (14.89 cm), rhizome width (15.54 cm) and rhizome weight (185.25 g) while, in case of different treatments chiller treatment (4°C for 2 hrs.) given significantly maximum result for germination per cent (85.98 %), plant height (65.35 cm), leaf length (34.51 cm), mother rhizome plant⁻¹ (2.38), Primary fingers plant⁻¹ (6.70), secondary fingers plant⁻¹ (7.95), rhizome length (14.13 cm), rhizome width (15.58 cm) and rhizome weight (187.23 g). Maximum leaf width (12.96 cm) and the number of tillers plant⁻¹ (3.24) found by Sodium hypochloride treatment (4% for 2 hrs.) T₅. Therefore, it is found that pre-sowing rhizome treatment can be useful for better turmeric crop cultivation.

KEYWORDS: Chiller treatment, *Curcuma longa*, Mother rhizome, Sodium hypochloride

INTRODUCTION

Turmeric (*Curcuma longa* L.) is a tropical perennial, underground rhizomatic crop belonging to the family *Zingiberaceae*. Is a cross-pollinated, triploid species ($2n = 3x = 63$). It is one of the ancient and sacred spices of India. It is the third most important spice crop of India next to chilli and black pepper. It is being used dates back nearly 4000 years to the Vedic culture in India as a culinary spice and dye that had a wide range of spiritual significance to the Hindu religion. Also known as “Golden Spice” as well as the “Spice of Life”. Turmeric is valued for its underground rhizome containing a yellow phenolic pigment called curcumin. Which is used as a natural coloring agent for food, cosmetics, and dye. Curcumin the main active ingredient of turmeric, functions as a medicine with anti-inflammatory, anti-mutagenic, anti-tumor, anti-bacterial, anti-oxidant, anti-fungal, anti-parasitic and detoxifying properties, due to its anti-oxidant properties, controls Alzheimer’s disease in human beings [23]. Alleppey turmeric is the world's most outstanding and demanded grade. Which is the richest source of curcumin and is extensively cultivated in Kerala.

India is the largest producer, consumer and exporter of turmeric supplying 94% of the world demand. Its crop duration is generally 7-9 months depending on the variety. In India, sowing takes place in July and harvesting commences from December to February. March-April months are the peak arrival period in the market for turmeric. Turmeric can be grown under diverse tropical conditions with temperatures ranging between 20-30°C and with an annual rainfall of 1500 mm.

The global production of turmeric is around 11 lakh tonnes per annum. India dominates the world production scenario contributing (80%) followed by China (8%), Myanmar (4%), Nigeria (3%) and Bangladesh (3%) [1]. In India, during 2021-22 about 306 ha area was covered under turmeric. The

important turmeric growing states in India is Maharashtra, Telangana, Odisha, Andhra Pradesh, Tamilnadu, Karnataka, West Bengal, and Assam. In Gujarat during 2021-22 about 4.28 ha area and production of 16.83 MT [2] was covered under turmeric. Major turmeric growing districts in Gujarat are Dahod, Navsari, Surat, Panchmahal and Mahisagar etc. India is the largest producer, consumer and exporter of turmeric that accounting for about 80 per cent, 90 per cent and 60 per cent share respectively of the world total.

Evaluating the effect of rhizome size on the growth and development of plants is very important for increasing yield in the plant species producing different sizes of seed [25, 26]. An optimum seed root in the size of a specific root crop may develop healthy seedlings and vegetative parts, which subsequently receive higher solar energy and maximize yield in ginger, potato and turmeric plants producing different sizes of propagules. The turmeric plant propagates by the mother rhizome and finger rhizome. The finger rhizomes of the species are considered to be different in size because primary finger rhizomes developed from the shoot base have secondary and tertiary finger rhizomes which are different in size due to the differences in developing time. In addition, all the primary finger rhizomes are not developed at a time from a shoot base. Therefore, it is necessary to determine the optimum size of seed rhizomes for turmeric cultivation.

To fulfill the increasing demand of people for turmeric and improve its quality, it is essential to the production of turmeric considerably. This can be achieved by bringing more area under cultivation and increasing productivity per unit area. The lack of a suitable cultivar for the particular agroclimatic condition is one of the constraints for low productivity. However other factors like high yielding varieties, nutrition, layout, planting material, spacing and time of planting influence the productivity of turmeric [20].

Materials and Methods

The experiment "Study of Different Rhizome Size and Treatment on Turmeric (*Curcuma longa* L.)" was conducted at the Department of Seed Technology, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar during *Kharif* 2021-22. Geographically, Sardarkrushinagar is positioned at 24° 19'26" North Latitude & 72° 18'53" East longitude with an altitude of 154.52 meters above mean sea level. The experiment was laid out in factorial randomized block design (FRBD) and replicated three times and a spacing of 45 x 30 cm. The variety used for the study was Gujarat Navsari Turmeric 2. Cultural practices were followed as per the recommended package of practices. The treatment comprised different sizes of rhizome based on weight as given below (Table 1, Fig.1).

The data on different characters will be analyzed on the mean value of five plants for experimental design. The characters viz., germination per cent, plant height, leaf length, leaf width, tillers per plant, mother rhizomes per plant, primary fingers per plant, secondary fingers per plant, rhizome length, rhizome width, fresh rhizome weight. The observed data were statistically analyzed by appropriate statistical procedures as suggested for Randomized complete block design (Factorial) by [18].

RESULT AND DISCUSSION

The analysis of variance (mean sum of the square) for four rhizome sizes and seven treatments for different characters are given in Table 2 and Table 3.

Growth parameters, yield and its attribute traits

The mean sum squares due to treatments and rhizome size were found to be significant for most of all the characters, indicating the existence of sufficient variability in the experimental materials uses of

different rhizome sizes and treatments use under this investigation significantly effect on germination per cent as well as different plant growth parameters. The analysis of the variance of rhizome size showed significant variation. However, maximum germination per cent was found in R₄ (80.48%) and it was at par with the R₁ (80.00%) and R₂ (79.63%), while in the case of different treatments T₄ (85.98%) give significantly higher germination per cent and it was at par with T₅ (85.84%), T₆ (83.61%) and T₁ (81.75%). The results have confirmed the reports of [24, 30, 32].

In the turmeric crop, different growth parameters are also important to obtain a good rhizome yield. In the investigation among the different sizes of rhizome, the mother rhizome gave significant results for plant height R₄ (64.52 cm) which was at par with the R₁ (62.02 cm), leaf length R₄ (36.96 cm), leaf width R₄ (13.50 cm) and the number of tillers plant⁻¹ R₄ (3.21) while different responsible traits for economic yield also found significantly due to mother rhizome maximum mother rhizome plant⁻¹ R₄ (2.32) and it was at par with the R₁ (2.24), primary rhizome plant⁻¹ R₄ (7.33) followed by the R₁ (6.67), secondary fingers per plant⁻¹ R₄ (7.94) followed the R₁ (7.32), rhizome length T₄ (14.89 cm) followed by the R₁ (14.06 cm), rhizome width R₄ (15.54 cm) followed by the R₁ (14.30 cm) and rhizome weight in R₄ (185.25g) followed by the R₁ (162.15 g). Similar results were reported by [3, 6, 13, 24, 31].

The analysis of the variance of the treatments found significant except for the plant height and rhizome length. Among the different treatments used in this experiment, the chiller treatment gave maximum plant height T₄ (62.35 cm) followed by the T₃ (60.82 cm), leaf length in T₄ (34.51 cm) and it was at par with the T₅ (34.37 cm), T₃ (33.92 cm) and T₆ (32.82 cm), mother rhizome per plant⁻¹ T₄ (2.38) and it was at par with the T₅ (2.33), primary fingers per plant⁻¹ T₄ (6.70) and it was at par with the T₅ (6.61) and T₆ (6.34), secondary fingers per plant⁻¹ T₄ (7.95) and it was at par with the T₅ (7.43), rhizome width treatment T₄ (15.58 cm) and it was at par with the T₅ (14.85 cm) and T₃ (14.59 cm). Maximum leaf width found by Sodium hypochloride treatment (4% for 2 hrs.) T₅ (12.96 cm) and it was at par with the T₄ (12.74 cm) and T₃ (12.44 cm) and number of tillers per plant⁻¹ T₅ (3.24) and it was at par with the T₄ (3.20). The results have confirmed the reports of [7, 8, 9, 17, 19, 30, 31].

CONCLUSION

It may be concluded from the above discussion that of the study showed that seedlings from different size finger rhizomes and mother rhizomes were similarly healthy. From this experiment, the mother rhizome gave the best performance in all parameters because of sufficient food reserves which probably encouraged vigorous plant growth that should have eventually translated into yield. An optimum seed root in size of a specific root crop may develop healthy seedlings and vegetative parts, which subsequently receive higher solar energy and maximize yield. The length of chilling treatment to break dormancy to improve the germination and seedling emergence. It was a best for vigor enhancement in turmeric. Almost all characters are positively improved by the application of rhizome R₄ (mother rhizome) and followed by >35 g finger rhizome, while in the case of rhizome treatment T₄ (Chiller treatment) and sodium hypochloride treatment (4% for 2 hrs.) given the significant result for germination per cent as well as in different growth attributes, Therefore, it is found that pre-sowing rhizome treatment can be useful for better turmeric crop cultivation.

ACKNOWLEDGEMENTS

Authors gratefully acknowledge Department of Seed Technology, Sardarkrushinagar Dantiwada Agricultural University, Dantiwada, Gujarat for the facilities provided

REFERENCES

- 1 Anonymous 2021^a Turmeric Outlook, <https://pitsau.edu.in/files/AgriMkt/Turmeric-August-2021>.
- 2 Anonymous 2021^b <https://dag.gujarat.gov.in/images/directorofagriculture/pdf/Fourth-Advance->

Estimate-Web.pdf. accessed on 09 July, 2022.

3. Bandara M S, Krieger K, Slinkard A E & Tanino K K 2000 Pre-plant chilling requirements for cloving of spring-planted garlic. *Canadian Journal of Plant Science*. 80(2): 379-384.
4. Butola J S & Badola H K 2008 Use of Sodium hypochlorite to enhance seedling emergence, vigor and survival of *Angelica glauca* and *Aconitum heterophyllum*. *Journal of Herbs, Spices & Medicinal Plants*. 13(4): 1-10.
5. Fisayomi O O, Adekanmi A A, Ayoade J O, Omole F O & Adekanmi Adeniyi S 2021 Effects of Low and Medium Microwave Radiation on Germination of Bean and Maize Seedlings.
6. Gracie A J, Brown P H, Burgess S W & Clark R J 2000 Rhizome dormancy and shoot growth in myoga (*Zingiber mioga* Roscoe). *Scientia horticultrae*. 84(1-2): 27-36.
7. Hailemichael G & Tesfaye K 2008 The effects of seed rhizome size on the growth, yield and economic return of ginger (*Zingiber officinale* Rosc.). *Asian Journal of Plant Sciences*.
8. Hore J K, Chattopadhyay N, Samanta M K, Murmu D & Ghanti S 2014 Effect of rhizome treatment on growth and yield of turmeric. *Journal of Crop and Weed*. 10(1): 121-125.
9. Hossain A, Ishimine Y, Akamine H & Motomura K 2005 Effects of seed rhizome size on growth and yield of turmeric (*Curcuma longa* L.). *Plant production science*. 8(1): 86-94.
10. Jage S, Malik Y S, Nehra B K & Partap P S 2000 Effect of size of seed rhizomes and plant spacing on growth and yield of turmeric (*Curcuma longa* L.). *Haryana Journal of Horticultural Sciences*. 29(3/4): 258-260.
11. Jakubowski T 2010 Effect of seed potatoes microwave stimulation on the growth and yields of potato (*Solanum tuberosum* L.). *Acta Agrophysica*. 16(2): 295-313.
12. Jakubowski T 2015 Evaluation of the impact of pre-sowing microwave stimulation of bean seeds on the germination process. *Agricultural Engineering*. 19.
13. Khan M B, Ghurchani M, Hussain M & Mahmood K 2010 Wheat seed invigoration by pre-sowing chilling treatments. *Pak. J. Bot.* 42(3): 1561-1566.
14. Kumari N, Verma S & Sharma V 2018 Manipulating tomato plant electric signaling system by microwave radiation to enhance crop productivity and nutritional value. *Computers and Electronics in Agriculture*. 154: 330-340.
15. Mahender B, Reddy P S S, Sivaram G T, Balakrishna M & Prathap B 2015 Effect of seed rhizome size and plant spacing on growth, yield and quality of ginger (*Zingiber officinale* Rosc.) under coconut cropping system. *Plant Archives*. 15(2): 769-774.
16. Monnaf M A, Rahim M A, Hossain M M A & Alam M S 2010 Effect of planting method and rhizome size on the growth and yield of ginger. *Journal of Agroforestry Environment Science*. 4(2): 73-76.
17. Padmadevi K, Jeeva Jothi L, Ponnuswami V, Durgavathi V & Rijwana Parveen I 2012 Effect of different grades of rhizomes on growth and yield of turmeric (*Curcuma longa* L.), *Asian Journal of Horticulture*. 7(2): 465-467.
18. Panse V G & Sukhatme P V 1978 "Statistical methods for agricultural workers". ICAR, New Delhi.
19. Patel B N, Patel R K & Bhalerao P P 2018 Effect of rhizome size on growth and yield of turmeric CV. GNT-1. *International Journal of Tropical Agriculture*. 36(3): 771-777.
20. Patil R B & C D Borse 1980 Effect of planting material and transplanting of seedling of different age on the yield of turmeric. *Indian Arecanut, Spices and Cocoa Journal (India)*. 4: 3-6.
21. Preetham, Ashwini, Pavan & Ravi kondle 2018 Evaluation of turmeric variety salem with different rhizome types and population levels on raised beds. *International Journal of Current Microbiology Applied Science*. 7(4): 3202-3211.
22. Radzevicius A, Sakalauskiene S, Dagys M, Simniskis R, Karkleliene R, Bobinas C & Duchovskis P 2013 The effect of strong microwave electric field radiation on vegetable seed germination and seedling growth rate. *Zemdirbyste-Agriculture*. 100(2): 179-184.
23. Risch S J & Ho C 1997 *Spices: flavor chemistry and antioxidant properties*. American chemical society.
24. Singh A 1985 Enhancement of turmeric (*Curcuma longa*) growth, Rhizome yield and its yellow pigment content by presowing chilling treatment of rhizomes.
25. Singh D K & Singh V 2003. Seed size and adventitious (nodal) roots as factors influencing the tolerance of wheat to waterlog. *Australian Journal of Agricultural Research*. 54(10): 969-977.
26. Stougaard R N & Xue Q 2004 Spring wheat seed size and seeding rate effects yield loss due to wild

- oat (*Avena fatua*) interference. *Weed Science*. 52: 133-141.
27. Tiwari T N, Kamal D, Singh R K & Prasad S R 2014 Relative efficacy of seed priming with potassium nitrate and tap water in relation to germination, invigoration, growth, nitrate assimilation and yield of pigeon pea (*Cajanus cajan* L.). *Annals of Agricultural Research*.35(2).
 28. Wang S, Wang J & Guo Y 2018 Microwave irradiation enhances the germination rate of Tartary buckwheat and the content of some compounds in its sprouts. *Polish Journal of food and nutrition sciences*. 68(3).
 29. Warkad D P, Bhingarde M T & Shelar V R 2018 Effect of microwave intensity on plant growth, seed yield, quality and biochemical parameters of rabi sorghum [*Sorghum bicolor* (L.) Moench] under rainfed conditions.
 30. Watkinson J I & Pill W G 1998 Gibberellic acid and presowing chilling increase seed germination of Indiangrass [*Sorghastrum nutans* (L.) Nash.]. *HortScience*. 33(5): 849-851.
 31. Wu C, Wang M, Dong Y, Cheng Z & Meng H 2015 Growth, bolting and yield of garlic (*Allium sativum* L.) in response to clove chilling treatment. *Scientia Horticulturae*. 194: 43-52.
 32. Yothasiri A, Somwong T, Tubngon S & Kasirawat T 1997 Effects of types and sizes of seed rhizomes on growth and yield of turmeric (*Curcuma longa* L.). *Agriculture and Natural Resources*. 31(1): 10-19.

Table 1: Treatment details	
Rhizome size (R)	Treatment (T)
R ₁ = >35 g finger rhizome	T ₁ = 4-second microwave treatment
R ₂ = >20-25 g finger rhizome	T ₂ = 8-second microwave treatment
R ₃ = ≤ 5 g finger rhizome	T ₃ = Deep freezer treatment (-20°C for 2 hrs.)
R ₄ = Mother rhizome	T ₄ = Chiller treatment (4°C for 2 hrs.)
	T ₅ = Sodium hypochloride treatment (4% for 2 hrs.)
	T ₆ = Tap water treatment for 2 hrs.
	T ₇ = Control



Fig. 1: Different grades of rhizomes

Table 2: Influence of different rhizome size on germination and its attributes traits of turmeric cv. GNT 2

Rhizome size	Germination (%)	Plant height (cm)	Leaf Length (cm)	Leaf width (cm)	Tillers plant ⁻¹	Mother rhizome plant ⁻¹	Primary fingers plant ⁻¹	Secondary fingers plant ⁻¹	length Rhizome	width Rhizome	weight Rhizome
R ₁	80.00	62.02	33.87 32.20	12.57	2.92 2.92	2.24	6.67	7.32	14.06	14.30	162.15
R ₂	79.63	60.66	29.50	11.85	2.3	2.65 2.65	5.62	5.85	12.89	13.49	153.55
R ₃	80.48 80.48	54.65	36.49 36.49	10.91	5	2.32	4.86 4.86	5.94 5.94	12.04	13.45	137.30
R ₄	1.30	64.52	36.49 36.49	10.20 10.20	0.05 0.05	0.06	0.11 0.11	0.17 0.17	10.89 10.89	15.54 15.54	184.25 184.25
C.B.S%	3.69	2.83	1.40	0.	0.	0.16	1	0.17	0.78	0.78	16
				58	15						11.79

Table 3: Influence of different treatments on germination and its attributes traits of turmeric cv. GNT 2

Treatments	Germination (%)	Plant height (cm)	Leaf Length (cm)	Leaf width (cm)	Tillers plant ⁻¹	Mother rhizome plant ⁻¹	Primary fingers plant ⁻¹	Secondary fingers plant ⁻¹	length Rhizome	width Rhizome	weight Rhizome
T ₁		60.5			2.77	2.05			13	1-1	1
T ₂	84.93	59.08		11.	2.43	2.03	6.1	6.1			
T ₃	79.44		31.	11.		2.20	5	5	1	1	1
T ₄		62.3	34.5	1		2.3	5.70	6.95	13	1	1-1
T ₅	85	60.62	34.37	1	3	2.38	6.6	7.43	14.13	15-8	1
T ₆	83	59.92		1	3	2.23	6.34	7.06	1	1	1-1
T ₇	58.68		32.	1		2.00			13-21	1-1	1
		NS	0.63	10.27	0.07	0.07	5	6.28	13	13	1-15
C.D. 5%	1		1.	0.76	0.	0.2	0.45	0.64	NS ⁶		5
	4.				19	1	1			1	15

