

SACCULAR ASCENDING AORTA ANEURYSMS: RARE CASE REPORTS AND LITTERATURE REVIEW.

ABSTRACT

The saccular aneurysm at the level of the ascending aorta is an extremely rare clinical entity, considered as an effusion of part of the aortic circumference. This entity is often asymptomatic, and the pathophysiological mechanism is still poorly understood.

We describe two unusual clinical cases of saccular aneurysms of the ascending aorta, the first revealed fortuitously following hospitalization for a hemorrhagic accident with vitamin K antagonists and the second following a complication of a severe aortic insufficiency. Our experience confirms the fundamental role of modern cardiac imaging techniques in the differential diagnosis of these unusual cases and in planning the correct surgical intervention.

This type of rare presentation of the disease is little studied in the literature. Hopefully our experience can provide more in terms of diagnosis or therapeutic management.

Keywords: Saccular aortic aneurysm, transthoracic echocardiography, Cardiac MRI, aortoplasty.

Introduction:

A true arterial aneurysm is defined as a permanent, localized dilation with an increase of at least 50% in diameter compared to the normal diameter of the artery in question. The aorta and particularly the subrenal segment are the most common sites (1).

The typical fusiform form is the most frequent 1 whereas the saccular form at the level of the ascending aorta is an extremely rare clinical entity, considered as an effusion of part of the aortic circumference (2).

The etiology of saccular aneurysms is above all a focal infection of the aortic wall, or trauma, unlike the fusiform form where degeneration of the wall secondary to atherosclerosis is predominant (1).

Saccular aneurysms tend to develop in areas of healed microscopic dissection of the aortic wall that are more susceptible to rupture and therefore require urgent surgical management (3).

We present in this article, two cases of particularly rare saccular aneurysm, especially in their mode of revelation as well as in their anatomical form.

Case Presentation

Case 1:

An 85-year-old patient, followed for atrial fibrillation (AF) complicated by ischemia cerebral vascular accident (ICD) put on vitamin K antagonist anticoagulant therapy based on (Sintrom), with a normal blood pressure, presents to the cardiology emergency room for a haemorrhagic accident in the medium abundance melena-type vitamin K antagonist with a blood assessment : Normochromic normocytic anemia at 2.4 g/dl, and an INR above 12, well tolerated haemodynamically.

Clinical examination showed deep mucocutaneous pallor, NYHA stage 2 dyspnea, blood pressure (BP) at 129/73 mmHg, and heart rate (HR) at 93 bpm, and facial paralysis.

The rest of the examination was unremarkable.

On the ECG: there was an AF at 95 bpm, without electrical repolarization disorders.

On the echocardiography, we noted a slightly dilated left ventricle located at a septal wall measuring 12mm by 10mm of good global and segmental contractility LVEF: 60%, a biauricular dilatation with a left atrium at 40cm², and a right atrium at 32cm² without intracavity thrombus, with moderate mitral insufficiency and moderate Aortic Insufficiency (Figure 1), the right ventricle is of good function, with a moderate tricuspid regurgitation(TR) estimating the systolic pulmonary pressure (SPAP) at 35+5: 40mmHg. We also note a dilated aorta with: ASCENDING aorta measuring 47 mm, Sinus 48 mm with a doubt of aortic dissection or aneurysm (Figure 2).

With this finding, a thoracic CT angiography was carried out objectifying: an aspect of saccular aneurysm at the level of the initial part of the ascending aorta (Figure 3).

The initial therapeutic management consist of hospitalization in the cardiac intensive care unit, with implementation of vitamin K perfusion and a blood transfusion to correct the present anemia.

Subsequently, and given that the patient did not present any notable complication, as well as the risk of mortality for the surgery of her saccular aneurysm was high, after multidisciplinary staff discussion: a close monitoring was indicated.




Figure 1: Transthoracic echocardiography (TTE) shows a moderate aortic regurgitation, Vena contracta:4mm.




Figure 2: A -Transthoracic echocardiography (TTE) long axis view: -shows a saccular aneurysm on the posterior wall of the aorta next to the posterior cusp aortic valve

B- TTE short axis view: saccular aneurysm (yellow arrow).




Figure 3: shows a saccular aneurysm at the level of the initial part of the ascending aorta (A: CT angiography image, B and C: 3D aortic construction showing the localization of the aneurysm).

Case 2:

A 63-year-old male patient admitted to our structure for heart valve disease control. Having a particular history, arterial hypertension for one year under enzyme converting inhibitor (ECI). Note, he is a chronic smoker since the age of 20 at the rate of 10 packets par year.

On clinical examination, the patient presents a NYHA stage II dyspnea, with angina on exertion.

Overall, he is hemodynamically stable with BP at 130/60mmHg and HR at 65 bpm. On auscultation, there was a diastolic murmur in the intense aortic focus, with no sign of heart failure.

The electrocardiogram finds a complete left bundle branch block (LBBB) with secondary repolarization disorders.

The transthoracic echocardiography shows, severe aortic regurgitation (ROS=40cm², RV=59ml, Radius Piza=8mm, TDE=26ms) on a fine tricuspid aortic valve without stenosis (Figure 4), with a dilated LV (the end-diastolic diameter at 65mm), not hypertrophied, of good function (LVEF at 62%). And an aneurysmal dilation of the thoracic aorta. The rest was unremarkable.

In front of this outcome, a thoracic CT angiography was carried out objectifying an aneurysmal thoracic aorta with the aspect of a saccular aneurysm (small size) at the initial part of the aorta (Aortic ring: 32mm, aortic sinus: 38mm, sino-tubal junction: 31mm, ascending aorta: 49mm, aortic arch: 36mm, descending aorta: 26mm). (Figure 5)

The biological assessment, both inflammatory and infectious, does not report any particularity. The cause of the aneurysm was not yet elucidated.

As concerning the management of his valve disease, and after a multidisciplinary staff, an aortic surgical replacement by BENTHALL was performed. The post-operative follow-up was simple and uncomplicated.

The patient is still under cardiological follow-up.




Figure 4: TTE: Long axis view: Severe aortic regurgitation (Vena contracta:7mm)



Figure 5: CT angiography: shows a dilated thoracic aorta with presence of a saccular aneurysm (small size) at the initial part of the ascending aorta.

Discussion:

An arterial aneurysm is a permanent dilatation characterized by a diameter 50% greater than that of the normal vessel in question (4). True aneurysm can be divided into two types: fusiform (most common) and saccular (2, 9).

Regarding to the etiology, fusiform aortic aneurysms often occur as part of degeneration of the walls secondary to atherosclerotic disease, whereas saccular aneurysms may occur following aortic infections, degeneration of a penetrating ulcer, trauma or complicating aortic surgery (4, 10)

The saccular aneurysm has a greater risk of rupture than the fusiform form, and therefore they are often treated at smaller diameter (11).

Several studies have been performed with the aim of providing an accurate radiological characterization of the size, presentation and progression of saccular aortic aneurysms.

Pathophysiologically, saccular aneurysms are secondary to localized aortic dissection. This begins with zonal degeneration and fibrosis extending through the intima and first tier of media, the area of "healed microscopic dissection" of the aortic wall. This process involves ulceration of the intima followed by the formation of an aneurysm (3,12)

Pathologically, the edges of dissected saccular aneurysms protrude, which histologically consist of the medial part of the dissected media, and are associated with a medial thinning in the center of the aneurysmal wall. These lesions differ from organized thrombi in the false lumens (3)

Classically, patients with ascending aortic aneurysms report chest pain associated with ischemic abnormalities on the electrocardiogram, particularly in cases of sinus of Valsalva aneurysms (5). However, the saccular aneurysm is rarely symptomatic. Its discovery is often fortuitous following an assessment of other manifestations, as is the case of our patients described above, or following complications.

Imaging plays an essential role in the diagnosis and follow-up of patients with aortic aneurysms. It can also point to risk or causal factors. Echocardiography is usually the initial cost-effective imaging modality that is used in the diagnosis of aneurysms, as seen in our patients it is safe and widely available (6,13)

Cross-sectional imaging such as CT and MRI are integral to the diagnosis, follow-up and management of aneurysms. 6

The growth rate of the aneurysm is considered an indicator of the risk of rupture (14) as confirmed by Dapunt et al who examined the natural history of thoracic aortic aneurysms and determined a combined mean growth rate of 4.3 mm/year.

While many biomechanical studies implicate aortic asymmetry as an indicator of risk of rupture and aneurysm growth, (8) no increased saccular aneurysm growth rates have been noted compared to fusiform aneurysms. In fact, neither aortic diameter nor other saccular aneurysm diameters measured

predicted aneurysm growth, although the former was based on a small sample size. More importantly, the heterogeneity of growth rates detected suggests that individual aneurysm characteristics may be important in determining aneurysm stability (4,15).

In the report of the subcommittee of the Joint Council of the Society for Vascular Surgery and the International Society for Cardiovascular Surgery, it was recommended that saccular aneurysms could represent an indication for surgery, regardless of their size. This recommendation is widely shared by most vascular surgeons. Despite the few studies focusing on the natural history of saccular aortic aneurysms. But the focal and asymmetrical bulge characterizing the saccular aneurysm represents an area of extreme thinning of the aortic wall with a greater risk of rupture (1)

Surgical management and technique vary by location. If an obvious acute infection is encountered at the time of repair, debridement, aortic closure and bypass surgery will only be feasible if the involvement is infrarenal. It is more practical to consider the use of a substitute such as a bovine patch or a cadaveric aorta (16).

For thoracic aortic aneurysms, it is advantageous to perform patch aortoplasty when <50% of the wall circumference is involved and there is normal tissue at the edge. This ensures sparing of the thoracic intercostal arteries and minimizes the risk of spinal cord ischemia (17).

In summary, our case reports shows that saccular aortic aneurysms are an unusual variant with a poorly understood natural history. Transthoracic echocardiography and cardiac MRI are the most efficient diagnostic tools. Treatment decision should involve the heart team. Patients with aortic regurgitation should be screen automatically for aortic aneurysms.

Conclusion:

Saccular aneurysms are often asymptomatic. Multimodality imaging and echocardiography are ideal for accurate diagnosis and treatment. Early management, particularly surgery, is the standard approach with satisfactory results.

These were rare and original cases which broaden knowledge in general cardiology, vascular pathology, and surgical modalities.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

References

- 1- Brian V. Taylor and Peter G. Kalman, Toronto, Canada. Saccular Aortic Aneurysms : Annals of Vascular Surgery. 1999; 13: 6.
- 2- Mehrpooya, and Al. Diagnostic dilemma: Saccular aneurysm or pseudoaneurysm of the ascending aorta with dissection above level of leaflets Maryam. ARYA Atherosclerosis Journal. 2012; 8 : 167-169
- 3- Hazem Aljaseem | Mohammad Bashar Izzat. Dissected saccular aneurysms of the ascending aorta. Clin Case Rep. 2019; 7:2402–2404.
- 4- A modern experience with saccular aortic aneurysms. Eric K. Shang et al. JOURNAL OF VASCULAR SURGERY. J Vasc Surg. 2013; 57: 84-88
- 5- Bruno Borrello, Francesco Nicolini, Cesare Beghi, Tiziano. Saccular ascending aorta aneurysm: report of an unusual presentation. Gherli. Interactive CardioVascular and Thoracic Surgery.2008; 7: 508–509.
- 6- M.S. Durojaye, T.O. Adeniyi, and O.A. Alagbe. MULTIPLE SACCULAR ANEURYSMS OF THE ABDOMINAL AORTA : A CASE REPORT AND SHORT REVIEW OF RISK FACTORS FOR RUPTURE ON CT SCAN. Annals of Ibadan Postgraduate Medicine. 2020; 18 : 2.
- 7- Golledge J, Muller J, Daugherty A, Norman P. Abdominal aortic aneurysm: pathogenesis and implications for management. Arterioscler Thromb Vasc Biol 2006; 26 :2605-13.
- 8- Shabir H. Shah and Al. A Case Report of Giant Ascending Aortic Aneurysm: Role of Multimodality Imaging. Aorta (Stamford) 2021; 9 :106–109.
- 9- Fuster V, O'Rourke R, Walsh R, Poole-Wilson P. Hurst's the Heart. Philadelphia, PA: Saunders; 2011: 167-17.

10- Coady MA, Rizzo JA, Hammond GL, Pierce JG, Kopf GS, Elefteriades JA. Penetrating ulcer of the thoracic aorta: What is it? How do we recognize it? How do we manage it? *J Vasc Surg* 1998; 27:1006-16.

11- Maier A, Gee MW, Reeps C, Pongratz J, Eckstein HH, Wall WA. A comparison of diameter, wall stress, and rupture potential index for abdominal aortic aneurysm rupture risk prediction. *Ann Biomed Eng* 2010; 38: 3124-34.

12- Saito K, Shimizu H, Kawai T, Yamada S, Furuse M, Fukushima K. Pathology and morphogenesis of the saccular aneurysm of the aorta and large elastic artery with healed medial dissection. *Jpn J Chest Dis*. 1994; 53 :1-10.

13- **Lee SH**, Chang SA, Jang SY *et al*. Screening for abdominal aortic aneurysm during transthoracic echocardiography in patients with significant coronary artery disease. *Yonsei Med J*. 2015; 56 :38-44.

14- Dapunt OE, Galla JD, Sadeghi AM, Lansman SL, Mezrow CK, de Asla RA, et al. The natural history of thoracic aortic aneurysms. *J Thorac Cardiovasc Surg* 1994; 107:1323-32.

15- Zafar MA, Li Y, Rizzo JA, et al. Height alone, rather than body surface area, suffices for risk estimation in ascending aortic aneurysm. *J Thorac Cardiovasc Surg* 2018; 155 :1938–1950.

16- Coady MA, Rizzo JA, Haemmond GL, Pierce JG, Kopf GS, Elefteriades JA. Penetrating ulcer of the thoracic aorta: What is it? How do we recognize it? How do we manage it? *J Vasc Surg* 1998; 27 :1006-1016.

17- Hollier LH, Taylor LM, Ochsner J. Recommended indications for operative treatment of abdominal aortic aneurysms. *J Vasc Surg* 1992; 16 :274-278.