

## Original Research Article

### **“Influence of Molybdenum and Foliar application of Boron on Growth and Yield of Cowpea (*Vigna unguiculata*)”**

#### **ABSTRACT**

The field experiment was undertaken at Crop Research Farm (CRF) during the Zaid season (march to June – 2022). The Department of Agronomy at SHUATS in Prayagraj (UP). Cowpea (*Vigna unguiculata*) is one of the important pulse crops grown mainly in Zaid season under rainfed conditions. Cowpea is considered as an important crop of Uttar Pradesh because of its short duration, high yielding and quick growing capacity. Boron has emerged as an important micronutrient in Indian agriculture in the context of the spread of its deficiency. About one third of the cultivated soils in India are deficient in Boron. Likewise, molybdenum is another micronutrient leading to poor seed yield of pulses in molybdenum deficient soils of Uttar Pradesh. This study is looking into how molybdenum and foliar boron applications affect cowpea growth and yield on soil with a sandy loam texture. The treatments consist of three doses of Molybdenum viz., 0.5 kg ha<sup>-1</sup>, 0.75 kg ha<sup>-1</sup> and 1 kg ha<sup>-1</sup> and Boron 0.5% at 3 intervals viz., 30DAS, 45DAS and 30DAS & 45DAS whose effect is observed on Cowpea (var. Gomati). The experiment was laid out in Randomized Block Design with Ten treatments replicated thrice. The treatment with application of Molybdenum 1 kg ha<sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS recorded significantly higher plant height (72.55 cm), number of branches plant<sup>-1</sup> (15.33) and plant dry weight (43.68 g), number of pods per plant (16.73), number of seeds per pod (11.67), seed index (18.38 g), seed yield (1.59 t ha<sup>-1</sup>) and Haulm yield (4.16 t ha<sup>-1</sup>) compared to other treatment combinations.

**Keywords:** Boron, Cowpea, Growth, Molybdenum, Yield.

#### **1. Introduction**

One of the significant pulse crops is cowpea (*Vigna unguiculata*), which is primarily grown in the Zaid season under rainfed conditions. The crop is autogamous, but up to 5% outcrossing has been reported in the cultivated varieties, probably due to insect activities (Badiane et al. 2014). Cowpea grains have a nutritional profile of 23.4% protein, 60.3% carbohydrates and 1.8% fat, as well as a good source of vitamins (Venkatesan et al. 2003, Srivastava et al. 2016). Cowpea is also known for producing high-quality forage. It fixes nitrogen as it is a leguminous crop and improves fertility of the soil (Kumar et al. 2015). The young leaves are used as spinach in eastern and southern Africa (Boukar et al. 2015). Cowpea [*Vigna unguiculata* (L.) Walp] (2n=22) belong to family Leguminosae also known as southern pea and black eye pea, is one of the most important vegetables. Cowpea (*Vigna unguiculata* Walp.) is an important arid legume grown for its pods, leaves and grains (Ehlers and Hall, 1997, Boukar et al. 2011). Cowpea can be grown as fodder crop, while its tender pods can be used as vegetables. However, it is predominantly grown for its dried grains (Lioi et al. 2019). Cowpea is mainly cultivated in tropical and subtropical countries such as Africa, Asia, Central and South America. Because of its

**Comment [S1]:** Enter the purpose of this research

**Comment [S2]:** should include the motivating factors for conducting this research

**Comment [S3]:** layout customize journal template

high protein content, cowpea is referred to as “vegetable meat” and have biological value on a dry weight basis (Ram, 2014). Trends in the production of pulse is adversely affected the per capita availability of pulses. In India per capita/day availability of pulses had decreased from 69g during sixties to 35g as against the FAO/WHO’s current recommendation of 80 g per day (Ali and Gupta 2012). The most important functions of boron in plants are thought to be its structural role in cell wall development, cell division, seed development and stimulation or inhibition of specific metabolic pathways for sugar transport and hormone development (Ahmed et al. 2009). Legume crops required more amount of boron compared to most field crops as boron plays vital role in proper development of reproductive organs. Its deficiency leads to sterility in plants by malformation of reproductive tissues affecting pollen germination, resulting in increased flower drop and reduced fruit set (Subasinghe et al. 2003). Foliar feeding practice would be more useful in early maturing short duration crops, where the soil applied fertilizer may not become fully available before maturity of crop (Somla Naik et al. 2018). Boron insufficiency is one of the biggest challenges to agricultural output. Its deficiency has been noted as the second-largest micronutrient restriction in crops globally, after zinc (Zn) (Ahmad et al., 2012). Molybdenum is an essential element; it is constituent of the nitrogenase enzyme and every bacterium which fixes nitrogen needs molybdenum during the fixation processes. The functions of molybdenum in leguminous plants include nitrate reduction nodulation, nitrogen fixation and general metabolism (Togay et al. 2008). Molybdenum was required for normal plant growth, reduction supply with molybdenum to the growth media decreased the activities of nitrate reductase and glutamine synthesis involved in initial steps of nitrate assimilation (Hristozkora et al. 2006). Gungula and Gerjila (2006) reported that molybdenum at 0.13 kg/ha increased plant height, number of branches, dry weight and yield parameters of cowpea plants. Molybdenum enhance cowpea root nodule efficiency, growth, minerals composition, yield quantity and quality compared with control plants. (Nadia gad et al. 2013). As a result, this study focused on the use of both Molybdenum and Boron to ensure the long-term viability of Cowpea production.

## 2. Materials and Methods

A field experiment was conducted during *Zaid season* (march – June, 2022) at Crop Research Farm (CRF), Department of Agronomy, SHUATS, Prayagraj (UP). The Crop Research Farm is situated at 25.4137° N latitude, 81.8491° E longitude and at an altitude of 98 m above mean sea level. The soil of the experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.4), medium in organic carbon (0.48%), medium in available Nitrogen (278.93 kg/ha), low in available Phosphorous (19.03 kg/ha) and medium in available Potash (238.1 kg ha<sup>-1</sup>). The treatments consist of three doses of Molybdenum viz., 0.5kg ha<sup>-1</sup>, 0.75kg ha<sup>-1</sup> and 1kg ha<sup>-1</sup> and Boron 0.5% at 3 intervals viz., 30DAS, 45DAS and 30DAS & 45DAS whose effect is observed on Cowpea (var. Gomati). The experiment was laid out in Randomized Block Design with ten treatments replicated thrice. The experiment comprising ten treatment possible combination of above factor, viz., T<sub>1</sub>: Molybdenum 0.5kg ha<sup>-1</sup> + Boron 0.5% at 30DAS T<sub>2</sub>: Molybdenum 0.5kg ha<sup>-1</sup> + Boron 0.5% at 45DAS, T<sub>3</sub>: Molybdenum 0.5kg ha<sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS, T<sub>4</sub>: Molybdenum 0.75kg ha<sup>-1</sup> + Boron 0.5% at 30DAS, T<sub>5</sub>: Molybdenum 0.75kg ha<sup>-1</sup> + Boron 0.5% at 45DAS, T<sub>6</sub>: Molybdenum 0.75kg ha<sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS, T<sub>7</sub>: Molybdenum 1kg ha<sup>-1</sup> + Boron 0.5% at 30DAS, T<sub>8</sub>: Molybdenum 1kg ha<sup>-1</sup> + Boron 0.5% at 45DAS, T<sub>9</sub>: Molybdenum 1kg ha<sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS, T<sub>10</sub>: Control (R D F: 25: 50: 25). Observations regarding growth and yield attributes was recorded during the field experiment.

**Comment [S4]:** layout customize journal template

**Comment [S5]:** you should complete a map of the research location

**Comment [S6]:** better complete the climate data

### 3. Result and Discussion

#### 3.1 Growth

According to the recorded and tabulated data pertaining to growth parameters, the significantly higher plant height (72.55 cm), maximum number of Branches per plant (15.33) and higher plant dry weight (43.68 g) was recorded in treatment with application of Molybdenum  $1\text{ kg ha}^{-1}$  + Boron 0.5% at 30DAS & 45 DAS.

Application of Molybdenum along with Boron resulted in significant increase in nutrient contents might be due to greater availability of N due to molybdenum is a constituent of enzyme nitrogenase, which is essential for the process of symbiotic nitrogen fixation. These results are in close conformity with findings of Salwinder Singh *et al.*, (2022), Shivakumar and Kumutha (2003). On the other hand, the increased plant growth may be due to Boron is considered to be essential for actively growing regions of plants, such as root tips, new leaf and bud development. These results were in close conformity with findings of Ahmad *et al.*, (2009), Ranjit Chatterjee *et al.*, (2015).

**Comment [S7]:** layout customize journal template

**Comment [S8]:** better complete with minimum data as well

#### 3.2 Yield attributes

According to the yield characteristics data that was collected and analysed at harvest, maximum number of pods per plant (16.73), maximum number of seeds per pod (11.67) and higher seed index (18.38g) was recorded in treatment with the application of Molybdenum  $1\text{ kg ha}^{-1}$  + Boron 0.5% at 30DAS & 45 DAS.

The higher performance of yield attributes might be due to the fact that, boron is proved to increase pollen viability and significant effect on pollen formation and fertilization, hence higher grains number are direct index of pollen viability, whereas the prolong nutrient supplied by the inorganic and inorganic sources led to better translocation of photosynthates into form of grain resulting in better yield attributes. Similar findings were reported by Kannan *et al.* (2013).

**Comment [S9]:** better complete with minimum data as well

#### 3.3 Yield

After evaluated the data recorded post harvesting of crop show that significantly higher seed yield (1.59 t/ha), higher Haulm yield (4.16 t/ha) and harvest index (27.62%) was recorded in treatment with the application of Molybdenum  $1\text{ kg ha}^{-1}$  + Boron 0.5% at 30DAS & 45 DAS.

The judicious use of Molybdenum and Boron showed beneficial effect on physiological process of plant metabolism viz. growth and yield there by leading to higher grain yield. similar findings were reported by Ranjit Chatterjee *et al.*, (2015) who observed that Molybdenum and Boron significantly increased grain yield of Cowpea.

### 4. Conclusion

Based on the above experimental findings, it is concluded that application of nutrients in combination of Molybdenum  $1\text{ kg ha}^{-1}$  and Boron 0.5% at 30DAS & 45 DAS accomplished better growth parameters, yield attributes, higher seed yield, higher gross returns and net returns in cowpea under eastern Uttar Pradesh conditions.

**Comment [S10]:** layout customize journal template

**Comment [S11]:** preferably complete with data from the discussion

## 5. References

- Ahmad, W., Zia, M.H., Malhi, S.S., Niaz, A., Saifullah, 2012. Boron deficiency in soils and crops: A Review. Crop Plant, Aakash Goyal (Ed.), Intech Open, Page 78. DOI: 10.5772/36702.
- Ahmad\*, A.N., Kanwal, S., Rahmatullah, Khalid Rasheed, M., 2009. Role of boron in plant growth: a review. Journal of Agricultural Research 2009, 47(3) 329 - 338
- Ahmad\*, A. Niaz\*\*, S. Kanwal\*\*\*, Rahmatullah and M. Khalid Rasheed\*\* ROLE OF BORON IN PLANT GROWTH: A REVIEW. J. Agric. Res., 2009, 47(3)
- Ali, M. and Gupta, S. (2012) Carrying capacity of Indian Agriculture: pulse crops. Current Science 102, 874-881.
- Ahluwat IPS, Shivakumar BG. *Kharij Pulses*. In Textbook of Field Crops Production (R. Prasad, Ed.) ICAR, New Delhi, India 2005.
- Alves, J.M., E.C. Guimaraes, J.S. Alves and J.J. Neto, 2002. Molybdenum application in cowpea (*Vigna unguiculata* (L.) Walp) leaf. Revista Universidade Rural. Serie Ciencias da Vida. 2002. 22: Suplemento. 193-197.
- Anonymous., 2011. Annual report of Indian Council of Medical Research, Hyderabad.
- Badiane, F.A., Diouf, M., Diouf, D., 2014. Cowpea. In: Singh, M., Bisht, I.S., Dutta, M. (Eds.). Broadening the Genetic Base of Grain Legumes. Springer, New Delhi, 95-114.
- Boukar, O., Belko, N., Chamarthi, S., Togola, A., Batiemo, J., Owusu, E., Haruna, M., Diallo, S., Umar, M.L., Olufajo, O., Fatokun, C., 2019. Cowpea (*Vigna unguiculata*): Genetics, genomics, and breeding. Plant Breeding 138(4), 415-424
- Bhagiya, S.R., K.B. Polara and J.V. Polara, 2005. Effect of B and Mo on yield, quality and nutrient absorption by groundnut. Advances in Plant Sciences, 18(2): 803-806.
- Boukar, O., Massawe, F., Muranaka, S., Franco, J., Maziya-Dixon, B., Singh, B., Fatokun, C., 2011. Evaluation of cowpea germplasm lines for protein and mineral concentrations in grains. Plant Genetic Resources 9(4), 515-522.
- Das D.K. (2011). Micronutrients: their behavior in soil and plants. Kalyani publisher, New Delhi. Pp. 165-168
- Debnath\*, S.K. Pattanaik, D. Sah, G. Chandral and A.K. Pandey (2018). Effect of Boron and Zinc Fertilization on Growth and Yield of Cowpea (*Vigna unguiculata* Walp.) in Inceptisols of Arunachal Pradesh. Journal of the Indian Society of Soil Science, Vol. 66, No. 2, pp 229-234 (2018) DOI: 10.5958/0974-0228.2018.00029.4
- Ehlers, J.D., Hall, A.E., 1997. Cowpea (*Vigna unguiculata* L. walp.). Field Crops Research 53(1-3), 187-204.
- Gungula, D.T. and Y. Garjila, 2006. The effects of molybdenum application on growth and yield of cowpea in yola, Nigeria. American-Eurasian Journal of Agriculture and Environmental Sciences, 1(2): 96-101.
- Hristozkova, M., M. Geneva and I. Stancheva, 2006. Response of Pea Plants (*Pisum sativum* L.) to Reduced Supply with Molybdenum and Copper. International Journal of Agriculture and Biology, 8(2): 218-220.
- Jing Li, Y., Z. Xiao Ping, L. Deng Yu, C. Qiang and W. Ke Mei, 2006. Study of the effects of Mo and Bradyrhizobium inoculation in acid purple soil by *celB* gene marker. Plant Nutrition and Fertilizer Science, 12(2): 250-253.
- Kannan, R. L., Dhivya, M., Abinaya, D., Krishna, R. L., & Krishnakumar, S. (2013). Effect of

- integrated nutrient management on soil fertility and productivity in maize. *Bulletin of Environment, Pharmacology and Life Sciences*, 2(8), 61-67.
- Kumar, S., Phogat, D., Bhusal, N., 2015. Characterization of elite forage cowpea genotypes for various DUS traits. *Forage Research* 40(4), 232–236.
- Lioi, L., Morgese, A., Cifarelli, S., Sonnante, G., 2019. Germplasm collection, genetic diversity and on-farm conservation of cowpea [*Vigna unguiculata* (L.) Walp.] landraces from Apulia region (southern Italy). *Genetic Resources and Crop Evolution* 66(1), 165–175.
- Mandal, M.K., Pati, R., Mukhopadhyay, D. and Majumdar, K. (2009) Maximizing yield of cowpea through soil test based nutrient application in Terai alluvial soils. *Better Crops* 3, 28-30.
- Meena, H.K., Krishna, K.R., Singh, B., 2015. Character associations between seed yield and its component traits in cowpea [*Vigna unguiculata* (L.) Walp.]. *Indian Journal of Agriculture Research* 49(6), 567–570.
- Nadia Gad, and M.R. Abd El-Moez (2013) Influence of Molybdenum nodulation, Nitrogen fixation and yield of Cowpea. *Journal of Applied Sciences Research*, 9(3):1498-1504, 2013
- Ram, H., 2014. *Vegetable Breeding Principles and Practices* (3rd Edn.). Kalyani publishers New Delhi, 460–473. Ram, H., Hedau, N.K., Chaudhari, G.V., Choudhary, 2021. Genetic diversity assessment in M Pea (*Pisum sativum* L.) using Microsatellite Markers. *International Journal of Bio-resource and Stress Management* 12(4), 402–408.
- Ranjit Chatterjee, Subhendu Bandyopadhyay, Effect of boron, molybdenum and biofertilizers on growth and yield of cowpea (*Vigna unguiculata* L. Walp.) in acid soil of eastern Himalayan region. *Journal of the Saudi Society of Agricultural Sciences*, Volume 16, Issue 4, 2017, Pages 332-336, ISSN 1658-077X, <https://doi.org/10.1016/j.jssas.2015.11.001>.
- Salwinder Singh,; Sharma, V.; Shukla, A.K.; Kaur, J.; Verma, V.; Kaur, M.; Singh, P.; Brestic, M.; Gaber, A.; Hossain, A. Interactive Effects of Molybdenum, Zinc and Iron on the Grain Yield, Quality, and Nodulation of Cowpea (*Vigna unguiculata* (L.) Walp.) in North-Western India. *Molecules* 2022, 27, 3622. <https://doi.org/10.3390/molecules27113622>
- Shaik Karimunnisa, Umesha C., Shahazad Ahmed Khan. Effect of foliar application of Boron and soil application of Zinc levels on growth and yield of Cowpea (*Vigna unguiculata*). *Pharma Innovation* 2021; 10(8):1341-1344.
- Shivakumar; U.I. and Kumutha, K. (2003). Effect of Rhizobium and Mo on nodulation yield and yield contributing characters of groundnut. *Journal of Ecobiology*, 15 : 451-455.
- Singh, M. V. "Spread of micro-nutrient deficiencies specially boron in India and response of field crops." *Brain storming workshop on soil test based nutrients including boron and other micro nutrients* Organized by ICRISAT-Agriculture Directorate, Karnataka-Rio Tinto India at Bangalore, India. 2012
- Somla Naik R, Lora Anusha P, Siva Kumar B. Review on foliar nutrition enhancing the growth and yield characters in blackgram. *International Journal of Chemical Studies* 2018; 6(6):2212-2216.
- Srivastava, S., Chopra, A.K., Sharma, P., Kumar, V., 2016. Amendment of sugar mill wastewater irrigation on soil bio hydrological properties and yield of cowpea [*Vigna unguiculata* (L.) Walp.] in two seasons. *Communication in Soil Science and Plant Analysis* 48(5), 511–523.

Subasinghe, S., Dayatilake, G.A. and Senaratne, R., 2003. Effect of B, CO and MO on nodulation, growth and yield of cowpea (*Vigna unguiculata*). Tropical Agricultural Research and Extension, 6, pp.108–112.

DOI:<http://doi.org/10.4038/tare.v6i0.5448>

Timko, M.P., Ehlers, J.D., Roberts, P.A., 2007. Cowpea. In: Kole, C. (Ed.). Genome Mapping and Molecular Breeding in Plants, Pulses, Sugar and Tuber Crops (Vol.3). Springer, Berlin, Heidelberg, 49–67.

Venkatesan, M., Prakash, M., Ganesan, J., 2003. Correlation and path analysis in Cowpea [*Vigna unguiculata* (L.) Walp.]. Legume Research 26(2), 105–108

UNDER PEER REVIEW

**Table 1. Growth attributes of Cowpea at harvest as influenced by Molybdenum and Boron.**

Treatments	Plant Height (cm)	Plant Dry Weight (g)	No. of Branches per Plant	CGR (gm <sup>2</sup> day <sup>-1</sup> )	RGR (g <sup>-1</sup> g <sup>-1</sup> day <sup>-1</sup> )
Molybdenum 0.5kgha <sup>-1</sup> + Boron 0.5% at 30DAS.	63.20	34.27	12.07	12.15	0.0195
Molybdenum 0.5k ha <sup>-1</sup> + Boron 0.5% at 45DAS.	63.67	34.92	12.33	12.12	0.0190
Molybdenum 0.5k ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	64.63	36.20	12.87	12.93	0.0196
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	65.57	37.19	13.33	13.03	0.0192
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 45DAS.	66.58	38.46	14.07	13.11	0.0186
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	70.53	41.33	14.40	9.92	0.0123
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	65.45	39.38	13.53	11.87	0.0160
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 45DAS	69.55	40.36	13.47	10.56	0.0136
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	72.55	43.68	15.33	11.22	0.0133
Control	62.41	33.34	11.13	11.78	0.0194
F test	S	S	S	S	S
SEm(±)	0.06	0.03	0.11	0.05	0.00
CD(p=0.05)	0.19	0.09	0.33	0.15	0.00

Treatment	Yield attributes		
	No. of podsplant <sup>-1</sup>	No. of seeds Pod <sup>-1</sup>	Seed Index (g)
Molybdenum 0.5kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	11.93	10.47	16.53
Molybdenum 0.5kg ha <sup>-1</sup> + Boron 0.5% at 45DAS.	12.20	10.73	16.68
Molybdenum 0.5kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	12.80	10.87	17.19
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	12.93	11.13	16.91
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 45DAS.	14.40	10.93	17.19
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	16.53	11.4	18.15
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	14.33	10.47	17.53
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 45DAS	14.73	11.33	17.68
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	16.73	11.67	18.38
Control	10.53	10.27	16.25
F test	S	S	S
SEm(±)	0.08	0.1	0.03
CD(p=0.05)	0.23	0.3	0.10

**Table 2. Yield attributes of Cowpea at harvest as influenced by Molybdenum and Boron.**

Treatment	Seed Yield (tha <sup>-1</sup> )	Haulm Yield (tha <sup>-1</sup> )	Harvest Index (%)
Molybdenum 0.5kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	1.18	3.58	24.72
Molybdenum 0.5kg ha <sup>-1</sup> + Boron 0.5% at 45DAS.	1.19	3.68	24.50
Molybdenum 0.5kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	1.23	3.96	23.73
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	1.26	3.72	25.36
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 45DAS.	1.34	3.85	25.80
Molybdenum 0.75kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	1.55	4.14	27.18
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 30DAS.	1.35	3.88	25.79
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 45DAS	1.37	3.96	25.69
Molybdenum 1kg ha <sup>-1</sup> + Boron 0.5% at 30DAS & 45 DAS.	1.59	4.16	27.62
Control	1.07	3.33	24.36
F test	S	S	S
SEm(±)	0.01	0.01	0.21
CD(p=0.05)	0.04	0.02	0.62

**Table 3. Yield of Cowpea at harvest as influenced by Molybdenum and Boron.**