

Minireview Article

Phytochemical Screening ~~for~~of Medicinal Plants: Guide for Extraction Methods

Abstract

Sri Lankans have relied on traditional medicine to meet their primary health needs since the beginning of time. This island is blessed with an enormous number of medicinal plants, which play a very important role in traditional medicine. However, the lengthy documented history of traditional medicine is still composed of medicinal plants, which are not scientifically proven to have the mentioned abilities or activities. In order to obtain scientifically sound information from this documented history of traditional medicine, extraction of the biologically active compounds from these medicinal plants is very important. Also, to maintain the accuracy of results obtained from *in-vitro* and *in-vivo* assays, it is important to consider the pre-extraction procedures as well as the evaporation and storage conditions of the extract. There are several extraction methods accessible in Sri Lanka. This research aims to review the pre-extraction preparation, extraction methods, evaporation techniques, and storage conditions of the plant extract.

Keywords: Sri Lanka, pre-extraction, medicinal plants, traditional medicine, menstruum, sublimation

Introduction

Traditional medicine has a long history in Sri Lanka and medicinal plants play a major role in it. A variety of illnesses and conditions are treated with medicinal plants in Sri Lanka (1,2, 3, 4, 5, 6, 7). Sri Lanka is an island surrounded by the Indian Ocean. In ancient times this island nation was known to be ruled by King Ravana, who was the author of many books of traditional medicine like Arkaprakasya, Kumarathanthraya, UdishaThanthraya, and Nadivignanaya. During his reign, King Ravana represented Sri Lanka at a medical symposium at the base of the Himalayas in India, according to the Ramayana (8). Sri Lanka is blessed with an enormous number of medicinal plants, with 1430 species representing 181 families and 838 genera. There

are 174 endemic species discovered so far in Sri Lanka (8, 9). In 1988, British ecologist Norman Myers identified Sri Lanka as a biodiversity hotspot based on the specie's endemism and degree of threat (10). Folklore medicine continues to meet the foremost health-related needs of 60% to 70% of Sri Lankans in rural regions. Sri Lankans have a well-documented and lengthy history of traditional medicine practice (1, 9). Medicinal plants are found to be prominent in phytoconstituents and secondary metabolites which play a major role in exhibiting many biological potencies such as anti-bacterial, anti-cancer, anti-diabetic, anti-fungal, anti-inflammatory, antioxidant, and antiparasitic activities. Secondary metabolites like alkaloids, essential oils, phenols, quinones, resins, saponins, steroids, terpenes, and tannins have shown various biological activities both *invitro* and *invivo* over the years. These secondary metabolites can be present in any plant part such as bark, flowers, fruits, leaves, rhizomes, roots, seeds, stem, and tubers (11,12). Medicinal plants are one of the plentiful bioresources of medicinal drugs. Most current drugs involved in routine medications were originally extracted from plants, and now they are produced synthetically (13). According to pharmaceutical sciences, Extraction is the separation of medicinally vital portions of selected plant or animal organs with the use of selective solvents and the most appropriate standard extraction procedure. Extracts of herbal plants can be utilized directly to treat specific illnesses and disorders, or they may need to be further processed before attaining therapeutic potential. There are many techniques used for the extraction purpose. Aqueous-alcoholic extraction by fermentation, counter-current extraction, decoction, digestion, hot continuous extraction (Soxhlet), infusion, maceration, microwave-assisted extraction, percolation, phytonic extraction (with hydrofluorocarbon solvents), supercritical fluid extraction, and ultrasound extraction (sonication) are some of them (14). There can be many factors that contribute to the medicinal value of a plant. During the extraction of herbal plants, the active phytoconstituents of the plant or secondary metabolites like alkaloids, flavonoids, glycosides, saponins, steroids, and terpenes can be separated from the inert components of the plant (15).

The nature of the plant substance, the type of solvent being used, the pH of the solvent being utilized, the temperature of the system, and the solvent-to-sample ratio all influence the selection of the most suitable extraction method. It is also determined by how the final products will be utilized (16, 17). Accurate identification and authentication, appropriate and well-timed collection of plant material, and pre-extraction procedures also have a greater influence on the

end product (18). Numerous studies have proved that the storage condition of plant extracts has a major impact on their bioactivity. Therefore, considering the storage conditions are also very important after the extraction of medicinal plants (19, 20). Usually, extraction of medicinal plants includes the following steps: collection of the plant material, identification and authentication of the selected plant, size depletion or homogenization, extraction, filtration, concentration, drying, and reconstitution. This study aims at reviewing different types of extraction methods for plants.

Identification and authentication of the plant

Plant authentication is the process of confirming the authenticity of plant material. After the selection of the plant for extraction, it should be identified accurately according to organoleptic properties such as color, odor, taste, and morphological characteristics such as appearance, shape, and size of the plant, before collection of the plant sample (21). Then the plant material will be collected, prepared, packaged, and stored for the authentication process. In the Sri Lankan setting, according to the national herbarium guidelines the plant material should be dried and pressed for a week. Then the identification and authentication of the plant material are performed by a scientific officer with the use of the voucher specimen. In Sri Lanka, the authentication of plants can be done by the National Herbarium in Royal Botanical Garden, Peradeniya, and Bandaranaike Memorial Ayurveda Research Centre in Nawinna, Maharagama (22). Some literature reveals that the authentication process is also done by a senior professor or Botanist/ Taxonomist (23, 24, 25).

Pre-treatment processes of plant materials

Several processes are involved in the pre-extraction preparation of the plant material, and it is very important to preserve bioactive components inside the plant extract before extraction. The initial step is the pre-washing of the sample collected, and it is very important when using organs of plants like roots, rhizomes, tubers, etc. (26). The extraction process can be done using both fresh and dried samples. However, in many instances dried samples are preferred because it is difficult to maintain the freshness of the sample and fresh samples are delicate and are prone to

deteriorate more quickly. It is also very cumbersome to design the experimental setting within a few hours of the sample collection (18, 27).

Fresh vs dried samples used for extraction

1. A study conducted in Nigeria **revealsrevealed** that extracts produced by raw and dried leaves of *Carica papaya* show different antimicrobial potentials. Dried leaf extract shows significant antimicrobial activity against both Gram-positive and Gram-negative bacteria while fresh leaf extract shows significant antimicrobial activity against only Gram-negative bacteria (28).
2. A study was conducted in Poland to **revealrevealed** the total phenolic count, scavenging activity, and ferric-reducing antioxidant power (FRAP) of dried and fresh extracts of aerial parts of *Coriandrum sativum* L., *Levisticum officinale* Koch., and leaves of *Plectranthusamboinicus*. Results revealed that dried extracts of the herbs exhibited higher total phenolic content and greater antioxidant activity compared with the fresh extracts (29).
3. A study was conducted in South Africa to reveal the phytochemical content and antioxidant properties of dried and fresh extracts of rhizomes of *Tulbaghiaviolacea* Harv. Results revealed that fresh methanolic extract contains higher flavonoids, flavanols, phenolics, tannin, and proanthocyanidin concentrations and it also exhibited higher antioxidant activity compared with the dried extract (30).

After prewashing, excess water should be blotted off with a clean cloth otherwise fungi and other microorganisms can grow in the plant material during drying. Fresh samples are most often allowed to dry to obtain a constant weight. There are several methods used in drying plant materials for the preparation of extracts namely, air-drying, freeze-drying, microwave-drying, and oven-drying.

Air-drying

Time **needsneeded for air dryingto air dry** can vary from several days to months or even **can go** beyond a year depending on the plant species and the plant material being dried (e.g., bark, leaves, seeds, and tubers). In this method, plant materials are allowed to air-dry in a dry, shaded,

and well-ventilated place at ambient temperature. Air drying does not utilize high temperatures, thereby enabling the preservation of heat-labile components inside the plant. This technique is widely used in the extraction of medicinal plants in Sri Lanka, but it consumes a lot of time and is susceptible to contaminations in fluctuating temperature conditions, compared with other drying methods like freeze-drying, microwave-drying, and oven-drying (18, 31).

Microwave-drying

Microwave-drying method is a quick-drying method, which utilizes electromagnetic radiation. The above method can reduce the time required for drying but at the same time promote the degradation of phytoconstituents (18).

Oven-drying

This is also another drying technique that employs heat to remove water from plant materials. It is considered one of the simplest and quickest drying processes to preserve phytoconstituents (18).

Freeze-drying

Another form of product drying is freeze-drying, and this method utilizes the principle of sublimation. This process is also known as lyophilization or molecular drying. Sublimation can be defined as the direct transfer of a substance from the solid phase to the gas phase without passing through the transitional liquid phase. In the ice sublimation process, product drying is accomplished by the removal of water from the frozen material (18, 31, 32).

Before extraction, and after the drying process, plant material is subjected to shredding and grinding. The particle size influences the degree of extraction; when the particle size is small, the surface area becomes high resulting in a greater area of exposure among the plant sample and respective solvents. This results in a higher extraction yield. Shredding produces coarse particles, whereas powdered specimens have more homogeneity levels and finer particles. This results in greater interaction of solvents with the plant material. Past research has demonstrated that when the particle size is less than 0.5 mm, it can give a higher yield of extraction of active components

(33). For the homogenization process of plant material, traditional mortar and pestle, and laboratory blenders are frequently used in routine laboratory setups (18).

Solvents for Extraction

It is important to select the most appropriate solvent (also known as the menstruum) during an extraction process. It is determined by the plant species, which plant part is extracted, the makeup of biologically active compounds, and the accessibility of the menstruum. To extract polar components, a polar solvent such as ethanol, methanol, or water can be used. Non-polar components should be extracted using non-polar menstruum (e.g., dichloromethane and hexane) (15, 17, 34). Usually, two solvents that are capable of being mixed are used during liquid-liquid extraction (e.g.: water-dichloromethane, water-ether, and water-hexane). Water is involved in all combinations because of its high polarity and the possibility of being mixed with organic solvents. It is important that the compound which needs to be extracted should dissolve in the organic solvent but not in the water for a successful separation (35). Furthermore, menstruum is classified according to the polarity, with n-hexane having the lowest polarity while water has the highest polarity of them all (15, 17, 34).

Below are some different extraction solvents listed in increasing polarity order (15, 36).

Table 1: Polarities of different extraction solvents.

Solvents	Polarity
n-Hexane	0.009
Petroleum ether	0.117
Diethyl ether	0.117
Ethyl acetate	0.228
Chloroform	0.259
Dichloromethane	0.309
Acetone	0.355
n-Butanol	0.586
Ethanol	0.654

Methanol	0.762
Water	1.000

Properties of Solvents used in the extraction

- i. Alcohol: This is a polar substance that can be mixed with water and is capable of extracting polar constituents from plant materials (36, 37). Self-preservation of the plant extract in concentrations above 20%, non-toxic properties at low concentrations, and only a minimal amount of heat required to concentrate can be listed as the advantages of using alcohol as a solvent. There are also some disadvantages such as being flammable, evaporative, and failing to dissolve waxy substances, gums, or fats. (36, 37).
- ii. Chloroform: This solvent is nonpolar and used in the extraction of compounds like flavonoids, fats, oils, and terpenoids. Being colorless, odorless, soluble in alcohols, and easily absorbed and metabolized by the body are the advantages of chloroform. Disadvantages are, it is both sedative and carcinogenic (17, 37, 38).
- iii. Dichloromethane: This is a solvent that is widely involved in the extraction of bioactive compounds from medicinal plants because it can extract both nonpolar as well as polar compounds (39). Dichloromethane is widely used in the extraction of caffeine as it is found to be more soluble in dichloromethane (140 mg/ml) than in water (22 mg/ml) (40). This solvent has many advantages. Some of them are, it is not miscible in water and therefore it has the capacity to dissolve a scope of organic compounds. When these properties are combined with its volatility, it makes dichloromethane a very effective solvent in many industries. Along with these advantages, there are many disadvantages too. Very high volatility and stability, ability to cause CO poisoning, neurotoxicity, and carcinogenicity are some of them (41).
- iv. Ether: Can be identified as a solvent that is nonpolar and capable of being utilized to extract components such as fatty acids, terpenoids, alkaloids, and coumarins. Ether has some benefits, including the ability to mix with water, having a low boiling point, and the absence of taste. It is also a substance that is extremely stable and is not influenced by metals, acids, or bases. Its disadvantages include its high flammability and volatility (17, 37, 38).

- v. Hexane: Considering its ease of recovery, a low boiling point (63–69 °C), and exceptional dissolving properties, hexane has been frequently utilized to extract oil from oil seeds (17). Hexane gets released into the environment over the procedure of extraction and recovery steps, where it can interact with pollutants to produce ozone and photochemicals (42). Hexane is found to be soluble in neural lipids which can have a significant impact on the nervous system when inhaled by individuals (15). Thus, to replace n-hexane without compromising oil yield, researchers are looking for alternatives that are safer for the environment, human health, and safety. Green solvents have thus offered an effective replacement for the extraction of oil.
- vi. Green solvent (ionic liquid): This is a one-of-a-kind menstruum that has a high polarity and is steady against high temperatures (e.g.: 3,000°C). It has the ability to mix with water and other polar solvents, making it ideal for polar compound extraction. The benefits of green solvents can be listed as follows, it is ideal for microwave-assisted extraction, non-flammable, have high polarity, and are useful for liquid-liquid extraction. The disadvantages are, not being ideal to create tinctures (43).
- vii. Water: It can be nominated as the solvent with the highest polarity among other solvents and is utilized to extract diverse polar compounds. The advantages of using water as a solvent are, it dissolves many different substances, is inexpensive, has nonflammability, is nontoxic, and has high polarity. Disadvantages are, it encourages bacterial and fungal growth, could result in hydrolysis, and needs a lot of heat to concentrate the extract (36, 37).

Factors that need to be considered in the selection of solvents for extraction

Numerous factors have to be considered while selecting a solvent for extraction.

- i. Boiling temperature: The boiling temperature of the solvent should be minimum to prevent degradation by heat.
- ii. The cost should be as affordable as possible.
- iii. Safety: Non-toxicity and non-flammability are the preferred characteristics of a perfect extraction solvent.

- iv. **Selectivity:** The solvent used for extraction should have a high selectivity, which is the capability to separate biologically active compounds from inert constituents in plant material.
- v. **Reactivity:** The reactivity of the solvent has to be very low so it doesn't react with the extract.
- vi. **Recovery:** It should be possible to separate the solvent from the extract quickly.
- vii. **Viscosity:** It's ideal for a solvent to have a low viscosity to facilitate easy penetration (17, 36, 44).

Commonly used extraction methods

1. Maceration




Figure 1: Extraction by maceration method, adapted from (63).

In this routinely used extraction method, an airtight vessel (preferably amber in color) is used. Then the vessel is packed with plant material such as barks, flowers, leaves, roots, seeds, and tubers which are coarsely powdered. Menstruum, which is the solvent used in extraction, is subsequently poured over the plant material until being fully covered. Then the vessel is tightly sealed and stored for 3-7 days with continuous agitation regularly. The days kept for maceration depend on the type of plant material being used and the method of evaporation chosen. After the extraction process, the mixture is filtered using filter paper or a muslin cloth before evaporation. Then, the micelle can be evaporated with the help of an oven or a water bath to isolate it from the extraction solvent. This method is considered very suitable for heat-stable plant materials. Based on the requirement, the sample-to-solvent ratio can vary from 1:4, 1:5, and 1:10 (16, 17, 18, 35, 42, 45).

2. Infusion

This process is more similar to maceration. Here, finely powdered plant material is used for the extraction. Hot or cold menstruum is then added over the finely powdered plant material and the mixture is stored for a short time period. Infusion is considered more appropriate for the extraction of readily soluble biologically active compounds. Also, it is found to be a very suitable technique to obtain fresh extracts prior to use. Here, the sample-to-solvent ratio is generally 1:4 or 1:16 which depends on the requirement (16, 17, 18, 35).

3. Digestion

Digestion is a type of maceration where light heat is used for the extraction procedure. This technique is utilized when increased temperatures are not an issue for the plant constituents. The extraction of the solvent is more effective because of the higher temperatures (46).

4. Decoction

This is a continuous hot extraction technique where the powdered crude plant material is boiled with a specific amount of water for a period of specific time (about 15 minutes). Heat is applied throughout the procedure to speed up the extraction. This method is ideal when extracting water-soluble and thermally stable compounds from plant materials. In this process, the most commonly used sample-to-solvent ratios are 1:4 or 1:16 (16, 17, 18, 35).

5. Percolation




Figure 2: A diagram of the percolation extraction, adapted from (64).

A percolator is a narrow cone-shaped container made with glass which has openings on both sides. The finely powdered plant material will be added to a clean vessel followed by adding a higher quantity of menstruum to soak the powder. This mixture is then stored for a certain time period (4 hours). This mixture is then introduced to the percolator (during this procedure, the lower end has to be closed). Now, the system will be kept to stand for 24 hours. After 24 hours, the solvent used for extraction is added from the upper end to flow down until the plant material is saturated with the solvent. The stop cork is opened at the lower end while adding menstruum from the top. The liquid is then collected from the bottom. This procedure is carried out with the help of gravitational force which helps to push down the solvent through the plant material. Pouring of the menstruum should end when the amount of menstruum reaches around 75% of the total preparation. Finally, the collected micelle will be filtered and concentrated (17, 18, 35).




Figure 3: The process of percolation, adapted from (64).

6. Soxhlet extraction




Figure 4: A diagram of the Soxhlet extraction instrument, adapted from (65).

This is an extraction method that uses continuous heat. In this procedure, a glass instrument known as the Soxhlet extractor is utilized. This instrument comes with a solvent flask, a siphon tube, a condenser tube, and many other parts. The plant material which needs to be extracted is introduced to a part known as the thimble, which is a porous bag. First, the round bottom flask should be filled with the menstruum. Then heat is applied from the bottom of the flask. This results in to production of solvent vapor which reaches the condenser through the distillation tube. inside the condenser, the vapor gets condensed and falls to the thimble. The plant material which is already inside the thimble will get contact with the fallen solvent. Components of the plant material that is soluble in the solvent will get mixed. Then, the solvent mixture starts to fill both the thimble and siphon tube. When the solvent mixture fills up to the maximum height of the siphon tube it starts to overflow and fall back to the flask.

The advantages of this method are,

It can be involvedutilized forinthe extraction ofextracting large numbers of drugs with low volumes of solvents. This is valid for plant material that can withstand high heat. It does not require filtration.

The disadvantages of this method are,

Requires a large amount of heat.

Impossibility of regular shaking and not being suitable for thermolabile materials (16, 17,18, 35, 42, 47, 48).

7. Microwave-assisted extraction




Figure 5: A diagram of Microwave-assisted extraction, adapted from (66).

Microwave- assisted extraction is an advanced extraction technique used in the extraction of herbal medicines. It involves the mechanism of dipole rotation and ionic transfer to displace charged ions from the solvent and drug materials. This technique employs electromagnetic radiation having frequencies ranging from 300 MHz to 300 GHz and wavelengths ranging from 1 cm to 1 m. This approach uses microwave radiation to bombard a substance that absorbs

electromagnetic energy and converts it into thermal energy. Due to the heat generated, it facilitates the solvent to penetrate the plant material.

This method requires a minimum quantity of solvents and time for extraction. It also can give a higher output. However, being suitable only for flavonoids and phenolic compounds and the possibility of degrading tannins and anthocyanins due to the high heat can be some downsides of this method (16, 34, 42, 43).

8. Sonication (Ultrasound assisted extraction).




Figure 6: A diagram of extraction by sonication, adapted from (67).

In this extraction technique, ultrasound frequencies are used that are ranging from 20 KHz to 2000 KHz. They have the capability to disturb the plant cell walls and thereby increase the surface area of plant material to facilitate the penetration of solvents. Subsequently, this facilitates the release of biologically active compounds.

The advantages of this method are,

This method can be used for involved in small sample amounts. It gives us a high yield while reducing extraction time and the quantity of menstruum.

Even though this method is useful in some incidents such as the extraction of rauwolfia roots, involving this process on a large scale can be a disadvantage because of the high cost. Also, the high ultrasound energy can sometimes degrade the phytochemicals due to the production of free radicals (17, 18, 34, 38).

9. Aqueous- alcoholic extraction by fermentation.




Figure7: A diagram of extraction by aqueous alcoholic fermentation, adapted from (68).

To extract the active ingredients, some Ayurveda medicinal preparations (such as āsava and arista) use fermentation as a process. For the extraction of active components from the plant, the crude drug must be soaked for a designated amount of time, either as a powder or a decoction (kasaya). During this time the drug mixture will undergo fermentation and produce alcohol. This action will enable the active compounds present in our plant to be extracted. However, this extraction technique is not yet standardized in Ayurveda, but when considering the extraordinary level of improvement in fermentation technology, it can be used to produce herbal drug extracts (18).

10. Counter-current extraction




Figure 8: A diagram of counter-current extraction, adapted from (69).

In this method, to create a fine slurry, the wet material is ground using a toothed disc disintegrator. The material to be extracted is allowed to move in a single direction inside a cylindrical extractor. The plant material will contact the menstruum. The more the plant material moves, the extracts become more concentrated. When the solvents, material quantities, and their rate of flowing are optimized, full extraction is attainable. This procedure is considered very quick and efficient, raising fewer adverse effects from high temperatures. Then, the extract which is sufficiently concentrated exits the extractor at one end while the marc, which is now free of solvents, exits at the other end (14).

This method comes with many advantages.

When compared with methods like maceration, decoction, and percolation, this method requires less volume of solvent. As the extraction happens at room temperature, the thermolabile compounds will be preserved by heat. Again, as the grinding happens in wet conditions, the

thermal energy generated during grinding will be transferred to water thereby, protecting the thermolabile compounds (14).

11. Superficial fluid extraction




Figure 9: Schematic diagram of extraction by Superficial fluid, adapted from (70).

Superficial fluid extraction (SFE) is an advanced extraction technique that utilizes supercritical fluid as the menstruum. Even though it can also be done with liquids, extraction is typically done from a solid matrix. SFE is utilized to remove unwanted compounds or collect wanted compounds from a sample mixture. Carbon dioxide (CO_2) is the most commonly used supercritical fluid but sometimes, alcohols like methanol and ethanol can also be used. This procedure of extraction, in contrast to other methods, leaves no solvent residue behind (49).

This method comes with great advantages.

SFE can be named as a replacement for liquid extraction where it involves hexane or dichloromethane as the solvent. The extract and matrix will always contain a small quantity of residual solvent; therefore, their use will always have some negative effects on the environment. Contrarily, CO_2 is simple to remove by simply lowering the pressure, virtually undetectable, and environmentally safe. Therefore, this method helps in the improvement of the environment with a reduced number of contaminations.

Also, this method gives pure products with high speed and recovery (49).

12. Reflux extraction method




Figure 10: A schematic diagram of the heat reflux extraction method, adapted from (71).

The reflux method is a popular extraction method in Sri Lanka, and studies show that it can increase extraction yield whilst lowering the amount of solvent utilized. Reflux extracting refers to a solid-liquid extraction method at a uniform temperature with replicated solvent evaporation as well as condensation for a set period of time without loss of solvent. The technique is commonly utilized in the herbal industry since it is effective, simple to operate, and affordable (50, 51).

Evaporation of plant extracts

After extracting a plant material, it has to be evaporated to obtain a dry extract. This can help to maintain the viability of the constituents in the extract.

Several evaporation methods are used such as,

- i. Evaporation by air at room temperature underneath a hood for twenty-four hours in the light.
- ii. Evaporation using a rotary vacuum evaporator under reduced pressure. (The temperature and time can change according to the type of solution).
- iii. Evaporating inside an oven in the dark (The temperature and time can change according to the type of solution) (52, 53).

The rotary vacuum evaporator




Figure 11: A labeled diagram of the rotary vacuum evaporator, adapted from (72).

This instrument was invented by Lyman C. Craig. It is a device used in chemical laboratories to effectively evaporate solvents from samples under low pressure. The rotary evaporator's function is to heat the sample and lower the boiling point of molecular compounds while operating in a

vacuum. To increase the rate of evaporation by expanding the evaporation area, the round-bottomed flask where evaporation occurs is kept rotating while in operation (54).

Depending on the solvent being used, a rotary evaporator's optimal distillation parameters change.

The water bath temperature should be set to 60 °C (It should not exceed this temperature). The temperature of the cooling water should be below 20 °C. The required vacuum for a solvent boiling point of 40 °C, 30 °C, and 50 °C can be adjusted using the table below (55).

Table 2: The list of vacuum pressures common solvents in different temperatures (55).

Solvent	Vacuum [mbar] for bp of 30°C	Vacuum [mbar] for bp of 40°C	Vacuum [mbar] for bp of 50°C
Acetic acid	26	44	72
Acetone	370	556	815
Acetonitrile	153	226	315
n-Amyl alcohol, n-pentanol	6	11	20
n-Butanol	14	25	44
tert-Butanol	78	130	231
Chlorobenzene	22	36	56
Chloroform	332	474	665

Cyclohexane	154	235	347
Dichloromethane	699	Atm. press.	Atm. press.
Diethyl Ether	838	Atm. press.	Atm. press.
Di isopropyl Ether	251	375	545
Dioxane	68	107	165
Dimethylformamide (DMF)	6	11	17
Ethanol	97	175	276
Ethyl Acetate	153	240	366
Heptane	77	120	183
Hexane	264	335	525
Isopropyl alcohol	78	137	231
Isoamyl alcohol	9	14	29
Methanol	218	337	607

Pentane	834	Atm. press.	Atm. press.
n-Propanol	37	67	115
Pentachloro ethane	8	13	21
Tetrachloromethane	179	271	398
Tetrahydrofuran (THF)	234	357	539
Toluene	48	77	118
Trichloroethylene	119	183	275
Water	42	72	120
Xylene	15	25	40

The Storage Conditions of Plant Extracts

There are many different ideas on how to store plant extracts. Ideal storage of plant extracts is very important to preserve the enzyme activities, biological properties, phytochemical constituents, and various other plant characteristics. Proper storage will prevent microbial growth and preserve the plant extracts. Several studies have investigated the importance of ideal storage conditions having a great effect on a plant's biological activities (56, 57).

Any kind of extract should be dried using a rotary evaporator or cephalization (if aqueous extraction is used). After that, you can keep the extracts in a refrigerator for more than two years by sealing the container tightly (58, 59).

“The type of active compounds you're looking for will have a big impact on how long you should store your medicinal plant extract. Some substances, like alkaloids, are stable, but others are not. I believe that inside plants, particularly those that have been freshly lyophilized or freeze-dried using liquid nitrogen, is the best environment for storing natural compounds. Additionally, I advise keeping the dried plant sample in a cool, dark environment. Compared to storing the extract, it is preferable” says Majid Azizi, The Academy of Sciences of the Islamic Republic of Iran (60).

“Extract of the plant can be kept at 4 °C for as long as it is in dry condition (i.e., the sample has been completely dried using a rotavapor or freeze dryer after extraction). To prevent degradation, it is best to keep your extract or solution at -20 °C or lower. When performing a bioassay, make a fresh extract solution” says Narayan D Chaurasiya, Southern Research Institute (61).

“It does rely heavily on the active ingredients, but the solvent is also important. The best course of action is to get your analysis as soon as you can, especially if you're thinking about getting antioxidant tests. The activity is not always maintained when stored at 4 °C. It would be best to store it at -20 °C, at the very least, if there are essential oil compounds present. For instance, matricin concentration can remain constant at -20 °C for more than 6 months, but at 4–8 °C, it can drop to half or even less of its initial value. To have a good activity, you must take the chemical compound's stability into account” says OanaCioanca(**provide the year Oana published tthis statement in a bracket**)Universitatea de MedicinasiFarmacieGrigore T. Popa Iasi, Faculty of Pharmacy (62).

References

1. Peiris, D.S.H.S., Fernando, D.T.K., Senadeera, S.P.N.N., Chandana, A.K. and Ranaweera, C.B. (2022). *Mirabilis jalapa* Linn.: A Folklore Ayurvedic Medicinal Plant in Sri Lanka. *Asian Plant Research Journal*, pp.21–41. doi:10.9734/aprj/2022/v10i2187.
2. Ranaweera, C. B., & Chandana, A. K. (2021). *Clitoriaternatea* - shifting paradigms: From laboratory to industry. *South Asian Journal of Research in Microbiology*, 18–26. <https://doi.org/10.9734/sajrm/2021/v11i230247>
3. Silva, A.R.N., Ranaweera, C.B., Karunathilaka, R.D.N., Pathirana, R. and Ratnasooriya, W.D. (2016). Antibacterial Activity of Water Extracts of Different Parts of *Morinda Citrifolia* Grown in Sri Lanka. *International Journal of Scientific and Research Publication*, 6(5), pp.124–127.
4. Ratnasooriya, R., Ranaweera, W.D., Abeysekara, C.B. and Pathirana, W.P.K.M. (2015). *In vitro* Antioxidant Activity of Methanolic Extracts of Leaves of *Indigofera indica* and Stem Bark of *Stereospermumsuaveolens* Grown in Sri Lanka. *International Journal of Institutional Pharmacy and Life Sciences*, 5(2), pp.128–138.
5. Ranaweera, C., Abeysekara, W., Pathirana, R. and Ratnasooriya, W. (2015) 'Lack of *in vitro* antihyaluronidase activity of methanolic leaf extract of *Indigofera tinctoria* L and methanolic stem bark extract of *Stereospermumsuaveolens* DC', *Journal of Pharmaceutical Negative Results*, 6(1), 40, available: <https://link.gale.com/apps/doc/A415909779/HRCA?u=anon~6214fe20&sid=googleScholar&xid=fd87970f> [accessed 25 May 2023].
6. Samaraweera, T.U., Samaraweera, T.U., Senadeera, S.P.N.N. and Ranaweera, C.B. (2022). Rich Diversity & Potential Medicinal Value of Endemic Sri Lankan Plant: *Jeffreycazeylanica*. *Asian Plant Research Journal*, 10(4), pp.21–34. doi:<https://doi.org/10.9734/aprj/2022/v10i4197>.
7. Ranaweera, C.B., Vidanagamage, A.S., Abeysekara, W.P.K., Silva, A.R.N., Chandana, A.K., Premakumara, S., Pathirana, R. and Ratnasooriya, W.D. (2015). *In vitro* Effects of Aqueous Extracts of Five Sri Lankan Medicinal Plants on Human Erythrocyte Membrane Stabilization

- Activity. *International Journal of Recent Advances in Multidisciplinary Research*, 2(6), pp.486–489.
8. Gunaratne, M.D.N. and Gunarathna, P.L.J. (2015). *Medicinal Plants of Sri Lanka*. M. D. NandanaGunaratne.
 9. Mahendranathan, C. and Priyashantha, A.K.H. (2020). Traditional Uses of Medicinal Plants in Sri Lanka. In: International Conference on Bio Engineering and Life Sciences.
 10. Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. and Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403(6772), pp.853–858. doi:10.1038/35002501.
 11. Ynalvez, R.A. and Compean, K.L. (2014). Antimicrobial Activity of Plant Secondary Metabolites: A Review. *Research Journal of Medicinal Plant*, 8(5), pp.204–213. doi:https://doi.org/10.3923/rjmp.2014.204.213.
 12. CoşgeŞenkal, B. (2020). The Role of Secondary Metabolites Obtained from Medicinal and Aromatic Plants in Our Lives. *ISPEC Journal of Agricultural Sciences*, 4(4), pp.1071–1079. doi:https://doi.org/10.46291/ispecjasvol4iss4pp1069-1077.
 13. Kirtikar KR, Basu BD. Indian medicinal plants. 2nd ed Uttaranchal: Oriental Enterprises; 2001.
 14. Handa, S.S., Khanuja, S.P.S., Longo, G. and Rakesh, D.D. (2008). *Extraction Technologies for Medicinal and Aromatic Plants*. Trieste (Italy): Earth, Environmental and Marine Sciences and Technologies.
 15. S Sasidharan, Y Chen, D Saravanan, K M Sundram and L Yoga Latha (2010). Extraction, Isolation and Characterization of Bioactive Compounds from Plants' extracts. *African Journal of Traditional, Complementary and Alternative Medicine*, 1(8), pp.1–10.
 16. Ingle K P, Deshmukh A G, Padole D A, Dudhare M S, Moharil M P and Khelurkar V C (2017). Phytochemicals: Extraction methods, identification, and detection of bioactive compounds from plant extracts. *Journal of Pharmacognosy and Phytochemistry*. 6, pp.32–36.
 17. Pandey, A. and Tripathi, S. (2014). Concept of standardization, Extraction and Pre-Phytochemical Screening Strategies for Herbal Drug. *Journal of Pharmacognosy and Phytochemistry*, 2(5), pp.115–119.

18. Azwanida, N. (2015). A Review on the Extraction Methods Use in Medicinal Plants, Principle, Strength and Limitation. *Medicinal & Aromatic Plants*, 04(03). doi:<https://doi.org/10.4172/2167-0412.1000196>.
19. Hartmann, J. and Asch, F. (2019). Extraction, Storage Duration, and Storage Temperature Affect the Activity of Ascorbate Peroxidase, Glutathione Reductase, and Superoxide Dismutase in Rice Tissue. *Biology*, 8(4), p.70. doi:<https://doi.org/10.3390/biology8040070>.
20. Del-Toro-Sánchez, C.L., Gutiérrez-Lomelí, M., Lugo-Cervantes, E., Zurita, F., Robles-García, M.A., Ruiz-Cruz, S., Aguilar, J.A., Morales-Del Rio, J.A. and Guerrero-Medina, P.J. (2015). Storage Effect on Phenols and on the Antioxidant Activity of Extracts from *Anemopsis californica* and Inhibition of Elastase Enzyme. *Journal of Chemistry*, 2015, pp.1–8. doi:<https://doi.org/10.1155/2015/602136>.
21. Sharma, D.R. (2022). *What Is Plant authentication? Plant Authentication in Research ? - ACME Research Solutions*. [online] Available at: <https://acmeresearchlabs.in/2022/07/01/plant-authentication/>.
22. . Senadeera, S.P.N.N., Fernando, K.S.K., Wickramasekara, W.L.L.N., Fernando, M.Y.S., Ranaweera, C.B., Rajapaksha, W. and Silva, A.R.N. (2021). *In vitro* Anti-inflammatory Activity of Endemic *Artocarpus nobilis* Thw Found in Sri Lanka. *Asian Plant Research Journal*, pp.116–122. doi:<https://doi.org/10.9734/aprj/2021/v8i430192>.
23. Perianayagam, J.B., Sharma, S.K. and Pillai, K.K. (2006). Anti-inflammatory Activity of *Trichodesma Indicum* Root Extract in Experimental Animals. *Journal of Ethnopharmacology*, 104, pp.410–414. doi:<https://doi.org/10.1016/j.jep.2005.08.077>.
24. Lalitha, K.G. and Sethuraman, M.G. (2010). Anti-inflammatory Activity of Roots of *Ecbolium Viride* (Forsk) Merrill. *Journal of Ethnopharmacology*, 128, pp.248–250. doi:<https://doi.org/10.1016/j.jep.2009.12.030>.
25. S. Vijaya Kumar, P. Sankar & R. Varatharajan (2009) Anti-inflammatory activity of roots of *Achyranthes aspera*, *Pharmaceutical Biology*, 47:10, 973-975, DOI: 10.1080/13880200902967979.
26. Krakowska-Sieprawska, A., Kielbasa, A., Rafińska, K., Ligor, M. and Buszewski, B. (2022). Modern Methods of Pre-Treatment of Plant Material for the Extraction of Bioactive Compounds. *Molecules*, 27(3), p.730. doi:<https://doi.org/10.3390/molecules27030730>.

27. Sulaiman, S.F., Sajak, A.A.B., Ooi, K.L., Supriatno and Seow, E.M. (2011). Effect of solvents in extracting polyphenols and antioxidants of selected raw vegetables. *Journal of Food Composition and Analysis*, 24(4-5), pp.506–515.
doi:<https://doi.org/10.1016/j.jfca.2011.01.020>.
28. Alabi, O.A., Haruna, M.T., Anokwuru, C.P., Jegede, T., Abia, H., Okegbe, V.U. and Esan, B.E. (2012). Comparative Studies on Antimicrobial Properties of Extracts of Fresh and Dried Leaves of *Carica papaya* (L) on Clinical Bacterial and Fungal Isolates. *Advances in Applied Science Research*, [online] 3(5), pp.3107–3114. Available at:
www.pelagiaresearchlibrary.com.
29. Kozłowska, M., Ścibisz, I., Przybył, J.L., Ziarno, M., Żbikowska, A. and Majewska, E. (2021). Phenolic Contents and Antioxidant Activity of Extracts of Selected Fresh and Dried Herbal Materials. *Polish Journal of Food and Nutrition Sciences*, 71(3), pp.269–278.
doi:<https://doi.org/10.31883/pjfn/139035>.
30. Olorunnisola, O.S., Bradley, G. and Afolayan, A.J. (2011). Antioxidant Properties and Cytotoxicity Evaluation of Methanolic Extract of Dried and Fresh Rhizomes of *Tulbaghia violacea*. *African Journal of Pharmacy and Pharmacology*, 5(22), pp.2490–2497.
doi:<https://doi.org/10.5897/AJPP11.620>.
31. Krakowska-Sieprawska, A., Kielbasa, A., Rafińska, K., Ligor, M. and Buszewski, B. (2022). Modern Methods of Pre-Treatment of Plant Material for the Extraction of Bioactive Compounds. *Molecules*, 27(3), p.730. doi:<https://doi.org/10.3390/molecules27030730>.
32. Speight, J.G. (2018). Mechanisms of Introduction into the Environment. *Reaction Mechanisms in Environmental Engineering*, pp.115–161. doi:<https://doi.org/10.1016/b978-0-12-804422-3.00004-3>.
33. Horablaga, N.M., Cozma, A., Alexa, E., Obistioiu, D., Cocan, I., Poiana, M.-A., Lalescu, D., Pop, G., Imbrea, I.M. and Buzna, C. (2023). Influence of Sample Preparation/Extraction Method on the Phytochemical Profile and Antimicrobial Activities of 12 Commonly Consumed Medicinal Plants in Romania. *Applied Sciences*, 13(4), pp.1–25.
doi:<https://doi.org/10.3390/app13042530>.
34. Altemimi, A., Lakhssassi, N., Baharlouei, A., Watson, D. and Lightfoot, D. (2017). Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants*, [online] 6(4), p.42. doi:<https://doi.org/10.3390/plants6040042>.

35. Majekodunmi, S.O. (2015). Review of Extraction of Medicinal Plants for Pharmaceutical Research. *Merit Research Journal of Medicine and Medical Sciences*, 3(11), pp.521–527.
36. Das, K., Tiwari, R.K.S. and Shrivastava, D.K. (2010) Techniques for Evaluation of Medicinal Plant Products as Antimicrobial Agent: Current Methods and Future Trends. *Journal of Medicinal Plants Research*, 4, 104-111.
37. Tiwari, P., Kumar, B., Kaur, M., Kaur, G. and Kaur, H. (2011). Phytochemical Screening and Extraction: A Review. *Internationale Pharmaceutica Scientia*, 1(1), pp.98–106.
38. Cowan, M.M. (1999). Plant Products as Antimicrobial Agents. *Clinical Microbiology Reviews*, 12(4), pp.564–582. doi:<https://doi.org/10.1128/cmr.12.4.564>.
39. Mworia, J.K., Kibiti, C.M., Ngugi, M.P. and Ngeranwa, J.N. (2019). Antipyretic Potential of Dichloromethane Leaf Extract of *Eucalyptus globulus* (Labill) and *Senna didymobotrya* (Fresenius) in Rats Models. *Heliyon*, 5(12), p.e02924. doi:<https://doi.org/10.1016/j.heliyon.2019.e02924>.
40. Chaugule, A., Patil, H., Pagariya, S. and Ingle, P. (2019). Extraction of Caffeine. *International Journal of Advanced Research in Chemical Science*, [online] 6(9), pp.11–19. doi:<https://doi.org/10.20431/2349-0403.0609002>.
41. Bell-Young, L. (2017). *What Is Dichloromethane?* [online] ReAgent Chemicals. Available at: <https://www.chemicals.co.uk/blog/what-is-dichloromethane>.
42. Hamuel, J. (2012). Phytochemicals: Extraction Methods, Basic Structures and Mode of Action as Potential Chemotherapeutic Agents. In: *Phytochemicals - a Global Perspective of Their Role in Nutrition and Health*. [online] doi:<https://doi.org/10.5772/26052>.
43. Bhan, M. (2017). Ionic Liquids as Green Solvents in Herbal Extraction. *International Journal of Advanced Research and Development.*, 2, pp.10-12.
44. Eloff, J.N. (1998). Which Extractant Should Be Used for the Screening and Isolation of Antimicrobial Components from plants? *Journal of Ethnopharmacology*, 60, pp.1–8.

45. Ujang, Z.B., Subramaniam, T., Diah, M.M., Wahid, H.B., Abdullah, B.B., Rashid, A.H.B.A. and Appleton, D. (2013). Bioguided Fractionation and Purification of Natural Bioactives Obtained from *Alpinia conchigera* Water Extract with Melanin Inhibition Activity. *Journal of Biomaterials and Nanobiotechnology*, [online] 04(03), pp.265–272.
doi:<https://doi.org/10.4236/jbnb.2013.43033>.
46. Bimakr, M., Rahman, R.A., Taip, F.S., Ganjloo, A., Salleh, L.M., Selamat, J., Hamid, A. and Zaidul, I.S.M. (2011). Comparison of Different Extraction Methods for the Extraction of Major Bioactive Flavonoid Compounds from Spearmint (*Mentha spicata* L.) Leaves. *Food and Bioproducts Processing*, 89(1), pp.67–72. doi:<https://doi.org/10.1016/j.fbp.2010.03.002>.
47. Hossain, M.A., Al-Hdhrami, S.S., Weli, A.M., Al-Riyami, Q. and Al-Sabahi, J.N. (2014). Isolation, Fractionation and Identification of Chemical Constituents from the Leaves Crude Extracts of *Mentha piperita* L. Grown in Sultanate of Oman. *Asian Pacific Journal of Tropical Biomedicine*, [online] 4, pp.S368–S372.
doi:<https://doi.org/10.12980/apjtb.4.2014c1051>.
48. Harborne, A.J., 1998. *Phytochemical methods a guide to modern techniques of plant analysis*. springer science & business media.
49. Sapkale, G.N., Patil, S.M., Surwase, U.S. and Bhatbhage, P.K. (2010). Supercritical Fluid Extraction- A Review. *International Journal Chemical Sciences*, 8(2), pp.729–743.
50. Ma, Y., Meng, A., Liu, P., Chen, Y., Yuan, A., Dai, Y., Ye, K., Yang, Y., Wang, Y. and Li, Z. (2022). Reflux Extraction Optimization and Antioxidant Activity of Phenolic Compounds from *Pleuroblastusamarus* (Keng) Shell. *Molecules*, 27(2), p.362.
doi:<https://doi.org/10.3390/molecules27020362>.
51. Chua, L.S., Latiff, N.A. and Mohamad, M. (2016). Reflux extraction and cleanup process by column chromatography for high yield of andrographolide enriched extract. *Journal of Applied Research on Medicinal and Aromatic Plants*, [online] 3(2), pp.64–70.
doi:<https://doi.org/10.1016/j.jarmap.2016.01.004>.
52. Bennour, N., Mighri, H., Eljani, H., Zammouri, T. and Akrou, A. (2020). Effect of Solvent Evaporation Method on Phenolic Compounds and the Antioxidant Activity of *Moringa*

- oleifera* Cultivated in Southern Tunisia. *South African Journal of Botany*, 129, pp.181–190.
doi:<https://doi.org/10.1016/j.sajb.2019.05.005>.
53. Mahour, K. (2016). Effect of Solvent Evaporation Methods on Biological Activity of Crude Extract of *Feronia elephantum* Correa Leaves. *Medicinal and Aromatic plants*, 2(2), pp.111–116. doi:<https://doi.org/10.48347/imist.prsm/ajmap-v2i2.6707>.
54. WkieLab.com. (2020). *Rotary Evaporator Parts and Functions - WKIE LAB*. [online] Available at: <https://wkielab.com/rotary-evaporator-parts-and-their-functions/>.
55. O’Driscoll, A. (2020). Putting the Rule of 20 into Practice | Blog | Rotovaps.net. [online] rotovaps.net. Available at: <https://rotovaps.net/blogs/blog/putting-the-rule-of-20-into-practice#:~:text=In%20some%20cases%2C%20the%2020> [Accessed 5 May 2023].
56. Laher, F., Aremu, A.O., Van Staden, J. and Finnie, J.F. (2013). Evaluating the Effect of Storage on the Biological Activity and Chemical Composition of Three South African Medicinal Plants. *South African Journal of Botany*, 88, pp.414–418.
doi:<https://doi.org/10.1016/j.sajb.2013.09.003>.
57. Hartmann, J. and Asch, F. (2019). Extraction, Storage Duration, and Storage Temperature Affect the Activity of Ascorbate Peroxidase, Glutathione Reductase, and Superoxide Dismutase in Rice Tissue. *Biology*, 8(4), p.70. doi:<https://doi.org/10.3390/biology8040070>.
58. Awaad, Amani. (2014). Re: How long can we store plant extracts (solvent extracted) before using for bioassays?. Retrieved from:
https://www.researchgate.net/post/How_long_can_we_store_plant_extract_solvent_extracted_before_using_for_bioassays/5454ea9cd3df3ebc7b8b4760/citation/download.
59. Sivakumar, V. (2014). Re: How long can we store plant extracts (solvent extracted) before using for bioassays? Retrieved from:
https://www.researchgate.net/post/How_long_can_we_store_plant_extract_solvent_extracted_before_using_for_bioassays/54534c33d5a3f25a1f8b457b/citation/download.
60. Azizi, Majid. (2014). Re: How long can we store plant extracts (solvent extracted) before using for bioassays? Retrieved from:
https://www.researchgate.net/post/How_long_can_we_store_plant_extract_solvent_extracted_before_using_for_bioassays/54579a00d3df3e19268b45c7/citation/download.
61. Chaurasiya, Narayan D. (2014). Re: How long can we store plant extracts (solvent extracted) before using for bioassays? Retrieved from:

[https://www.researchgate.net/post/How long can we store plant extracts solvent extracted before using for bioassays/545bcc9dd3df3e37178b45aa/citation/download](https://www.researchgate.net/post/How_long_can_we_store_plant_extract_solvent_extracted_before_using_for_bioassays/545bcc9dd3df3e37178b45aa/citation/download).

62. Cioanca, Oana. (2014). Re: How long can we store plant extracts (solvent extracted) before using for bioassays? Retrieved from:
[https://www.researchgate.net/post/How long can we store plant extracts solvent extracted before using for bioassays/5461dfa9d685cc24718b4693/citation/download](https://www.researchgate.net/post/How_long_can_we_store_plant_extract_solvent_extracted_before_using_for_bioassays/5461dfa9d685cc24718b4693/citation/download).
63. Jha, A.K. and Sit, N. (2021). Extraction of Bioactive Compounds from Plant Materials Using Combination of Various Novel methods: a Review. Trends in Food Science & Technology. doi:<https://doi.org/10.1016/j.tifs.2021.11.019>.
64. Mukherjee, P.K. (2019). Chapter 6 - Extraction and Other Downstream Procedures for Evaluation of Herbal Drugs. [online] ScienceDirect. Available at:
<https://www.sciencedirect.com/science/article/pii/B9780128133743000065> [Accessed 18 Mar. 2021].
65. Aryal, D. (2022). *Soxhlet Extraction: Principle, Extraction procedure, and Apparatus - Chemistry Notes*. [online] <https://chemistnotes.com/>. Available at:
<https://chemistnotes.com/organic/soxhlet-extraction-principle-extraction-procedure-and-apparatus/>.
66. Kusuma, H.S. and Mahfud, M. (2016). Preliminary study: Kinetics of Oil Extraction from Sandalwood by microwave-assisted Hydrodistillation. IOP Conference Series: Materials Science and Engineering, 128, p.012009. doi:<https://doi.org/10.1088/1757-899x/128/1/012009>.
67. Shaterabadi, D., Aboonajmi, M., Ghorbani Javid, M. and Arabhosseini, A. (2020). Effect of Power Ultrasound on the Extraction of Black Caraway (*Carum carvi* L.) and Evaluation of Their Qualitative Properties Using Response Surface Methodology. Food Science & Nutrition, [online] 8(8), pp.4361–4369. doi:<https://doi.org/10.1002/fsn3.1733>.
68. Zentou, H., Abidin, Z., Yunus, R., Awang Biak, D. and Korelskiy, D. (2019). Overview of Alternative Ethanol Removal Techniques for Enhancing Bioethanol Recovery from Fermentation Broth. *Processes*, 7(7), p.458. doi:<https://doi.org/10.3390/pr7070458>.
69. Aleksic, V. and Knezevic, P. (2014). Antimicrobial and antioxidative activity of extracts and essential oils of *Myrtus communis* L. *Microbiological Research*, [online] 169(4), pp.240–254. doi:<https://doi.org/10.1016/j.micres.2013.10.003>.

70. Markom, M., Singh, H. and Hasan, M. (2001). Supercritical CO₂ Fractionation of Crude Palm Oil. *The Journal of Supercritical Fluids*, 20(1), pp.45–53.
doi:[https://doi.org/10.1016/s0896-8446\(00\)00104-2](https://doi.org/10.1016/s0896-8446(00)00104-2).
71. Aditha, S.K., Kurdekar, A.D., Chunduri, L.A.A., Patnaik, S. and Kamiseti, V. (2016). Aqueous Based Reflux Method for Green Synthesis of nanostructures: Application in CZTS Synthesis. *MethodsX*, 3, pp.35–42. doi:<https://doi.org/10.1016/j.mex.2015.12.003>.
72. WkieLab.com. (2020). Rotary Evaporator Parts and Functions - WKIE LAB. [online] Available at: <https://wkielab.com/rotary-evaporator-parts-and-their-functions/>.

UNDER PEER REVIEW