

Original Research Article

ASSESSMENT OF PHYSICO-CHEMICAL PROPERTIES OF SOIL FROM DIFFERENT DEPARTMENTS OF NAI, SHUATS, PRAYAGRAJ U.P.

Abstract

An Assessment of Physico-chemical properties of soil from different departments of NAI, SHUATS, Prayagraj carried out 2022. The prime objectives of this study were to carry out the physico-chemical properties of soil at different depths of various department research farms to determine the availability of macronutrients and micronutrients in soil of these soils sample provide the assessment 6 sampling locations were selected. Soil samples were collected with depth of 0-15 cm and 15-30 cm respectively. The soil colour dry condition varied from light yellowish brown, pale olive, light olive brown, yellowish brown, olive yellow, dark brown and in wet condition varied from olive brown, dark brown, olive yellow and dark yellowish. The sand, silt and clay varied from 53 to 65 %, 19 to 27 % and 15 to 20 % respectively, soil texture was dominantly sandy loam in every site. The bulk density ranged from 1.20 to 1.34 Mgm^{-3} . The practical density ranged from 2.68 to 2.94 Mgm^{-3} . The pore space ranged from 43.25 to 49.50%. The water holding capacity ranged from 41.75 to 48.25 %. The soil pH ranged from 7.37 to 7.71. The electrical conductivity ranged from 0.21 to 0.48 dS m^{-1} . The soil organic carbon ranged from 0.32 to 0.45 %. The available nitrogen ranged from 235.17 to 261.12 kg ha^{-1} . The available phosphorus ranged from 19.66 to 25.50 kg ha^{-1} . The available potassium ranged from 152.00 to 214.00 kg ha^{-1} . The available Sulphur ranged from 8.80 to 14.42 mg kg^{-1} . The available zinc ranged from 0.58 to 0.93 mg kg^{-1} . The available boron ranged from 0.58 to 1.10 mg kg^{-1} . The available iron ranged from 5.92 to 8.70 mg kg^{-1} . The available manganese ranged from 3.21 to 3.61 mg kg^{-1} . The available copper ranged from 2.00 to 2.94 mg kg^{-1} . There is an including awareness of the need to pay greater attention in the role of macronutrient enhancement in the soil for good soil health and proper nutrition of plants so as to attain optimum economic yield and soil is suitable for all major tropical and sub-tropical crops.

Key words: Physico-chemical properties, nutrients distribution, soil health etc.

INTRODUCTION

Soil and land, though related, are two different entities; Land is two dimensional entity representing geographical area and landscapes, while soil is a three-dimensional body with length, breadth and depth and is hidden below the land surface. It is largely hidden from the outside world until it is lost and goes out of the site **Kanwar (2002)**. Soils are formed by weathering processes of rocks which in the words of Hans Jenny can be depicted by the equation: $S = f(cl, r, p, o, t)$; where S soil, cl climate (rainfall, temperature, snow, *etc.*), r relief or topography, p parent material (rocks, minerals and geological formation), and o organisms - plants and animals, and t time or age. Some of these factors of soil formation are passive but the organisms (biosphere) and man are most active in the transformation of soils and in modifying its properties **Jenny (1994)**. Soil organic matter undergoes mineralization and releases substantial quantities of nitrogen, phosphorus, sulphur and smaller amount of micronutrients **Rahman et al. (2013)**. In most cases, the rapid loss of soil organic matter during the years immediately following the conversion was replaced by slower, but continuing declines due to inappropriate agronomic practices. Long-term records show soil nitrogen content falling by 25–70 % over periods ranging from 30 to 90 years; these records also show soil carbon declining by up to 50% over similar time spans **Bahadur et al. (2015)**. The quality of soil is rather dynamic and can affect the sustainability and productivity of land use. It is the end product of soil derivative and is controlled by chemical, physical, and biological components of a soil and their interactions, however, will vary according to the location, and the level of sophistication at which measurements are likely to be made **Riley (2001)**. Soil testing makes complete nutrient control possibility Fertilizer experiments are being patterned to determine economically optimum rates of nutrients application high yields with low production costs per unit area must in modern farming. Farmers of today are different in the failure is more certain and sooner unless they are obtaining reasonably high yields, improved drainage, many improved Cultural practices, disease have helped to set the stage for high yields. Soil provides food, fuel and fodder for meeting the needs of human and animal.

MATERIAL AND METHOD

Analysis of the soil samples were under the methods: Soil textural class was determined by using hydrometer **Bouyoucos (1927)**. Bulk density, particle density, water holding capacity was determined by using graduated measuring cylinder method **Muthuavalet et al. (1992)**. pH was estimated with the help of digital pH meter after making 1:2.5 soil water suspension **Jackson (1958)**. Electrical conductivity was estimated with the help of digital conductivity meter **Wilcox, (1950)**. Percent organic carbon was estimated by wet oxidation method **Walkley and Black (1947)**. Available nitrogen was estimated by alkaline potassium permanganate method, using Kjeldahl apparatus **Subbiah and Asija (1956)**. Available phosphorus was estimated by photoelectric colorimeter method **Olsen's et al. (1945)**. Available potassium was estimated by neutral normal ammonium acetate extraction followed by flame photometric method **Toth and Prince (1949)**. Available sulphur was estimated by reduction method **Johnson and Nishita (1952)**. Available boron estimation by **Kmiecik et al. (2016)** and copper, iron, manganese, zinc estimated by **Lindsay and Novell (1978)**.

RESULT AND DISCUSSION

Physical Properties

The soil colour dry condition varied from light yellowish brown, pale olive, light olive brown, yellowish brown, olive yellow, dark brown and in wet condition varied from olive brown, dark brown, olive yellow and dark yellowish. The soil textural classes identified as sandy loam. The sand, silt and clay varied from 53 to 65 % sand, 19 to 27 % silt and 15 to 20 % clay in Sandy Loam. Minimum bulk density was recorded 1.20 Mg m^{-3} in D₄- horticulture at 0-15 cm, and maximum 1.34 Mg m^{-3} in D₆- agro forestry at 15-30 cm. Minimum particle density was recorded 2.68 Mg m^{-3} in D₄- horticulture at 0-15 cm, and maximum 2.94 Mg m^{-3} in D₆- agro forestry at 15-30 cm. Minimum pore space was recorded 43.25 % in D₁- soil science and agricultural chemistry at 15-30 cm, and maximum 49.50 % in D₅- plant protection at 0-15 cm. Minimum water holding capacity was recorded 41.75 % in D₁- soil science and agricultural chemistry at 15-30 cm, and maximum 48.25 % in D₄- horticulture at 0-15 cm.

Macronutrients

Minimum nitrogen was recorded 261.12 kg ha⁻¹ in D₄- horticulture at 0-15 cm, and maximum 235.17 kg ha⁻¹ in D₁- soil science and agricultural chemistry at 15-30 cm. Minimum phosphorus was recorded 25.50 kg ha⁻¹ in D₄- horticulture at 0-15 cm, and maximum 19.66 kg ha⁻¹ in D₁- soil science and agricultural chemistry at 15-30 cm. Minimum potassium was recorded 214.00 kg ha⁻¹ in D₅- plant protection at 0-15 cm, and maximum 152.00 kg ha⁻¹ in D₃- genetic and plant breeding at 15-30 cm. Minimum sulphur was recorded 14.42 mg kg⁻¹ in D₂- agronomy at 0-15 cm, and maximum 8.80 mg ka⁻¹ in D₅- plant protection at 15-30 cm.

Table: 3. Evaluation of nitrogen, phosphorus, potassium and sulphur in various depths at different research farm of NAI, SHUATS, Prayagraj

Department	Nitrogen (kg ha ⁻¹)		Phosphorus (kg ha ⁻¹)		Potassium (kg ha ⁻¹)		Sulphur (mg kg ⁻¹)	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
D ₁ -Soil Science	242.24	235.17	22.32	19.84	195.00	182.00	12.36	11.90
D ₂ -Agronomy	253.84	242.37	23.14	20.86	203.00	188.00	14.42	13.96
D ₃ -GPB	246.85	242.44	24.21	21.18	166.00	152.00	10.36	08.96
D ₄ - Horticulture	261.12	250.26	25.50	21.82	192.00	177.00	13.26	11.20
D ₅ -Plant Protection	250.54	245.26	23.28	20.57	214.00	194.00	09.67	08.80
D ₆ - Agro-Forestry	248.30	241.43	22.86	19.66	171.00	164.00	11.26	09.96
	Depth	Site	Depth	Site	Depth	Site	Depth	Site
S. Ed. (±)	7.893	1.870	1.353	0.331	24.049	2.708	2.632	0.400
C.D. at 5%	16.732	3.964	2.868	0.701	50.983	5.741	5.581	0.849
F-test	S	S	S	S	S	S	S	S

Micronutrients

Minimum zinc was recorded 0.62 mg kg⁻¹ in D₆- agro forestry at 15-30 cm, and maximum 0.93 mg ka⁻¹ in D₄- horticulture at 0-15 cm. Minimum copper was recorded 2.00 mg kg⁻¹ in D₁- soil science and agricultural chemistry at 15-30 cm, and maximum 2.94 mg ka⁻¹ in D₂- agronomy at 0-15 cm. Minimum manganese was recorded 3.21 mg kg⁻¹ in D₁- soil science and agricultural chemistry at 15-30 cm, and maximum 3.61 mg ka⁻¹ in D₃- genetic and plant breeding at 0-15 cm. Minimum boron was recorded 0.58 mg kg⁻¹ in D₁- soil science and agricultural chemistry at 15-30 cm, and maximum 1.10 mg ka⁻¹ in D₅- D₂- agronomy at 0-15 cm. Minimum iron was recorded 5.92 mg kg⁻¹ in D₁- soil science and agricultural chemistry at 15-30 cm, and maximum 8.70 mg ka⁻¹ in D₄- horticulture at 0-15 cm.

Table: 4. Evaluation of zinc, copper, manganese, boron and iron in various depths at different research farm of NAI, SHUATS, Prayagraj

Department	Zinc (kg ha ⁻¹)		Copper (kg ha ⁻¹)		Manganese (kg ha ⁻¹)		Boron (kg ha ⁻¹)		Iron (kg ha ⁻¹)	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
D ₁ -Soil Science	0.74	0.69	2.20	2.00	3.24	3.21	0.64	0.58	6.10	5.92
D ₂ -Agronomy	0.89	0.78	2.94	2.86	3.28	3.24	1.10	0.94	8.14	8.03
D ₃ -GPB	0.67	0.58	2.30	2.26	3.61	3.59	0.98	0.78	6.90	6.55
D ₄ -Horticulture	0.93	0.81	2.08	2.02	3.57	3.54	1.04	0.94	8.70	7.90
D ₅ -Plant Protection	0.86	0.77	2.24	2.18	3.49	3.46	0.88	0.62	7.70	7.28
D ₆ -Agro Forestry	0.81	0.62	2.06	2.00	3.54	3.48	0.64	0.60	7.45	7.26
	Depth	Site	Depth	Site	Depth	Site	Depth	Site	Depth	Site
S. Ed. (±)	0.131	0.030	0.463	0.038	0.222	0.009	0.254	0.055	1.226	0.197
C.D. at 5%	0.278	0.064	0.981	0.080	0.471	0.019	0.538	0.117	2.599	0.417
F-test	S	S	S	S	S	S	S	S	S	S

CONCLUSION

It concluded that soil of all research farm of Naini Agricultural Institute have low level in organic carbon and nitrogen but medium in phosphorous and potassium content. Thus balance nutrient additions through organic manures, inorganic and bio-fertilizer sources are essential to maintain soil fertility in research farms of Naini Agricultural Institute, SHUATS.

REFERANCE

Bahadur Indra, Sonkar V. K., Kumar Sanjay, Dixit Jyoti and Singh Abhishek

Pratap(2015)Crop Residue Management for improving Soil Quality and Crop

Productivity in India. *Indian Journal of Agriculture and Allied Sciences*, **1(1):2395-1109.**

- Bouyoucos, G.J. (1927)** The Hydrometer as a new method for the mechanical analysis of soils, *Soil Science*, 23: 343-353
- Fisher, R. A. (1960)** Statistical Methods and Scientific Induction, *Journal of the Royal Statistical Society Series B*, 17: 69-78.
- Jackson, M.L. (1973)** Chemical Analysis of Soil pH. Prentice Hall of India Pvt. Ltd., New Delhi.
- Jenny, H. (1994)** Factors of Soil Formation: A System of Quantitative Pedology. Dover Publication, New York, 281.
- Kanwar, J.S. (2002)** Introduction, Fundamental of soil science *Indian Society of Soil Science*, (ISBN 81-901470-0-5), 1-4
- Lindsay, W.L. and Norvell, W.A. (1978)** Development of a DPTA soil test for heavy metals. *Soil Science Society of America Journal*, 42:402-403.
- Munsell, A.H. (1954)** Munsell Soil Color Charts, Munsell Color Company Inc., Baltimore.
- Rahman, M.H., Islam, M.R., Jahiruddin, M., Puteh, A.B. and Mondal, M.M.A. (2013)** Influence of Organic Matter on Nitrogen Mineralization Pattern in Soils under Different Moisture Regimes. *International Journal of Agriculture and Biology*, 15:55-61.
- Riley, J. (2001)** The Indicator explosion: local needs and international challenges. *Agriculture, Ecosystems, and Environment* 87: 119-120.
- Subbiah, B.V. and Asija, C.L. (1956)** A Rapid procedure for the estimation of available nitrogen in soils. *Current Sci.*, 25:259-260.
- Toth, S.J. and Prince, A.L. (1949)** Estimation of Potassium content of soil by flame photometer technique. *Soil Sci.*, 67: 439-445.
- Walkley, A. (1947)** Critical Examination of rapid method for determining organic carbon in soils, *Soil Sci.*, 63: 251.
- Wilcox, L.V. (1950)** Electrical conductivity, *Amer. Water Works Assoc. J.*, 42:775-776.