

*Original Research Article*

**INTAKE OF FRUITS AND VEGETABLES IN SELF PRESCRIBED WEIGHT REDUCING AND DASH DIETS AMONG OBESEHYPERTENSIVE INDIVIDUALS ATTENDING IRRUA SPECIALIST TEACHING HOSPITAL IRRUA, EDO STATE**

**ABSTRACT**

**Background:** Five servings of fruits and vegetables each day, along with relatively unprocessed whole grains or legumes with each meal, are essential. While being low or relatively low in calories, fruits and vegetables include a large amount of dietary fiber and a range of micronutrients, antioxidants and phytochemicals which are all essential for health.

**Aims:** To investigate the intake of fruits and vegetables in self prescribed weight reducing and DASH (Dietary Approach to Stop Hypertension) diets among obese-hypertensive individuals attending Irrua Specialist Teaching Hospital Irrua, Edo State.

**Methods:** A descriptive cross sectional study design was used with a sample size of 440 obese hypertensive individuals. Questionnaires, personal interviews and anthropometric measurements were used to collect data using purposive sampling technique. Data collected was were analyzed using Statistical Package for the Social Sciences (IBM SPSS) 22 Version.

**Results:** A total of 440 respondents were used in the study and data were elicited from all of this number giving a 100% response rate. 242 (55.0%) of the respondents consumed fruits only one to two days per week with only 36 (8.2%) consuming fruits every day. 215

**Comment [A1]:** I suggest your title to be rewritten as "INTAKE OF FRUITS AND VEGETABLES AS SELF PRESCRIBED WEIGHT REDUCING AND DIETARY APPROACH TO STOP HYPERTENSION(DASH) DIETS AMONG OBESE-HYPERTENSIVE INDIVIDUALS ATTENDING IRRUA SPECIALIST TEACHING HOSPITAL IRRUA, EDO STATE"

**Comment [A2]:** Are essential for what?

**Comment [A3]:** You need to justify that they were both obese and hypertensive (Obese-hypertensive)

**Comment [A4]:** You are not supposed to start a sentence with figures

(48.9%) consumed fruits one to two days per week, with only 26 (5.9%) consuming vegetables every day.

**Conclusion:** It can be concluded based on the key findings of this study that there is statistically significant mean difference in fruits and vegetable consumption in the management of obese hypertensive amongst individuals attending Irrua Specialist Teaching Hospital, Edo State, Nigeria ( $p < 0.05$ ).

**Keywords:** Fruit, Vegetable, DASH, Obese Hypertensive, Hospital

## INTRODUCTION

In Nigerian marketplaces, certain fruits and vegetables are always easily accessible. Garlic, cashews, almonds, fruits (papaya, mango, avocado, watermelon, etc.), vegetables, potatoes (high in potassium, magnesium, and fiber), fish, bananas (high in potassium), beans and cowpeas (nutritious chock-full of magnesium, soluble fiber, and potassium), fish oil and almonds seem to be affordable and readily available to those with low incomes.<sup>1</sup>

Robert et al., made the following dietary recommendations to control led obesity and hypertension: A diet with a high intake of vegetables, fruits, and whole grains. Other recommendations include consuming low-fat dairy products, poultry, fish, legumes, non-tropical vegetable oils, and nuts; and limiting intake of sweets, sugar-sweetened beverages, and red meat.<sup>2</sup> This dietary pattern should be adapted to appropriate calorie requirements, personal and cultural food preferences, and nutritional therapy for other medical conditions, including diabetes mellitus. One way to achieve this is by following plans such as the Dietary Approaches to Stop Hypertension (DASH) diet.<sup>2</sup> It is critical to consume five

servings of fruits and vegetables per day, as well as relatively unprocessed whole grains or legumes with each meal.<sup>3</sup>

A study by Emefa et al. concludes that increased physical activity, abstaining from alcohol and smoking, increased intake of fruits and vegetables, and reduced intake of carbohydrates, meat and fat have a positive influence on blood pressure control.<sup>4</sup> According to the World Health Organization, potassium is typically found in a variety of unprocessed foods, including fruits and vegetables. It provides foods like cowpeas, pigeon peas, lima beans, and African yam beans, which have a potassium content of 1300 milligrams per 100 grams of fresh weight.<sup>5,6,7</sup>

The present network meta-analysis suggests that ~~the~~ DASH might be the most effective dietary measure to reduce blood pressure among obese hypertensive and pre-hypertensive patients based on high quality evidence.<sup>8</sup> Findings showed that intake of fruits, vegetables and dairy products is associated with a healthier weight status in adults.<sup>9</sup> In another study, Arnotti and Bamber revealed that fruit and vegetable consumption promotes weight loss and prevents weight gain, thereby reducing risks for chronic health conditions.<sup>10</sup>

## **2. MATERIALS AND METHODS**

### **2.1 Study area**

### **2.2 Study population**

#### **2.2.1 Inclusion criteria**

#### **2.2.2 Informed consent statement**

#### **2.2.3 Ethical clearance**

### **2.3 Sampling and sample size**

**Comment [A5]:** What provides foods???

**Comment [A6]:** It is expected that you narrow down to your aim at this juncture. Talk about the aim of the study here please.

**Comment [A7]:** YOU NEED TO RE-WRITE THE WHOLE OF THIS SECTION IN TO SUB-SECTIONS IN A SELF EXPLAINED LANGUAGE. WHAT YOU HAVE HERE UNDER YOUR MATERIALS AND METHOD DOES NOT CONFORM WITH WHAT IS SPECIFIED IN THE AUTHORS GUIDE

**Formatted:** Tab stops: 1.88", Left

#### **2.4 Study tool(s)**

#### **2.5 Data collection**

#### **2.6 Data analysis**

This study complies with a reporting guideline by Lachat et al.<sup>11</sup> A descriptive cross sectional study design was used with a sample size of 440 obese hypertensive individuals. A purposive sampling technique was used in recruiting the study subjects. Questionnaires, personal interview and anthropometric measurements were used to collect data. The questionnaires and the research instruments were approved by the supervisors; they were pre-tested on 40 individuals attending Irrua Specialist Teaching Hospital Irrua; the questionnaires were administered by the researcher after the respondents have signed the written informed consent form; it was ensured that the structured instrument/questionnaire was reliable and valid for the purpose of the research after validation by the supervisors and two experts from the community. The study was carried out by the researcher incorporating other scientists/professionals on the need arises basis. The written informed consent forms were given (and after consenting) was followed by the collection of samples and other research procedures. Data collected was analysed using Statistical Package for the Social Sciences (IBM SPSS) 22 Version and ENA Software Version 2011. Hypotheses were tested using Chi-square, t-test and Pearson Product Moment Correlation Coefficient at 95% confidence level.

Sample size determination:<sup>12</sup>

$$n = Z^2 pq/d^2$$

Z= Level of significant = 1.96<sup>2</sup>

p= Prevalence of indicators= 0.061(6.1%)

q= 1-p

d<sup>2</sup>=Degree of Precision = 0.05<sup>2</sup>

$$n = 1.96^2 \times 0.061 (1-0.061)$$

0.05<sup>2</sup>

n=88.01

n= 88

- Minimum sample size

Sample size of 400

The study area was Irrua Specialist Teaching Hospital located at KM87 Benin - Auchi expressway Edo State. Edo State is one of the 36 states of Nigeria, located in the south – south geopolitical zone of the country. It is bounded by the states of Kogi to the northeast and east, Anambra to the east, Delta to the southeast and south, and Ondo to the west and northwest; the Niger River flows along the state's eastern boundary. Benin City is the state

capital and largest urban center. Edo state lies at elevations between 500 feet (150 m) in the south and more than 1,800 feet (550 m) in the north. Tropical rain forest covers most of the area. The state is inhabited largely by the Edo (Bini) people, who are linked to the historic kingdom of Benin.<sup>13</sup>

Agriculture is the mainstay of the economy. Yams, cassava (manioc), oil palm produce, rice, and corn (maize) are the major subsistence crops, while rubber, timber, and palm oil and kernels are cash crops. Mineral resources include limestone and lignite. Industries produce pharmaceuticals, rubber, plywood, beer, sawn wood, and furniture. A network of trunk roads in the state and an airport at Benin City facilitate transportation. The Nigerian Institute of Oil Palm Research, the Rubber Research Institute of Nigeria, and the University of Benin (founded 1970) are located at Benin City, while a state university (founded 1981) is at Ekpoma. Population (2006) is 3,218,332.<sup>13</sup>

**Comment [A8]:** All these are not needed in your methodology.

The research was limited to individuals attending Irrua Specialist Teaching Hospital Irrua, Edo State who are screened and confirmed to be obese and hypertensive. Irrua Specialist Teaching Hospital is a federal tertiary healthcare institution where people from different communities mostly within Edo State seek for healthcare services.

The study population was the obese hypertensive individuals attending Irrua Specialist Teaching Hospital Irrua, Edo State. Population of the state is projected to be 4,777,000 as at 2022. The estimated obese hypertensive sub-population in the study area is 19%.

A purposive sampling technique was used for this study. The sample size was achieved within the period of six (6) months; it was based on individuals attending Irrua Specialist

Teaching Hospital Irrua, Edo State who were screened and confirmed to be obese and hypertensive.

The study included all obese hypertensive individuals between the ages of 20 to 65 years attending Irrua Specialist Teaching Hospital Irrua, Edo State who gave both verbal and written consent to be part of the research.

Obese and hypertensive that were below 20 years and above 65 years, critically ill and none consenting individuals were excluded from the study.

Questionnaires, personal interviews, instruments for anthropometric, biochemical and clinical assessments were used to collect samples and data. Section “A” of the questionnaire captured the socio-demographic data of the individuals while section “B” the nutritional status of the obese hypertensive individuals attending the Irrua Specialist Teaching Hospital, Edo State. 8mls of blood was drawn from each of the participant for the nutritional biochemistry analysis (lipid profile, electrolytes, urea and creatinine, fasting/random glucose level).

It was ensured that the structured instrument/questionnaire was pre-tested and valid for the purpose of the research after validation by the research supervisors and two experts from the department of community and public health.

The instrument/questionnaires were approved by the supervisors; they were pre-tested on 40 obese hypertensive individuals in Irrua community, Edo State; the questionnaires were administered by the researcher after the written informed consent form had been signed.

A follow-up study design could be much better to study the nutritional status of obese hypertensive individuals attending Irrua Specialist Teaching Hospital, Edo State.

**Comment [A9]:** You didn't report this work in your result nor did it form any part of your discussion. Please add the result to the result section and discuss where appropriate.

Ethical Consideration: The study was conducted according the National Code of Health Research Ethics in Nigeria (the CODE) and the Declaration of Helsinki. Every other institutional guideline was followed.<sup>14</sup>The study was approved by institutional/local review board) before the field visit. Appropriate culture-sensitive and specific written informed consent was sought and obtained from the parents of the children prior to participants' recruitment. The researcher ensured privacy in handling the participants and confidentiality in handling the data. All examinations were done in a safe, comfortable and private environment and the questionnaire was anonymous; the collected data was stored in a passworded computer only available to the principal investigator on a need to know basis. The data is available for inspection by regulatory authorities and for quality assurance for ten years.

Analysis of data was based on information obtained from the questionnaires, anthropometric measurement, biochemical information, clinical examination and personal interviews that was computed and presented in frequency distribution tables and histogram. Statistical Package for the Social Sciences (IBM SPSS) 22 Version and ENA Software Version 2011 will be used for analysing the data. Hypotheses were tested using Chi-square, t-test and Pearson Product Moment Correlation Coefficient at 95% confidence level.<sup>15</sup>

## **RESULTS**

A total of 440 respondents/participants were used in the study and data were elicited from all of this number giving a 100% response rate. Out of the 440 (100%) participants, 133 (30.2%) were males while 307 (69.8%) were females. The age distribution shows that the highest proportion 424 (96.4%) were 31 years and above, 13 (3.0%) were within the age

group 21-25 years, 3 (0.7%) were within the age bracket 26-30 years while none were recorded for age group 15 – 20 years. The mean age for the respondents was 33 years. Majority of the study participants 348 (79.1%) were married, followed by 40 (9.1%) who were widowed, 39 (8.9%) were single while 13 (3.0%) were divorced. Majority 409 (93.0%) of the respondents subscribed to Christianity, while 30 (6.8%) subscribed to Islam. Most of the respondents 254 (58.0%) had tertiary education, followed by 92 (20.9%) who had secondary education, 86 (19.5%) had primary education while 7 (1.6%) had no formal education. Few of the respondents 46 (10.5%) were into business as their occupation, followed by 43 (9.8%) who were civil servants and 13 (3.0%) who were farmers. Details of this result are presented in Table 1.

UNDER PEER REVIEW

**TABLE 1. Socio-demographic characteristics of respondents (N=440)**

<b>Variables</b>	<b>Frequency</b>	<b>Percent</b>	<b>Mean</b>	<b>Standard Deviation</b>
<b>Sex of respondents</b>			1.70	0.460
Male	133	30.2%		
Female	307	69.8%		
<b>Age of respondents</b>			*33.37	2.503
15-20 years	0	0.0%		
21-25 years	13	3.0%		
26-30 years	3	0.7%		
31 years and above	424	96.4%		
<b>Marital status</b>			2.12	0.684
Single	39	8.9%		
Married	348	79.1%		
Divorced	13	3.0%		
Widowed	40	9.1%		

<b>Religious affiliation</b>			1.07	0.269
Christianity	409	93.0%		
Islam	30	6.8%		
Others	1	0.2%		
<b>Educational status</b>			2.35	0.846
No formal education	7	1.6%		
Primary level of education	86	19.5%		
Secondary level of education	92	20.9%		
Tertiary level of education	254	58.0%		
<b>Occupation</b>			3.24	1.815
Civil servants	34	9.8%		
Housewives	5	1.1%		
Farmers	13	3.0%		
Business Women/men	46	10.5%		
Applicants	4	0.9%		
Others	23	5.2%		

---

\*Mean age

**Anthropometric status data**

**Comment [A10]:** From the title of the study, it is expected that all the respondents were obese and hypertensive individuals. But you didn't report at any point where respondents were proven hypertensive. Please provide justification for that.

On account to determine the Body Mass Index (BMI) of respondents, 234 (53.2%) were within 30-34.9 (class 1 obesity), 127 (28.9%) were within 35-39.9 (class 2 obesity) while 79 (18.0%) were within 40 and above (class 3 obesity). The waist circumference of the male response was 7 (1.6%), 26 (5.9%), and 92 (20.9%) for < 94cm (Normal), 94-102cm (High), and > 102cm (Very high) respectively while the waist circumference for the female respondents was 0 (0.0%), 10 (2.3%) and 305 (69.3%) for  $\leq$  80.9cm (Normal), 81-88.9cm (High), and  $\geq$  90cm (Very high) respectively (Table 2).

UNDER PEER REVIEW

**TABLE 2. Anthropometric status data (N=440)**

<b>Variables</b>	<b>Frequency</b>	<b>Percent</b>	<b>Mean</b>	<b>Standard Deviation</b>
<b>Body Mass Index (BMI)</b>			1.65	0.767
30-34.9 (Class 1 obesity)	234	53.2%		
35-39.9 (Class 2 obesity)	127	28.9%		
40 & above (Class 3 obesity)	79	18.0%		
<b>Waist circumference in males</b>			2.68	0.576
≤ 94cm (Normal)	7	1.6%		
95-101.9cm (High)	26	5.9%		
≥ 102cm (Very high)	92	20.9%		
<b>Waist circumference in females</b>			2.97	0.176
≤ 80.9cm (Normal)	0	0.0%		
81-88.9cm (High)	10	2.3%		
≥ 90cm (Very high)	305	69.3%		

### **Consumption of fruits, vegetable, nuts and DASH diet**

The frequency of fruits intake for a greater proportion of the respondents 242 (55.0%) was one to two days per week, followed by those 96 (21.8%) whose frequency was 3-4 days per week, and 66 (15.0%) had it for 5-6days/week while only 36 (8.2%) had a frequency of fruits intake to be every day. Most of the respondents 377 (85.7%) had fresh fruits, 10 (2.3%) had dried or processed fruits while 53 (12.0%) had both. The frequency of vegetable intake for most respondents 215 (48.9%) was one to two days per week, followed by those 149 (33.9%) whose frequency was 3-4 days per week, and 50 (11.4%) had it for 5-6days/week while only 26 (5.9%) had a frequency of vegetables intake to be every day. Majority of the respondents 393 (89.3%) had fresh vegetables, 7 (1.6%) had dried or processed vegetables while 40 (9.1%) had both. Majority 377 (85.7%) had less than 400 grams and few 63 (14.3%) had 400 grams of fruits and vegetables per day while none had above 400 grams (Table 3).

**TABLE 3. Consumption of fruits, vegetable, nuts and DASH diet (N=440)**

<b>Variables</b>	<b>Frequency</b>	<b>Percent</b>	<b>Mean</b>	<b>Standard Deviation</b>
<b>Frequency of fruits intake</b>			1.76	0.987
1-2 days/week	242	55.0%		
3-4 days/week	96	21.8%		
5-6 days/week	66	15.0%		
Every day	36	8.2%		
<b>Forms of fruits</b>			1.26	0.660
Fresh	377	85.7%		
Dried/processed	10	2.3%		
Both	53	12.0%		
<b>Frequency of vegetable intake</b>			1.74	0.880
1-2 days/week	215	48.9%		
3-4 days/week	149	33.9%		
5-6 days/week	50	11.4%		
Every day	26	5.9%		
<b>Forms of vegetables</b>			1.20	0.584
Fresh	393	89.3%		
Dried/processed	7	1.6%		
Both	40	9.1%		

<b>Quantity of fruits/vegetables consumed</b>			1.14	0.351
<b>daily</b>	377	85.7%		
Less than 400 Grams	63	14.3%		
400 Grams	0	0.0%		
More than 400 Grams				
<b>Cereals</b>			2.29	0.915
Whole grains	139	31.6%		
Processed/refined	36	8.2%		
Both	265	60.2%		
<b>Drink</b>			3.20	1.269
1-5% Alcohol	87	19.8%		
6-14% Alcohol	41	9.3%		
15 and above % Alcohol	26	5.9%		
Sugary/Soft/Energy drink	269	61.1%		
None	17	3.8		
<b>Frequency of drinking</b>			1.36	0.703
Occasionally	324	73.6%		
Frequently	92	20.9%		
More frequently	7	1.6%		
None	17	3.9%		

<b>Average salt intake per day</b>			2.21	0.751
1 leveled teaspoon	84	19.1%		
1 heaped teaspoon	181	41.1%		
1.5 teaspoon	172	39.1%		
More than 1.5 teaspoon	3	0.7%		
<b>Pasta</b>			0.91	0.593
None	96	21.8%		
Occasionally	291	66.1%		
Frequently	50	11.4%		
More frequently	3	0.7%		
<b>Nuts and seeds</b>			1.28	0.540
None	20	4.5%		
Occasionally	278	63.2%		
Frequently	142	32.2%		
More frequently	0	0.0%		

---

**Hypothesis:** There is no statistically significant mean difference in fruits, vegetable and nuts consumption in the management of obese hypertensive at what P-Value??? individuals in Edo State, Nigeria. The test statistic used in testing this hypothesis was the t-test at 95 CI (0.05 significance level). Analysis using t-test showed that the means actually differs. In other words, there is a difference. That is, the study was statistically significant (P = 0.0001; df = 439; t-tests = 8.565 & 10.764). Accessing the Confidence intervals (CI = 0.11-0.18 & 0.23-0.33) indicates that the values fell within the rejection region, therefore the null hypothesis was rejected. To this end, the researcher concluded that there is statistically significant mean difference in fruits, vegetable and nuts consumption in the management of obese hypertensive individuals in Edo State, Nigeria (Table 4).

**Comment [A11]:** You are limited to your study area not the entire Edo state

**Comment [A12]:** You don't need this please

**Formatted:** Tab stops: 2.01", Left

UNDER PEER REVIEW

**TABLE 4. t-test analysis of the predictive mean difference in fruits, vegetable and nuts consumption in the management of obese hypertensive individuals in Edo State, Nigeria (N=440)**

Variables	N	Mean	SD	t-test	p-value	CI
Fruits and vegetables consumption	440	1.14	0.351	8.565*	0.000*	0.11 – 0.18
Nuts and seeds consumption	440	1.28	0.540	10.764*		0.23 – 0.33

\*Statistical significance based on  $P$ -value  $< 0.05$  at 95% CI;  $df = 439$

**Comment [A13]:** Limit yourself to your study area

**Comment [A14]:** Socio-demographic data of the respondents were collected and analyzed. Use that result to test whether fruit and vegetable consumption (frequency and quantity) varies in the study population across socio-demographic profiles. One or more of the demographic variable may affect the fruit and vegetable intake

UNDER PEER REVIEW

## DISCUSSION

This study showed that 85.7% of the respondents consume fruits, 48.9% consume vegetables at least one to two days per week as 5.9% consume it frequently every day while 63.2% consume nuts as a means of managing obesity and hypertension. As part of intervention in the management of obesity and hypertension, a study by Emefa et al. concludes that increase in physical activity, abstaining from alcohol and smoking, increased intake of fruits, nuts and vegetables, and reduced intake of carbohydrates, meat, and fat have a positive influence on blood pressure control.<sup>4</sup> To further corroborate this finding, Jevon (2020) found out that, the frequent intake of vegetables, fruits and nuts is important for obese hypertensive individuals in weight reduction, control of blood pressure and helps intervention groups keep fit.<sup>16</sup>

## CONCLUSION

It can be concluded based on the key findings of this study that there is statistically significant association between obesity status by Body Mass Index (BMI) and hypertension among obese hypertensive individuals; there is statistically significant mean difference in fruits and vegetable consumption in the self prescribed weight reducing and DASH diet of obese hypertensive individuals attending Irrua Specialist Teaching Hospital Edo State, Nigeria ( $p < 0.05$ ).

The intake of fruits and vegetables in self prescribed weight reducing and DASH diets are low among obese hypertensive individuals attending Irrua Specialist Teaching Hospital Edo State, Nigeria.

**Comment [A15]:** Your entire discussion is so scanty and skeletal. Discussion entails the interpretation, significance, implication, application, relationship of your findings with similar studies, and significance of your findings. Most recent references are recommended across the entire write up please.

**Comment [A16]:** This is confusing please. You didn't check for hypertension status of your respondents, how did you arrive at this? And if the respondents are already Obese-hypertensive, then why are you testing for association?

**Comment [A17]:** This should be well captured, justified and explained in your discussion. fruits and vegetable consumption has to vary across the demography of the respondents

## REFERENCES

1. Ayo OyozeBaje (2019). Local preservation and packaging of fruits and vegetables. <https://businessday.ng/health/article/local-preservation-and-packaging-of-fruits-and-vegetables/>.
2. Robert H. Eckel, John M. Jakicic, Jamy D. Ard, Janet M. de Jesus, Nancy Houston Miller, Van S. Hubbard, I-Min Lee, Alice H. Lichtenstein, Catherine M. Loria, Barbara E. Millen, Cathy A. Nonas, Frank M. Sacks, Sidney C. Smith, Laura P. Svetkey, Thomas A. Wadden, Susan Z. Yanovski (2013), AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the *American College of Cardiology/American Heart Association Task Force on Practice Guidelines*, <http://circ.ahajournals.org/content/early/2013/11/11/01.cir.0000437740.48606.d1.full.pdf+html>. Accessed July 6, 2021.
3. van den Born B. H., Lip G. Y. H., Brguljan-Hitij J., Cremer A., Segura J., Morales E., Mahfoud F., Amraoui F., Persu A., Kahan T., AgabitiRosei E., de Simone G., Gosse P., Williams B. (2019), ESC Council on hypertension position document on the management of hypertensive emergencies, *Eur Heart J CardiovascPharmacother*; 5:37–46.
4. EmefaModeyAmoah, Darlene EsinamOkai, Adom Manu, Amos Laar, Joseph Akamah, KwasiTorpey (2020), "The Role of Lifestyle Factors in Controlling Blood Pressure among Hypertensive Patients in Two Health Facilities in Urban Ghana: A

Cross-Sectional Study", *International Journal of Hypertension*, Article ID 9379128, 8 pages.

5. World Health Organization (2020). Obesity and overweight, <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed 6th May,2021.
6. World Health Organization (2021), World Obesity Day 2021 aims to encourage practices to achieve and maintain healthy weight, <https://www.firstpost.com/world/world-obesity-day-2021-aims-to-encourage-practices-to-achieve-and-maintain-healthy-weight-9375661.html>. Accessed 6th May,2021.
7. World Health Organization (2016), World Obesity Day: WHO wants governments to get citizens healthy; proposes tax on sugary drinks; <https://www.firstpost.com/living/world-obesity-day-who-wants-governments-to-get-citizens-healthy-proposes-tax-on-sugary-drinks-3048724.html>. Accessed 6th May,2021.
8. Lukas Schwingshackl and Anna Chaimani (2019). Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: A systematic review and network meta-analysis, *Critical Reviews in Food Science and Nutrition*; 59(16),2674-2687.
9. MehranNouri, ZainabShateri and Shiva Faghieh. The relationship between intake of fruits, vegetables and dairy products with overweight and obesity in a large sample in Iran: Findings of STEPS 2016. *Front. Nutr.*, 17 January 2023 Sec. *Nutritional Epidemiology*; Volume 9 - 2022 | <https://doi.org/10.3389/fnut.2022.1082976>.

10. Arnotti K, Bamber M. Fruit and Vegetable Consumption in Overweight or Obese Individuals: A Meta-Analysis. *Western Journal of Nursing Research*. 2020 Apr; 42(4):306-314. doi: 10.1177/0193945919858699. Epub 2019 Jun 29. PMID: 31256714.
11. Lachat C, Hawwash D, Ocké MC, Berg C, Forsum E, Hörnell A, Larsson C, Sonestedt E, Wirfält E, Åkesson A, Kolsteren P, Byrnes G, De Keyzer W, Van Camp J, Cade JE, Slimani N, Cevallos M, Egger M, Huybrechts I. Strengthening the Reporting of Observational Studies in Epidemiology-Nutritional Epidemiology (STROBE-nut): An Extension of the STROBE Statement. *PLoS Med*. 2016 Jun 7;13(6):e1002036. doi: 10.1371/journal.pmed.1002036. PMID: 27270749; PMCID: PMC4896435.
12. Charan J, Biswas T. How to calculate sample size for different study designs in medical research? *Indian J Psychol Med*. 2013 Apr;35(2):121-6. doi: 10.4103/0253-7176.116232. PMID: 24049221; PMCID: PMC3775042.
13. Aakanksha Gaur and Amy Mckenna. Edo State, Nigeria. *Encyclopedia Britannica*. <https://www.britannica.com/place/Edo-state-Nigeria>. Accessed 24th February, 2023.
14. Federal Ministry of Health (2007). National Code of Health Research Ethics. ISBN: 978-978 080-708-5. [http://www.nhrec.net/nhrec/NCHRE\\_July%2007.pdf](http://www.nhrec.net/nhrec/NCHRE_July%2007.pdf).
15. Arifa Rahman and MuktedirMd.Golam (2021). SPSS: An Imperative Quantitative Data Analysis Tool for Social Science Research. DOI:10.47772/IJRISS.2021.51012.<https://www.researchgate.net/publication/35598>

6638\_SPSS\_An\_Imperative\_Quantitative\_Data\_Analysis\_Tool\_for\_Social\_Science  
Research.

16. Jevon P (2020) Blood pressure 2: procedures for measuring blood pressure. Nursing Times [online]; 116: 8, 26-28. Vol 116 Issue 8 [www.nursingtimes.net](http://www.nursingtimes.net).

**Comment [A18]:** Please your reference is not written in accordance with the referencing style of this journal. Ensure that all your references are written in line with the journal guide. You can check the authors guide for clarification.

UNDER PEER REVIEW