

1  
2  
3  
4  
5  
6  
7  
8  
9  
10

# SPAWNING TECHNIQUE OF TILAPIA (*Oreochromis niloticus*) AT THE FISH FRY CENTER IN CIBIRU, WEST JAVA

11 **ABSTRACT**

The demand for fish consumption continues to rise due to its high protein content, leading to an inadequate supply of fish stocks. To address this issue, the utilization of public water bodies for fish cultivation is seen as a potential solution. Cultivating fish in public waters can significantly boost fish production and maximize water utilization. One crucial stage in freshwater fish farming is freshwater fish hatchery development, aimed at meeting market demand for fish. Tilapia, a type of freshwater fish, holds great potential for cultivation due to its low production costs, high economic value, and adaptability to various aquatic conditions. The Cibiru Fish Fry Center in Bandung City plays a crucial role in tilapia fry production, with tilapia spawning being a significant endeavor. Tilapia is a freshwater fish species known for its elongated body shape, flattened sides, and blackish color. It exhibits a remarkable tolerance to different salinity levels and feeds on both animals and plants. Tilapia spawning at the Cibiru Fish Fry Center involves careful preparation of spawning ponds and the use of *kakaban* and water nets as substrates for egg attachment. Approximately 30,000 larvae are produced per spawning, with a hatching rate of 80%. Larvae management takes place in the same ponds as the spawning ponds and fertilizer application is used to promote the growth of natural food sources for the larvae. Feeding management is crucial for tilapia cultivation, with proper nutrition and feed quality influencing gonadal maturity and overall fish growth. The success of tilapia fry production at the center relies on various factors, including careful spawning preparation, optimal larval management, and effective feeding management.

12  
13 *Keywords: aquaculture, fish fry, freshwater, spawning, tilapia*

14  
15  
16 **1. INTRODUCTION**

17  
18 The demand for fish consumption has been steadily increasing due to its high animal protein content, both in seawater  
19 and freshwater fish. However, the growing consumer demand for fish is not being adequately met by the available fish  
20 stocks. In 2017, tilapia production reached 1.15 million tons, showing a 3.6% increase from the previous year's production  
21 of 1.14 million tons [1]. To address this issue, the utilization of public water bodies for fish cultivation is seen as a potential  
22 solution to enhance fishery production.

23 Cultivating fish in public waters can significantly boost fish production and maximize water utilization for increased  
24 productivity. Fish farming is a technological practice that encompasses various activities such as fish fry production,  
25 breeding, disease control, and ecosystem management. One crucial stage in freshwater fish farming is freshwater fish  
26 hatchery development, aimed at meeting the market demand for fish. The objective is to produce high-quality fish fries that  
27 fulfill the seven essential criteria: the right species, size, quality, quantity, timing, location, and price [2]. By adhering to  
28 these criteria, it is expected to meet the market's requirements and ensure sustainable fish production.

29 Tilapia, a type of freshwater fish, holds great potential for cultivation and is in high demand as a consumption  
30 commodity. Its popularity continues to rise due to its relatively low production costs and substantial economic value [3]  
31 (Arie, 2013). Tilapia is characterized by its elongated body shape, flattened sides, and blackish color. The name "Nile" was

officially assigned by the Indonesian Government through the Director General of Fisheries [4]. In Indonesia, fish cultivators have been successfully cultivating two types of tilapias: common tilapia and red tilapia. Besides being affordable, tilapia is rich in nutrients, exhibits rapid growth, displays resilience to environmental changes, has a high tolerance to various aquatic conditions, offers relatively large body sizes with good quality meat, is easy to breed, and boasts a high survival rate. Moreover, tilapia demonstrates adaptability to both freshwater and brackish water environments and shows resistance to diseases, as noted by [2].

The Cibiru Fish Fry Center, located in Bandung City, plays a crucial role as an agency focused on freshwater fish fry production and serves as a center for research and the application of fish farming technology. Among the various activities conducted at this center, tilapia (*Oreochromis niloticus*) spawning is a significant endeavor.

## 2. MATERIAL AND METHODS

The data collection method employed in this study encompasses both primary and secondary data. Primary data is obtained through recording observations, conducting informant interviews, and active participation. On the other hand, secondary data is gathered from books, reports, and other archival sources. The data analysis conducted for this study adopts a descriptive comparative approach, involving a comparison of field conditions with relevant literature. The parameters under observation include the spawning process, calculation of larval density in tilapia hatcheries, hatching rate (HR), survival rate (SR), and fecundity (F).

Hatching rate (HR) refers to the extent to which fish eggs successfully hatch. According to [5], determining the hatching rate of fish eggs can be accomplished using the following formula:

$$HR (\%) = \frac{P_t}{P_o} \times 100\%$$

Description:

HR: Degree of hatching of eggs

$P_t$ : The number of eggs that hatched

$P_o$ : The number of fertilized eggs

The survival rate (SR) represents the extent of fish survival from the initial stocking until the harvest stage. [5] suggests the following formula for calculating the fish survival rate:

$$SR (\%) = \frac{N_t}{N_o} \times 100\%$$

Information:

SR: Survival Rate of fish

$N_t$ : Number of live fish (frys harvested)

$N_o$ : Number of fish stocked (larvae)

Fecundity refers to the number of mature eggs released by the fish. According to [6], fecundity can be calculated using the following formula:

$$(F) = \frac{W}{w} \times n$$

Description:

F: Total number of eggs

W: Total egg weight (g)

w: Egg sample weight (g)

n: Total number of eggs counted during sampling (items)

## 3. RESULTS AND DISCUSSION

### 3.1 Biology and Morphology of Tilapia

Tilapia is a freshwater fish species commonly found in various aquatic environments, including rivers, lakes, reservoirs, swamps, rice fields, and irrigation canals. It exhibits a remarkable tolerance to different salinity levels. With an average weight ranging from 200 to 400 grams, tilapia is considered a large fish species. As an omnivorous fish, it feeds on both animals and plants, as noted by [7]. In its natural habitat, tilapia utilizes a diverse range of food sources, including phytoplankton, zooplankton, larvae, microorganisms present on the water bottom (molasses), aquatic plants like hydrilla, leftover food, as well as fruits and leaves that enter the water [2].



**Fig. 1. Nile tilapia**

According to [8] tilapia (*Oreochromis niloticus*) can be classified as follows:

Phylum: Chordata

Subphylum: Vertebrata

Class: Osteichthyes

Subclass: Acanthopterygii

Order: Percomorphi

Suborder: Percoidea

Family: Cichlidae

Genus: *Oreochromis*

Species: *Oreochromis niloticus*

Tilapia possesses an elongated and slender body shape with large, rough, and neatly arranged scales of the ctenoid type. The body color of tilapia is predominantly black with slight whitish hues. It features a distinctive lateral line that extends from the gill cover to the back of the dorsal fin. The eyes are prominent, large, and edged with white. There are five types of fins in tilapia: the dorsal fin, pectoral fin, ventral fin, anal fin, and caudal fin. The dorsal, pelvic, and anal fins are equipped with hard, sharp spines. The dorsal fin is characterized by its black coloration. Tilapia possess a pair of small pectoral and pelvic fins, a single dorsal fin, a lone anal fin, and a relatively elongated body shape. The caudal fin, on the other hand, is round and consists of a single lobe [7].

Tilapia is known for its euryhaline nature, enabling it to thrive in a wide range of salinity levels. It can tolerate water pH conditions ranging from 5 to 11; however, the ideal pH range for breeding and growth is 7 to 8 [8]. Tilapia can inhabit various freshwater habitats, including shallow waterways, ponds, rivers, and lakes. While tilapia can become problematic as an invasive species in warm water environments, it faces challenges in temperate climates due to its inability to survive in cold waters, typically below 21°C. Male tilapia generally exhibit higher salinity tolerance compared to females. Additionally, small tilapia tend to adapt more quickly to increased salinity levels compared to larger individuals [7].

### **3.2 Tilapia Spawning Activities**

#### **3.2.1 Preparation of Spawning Pools**

The process of tilapia spawning at the Cibiru Fish Fry Center involves the careful preparation of spawning ponds, and rearing ponds within a single pool. Rectangular-shaped concrete pools measuring 30x30x70m<sup>3</sup> are utilized for this purpose, with two ponds designated specifically for tilapia. To ensure optimal conditions, the first step is to clean the dry pond bottom, removing any snail shells and dried organisms. This meticulous cleaning process aims to eliminate potential sources of diseases. However, it is worth noting that cleaning the pond bottom is a practice rarely observed at the Cibiru Fish Fry Center.

Subsequently, the pool is filled with water by opening the inlet channel while closing the outlet channel. The water is then carefully filled to a height of 70 cm from the bottom of the pond. The next stage involves the installation of a water net and *kakaban*. The water net is secured with weights and firmly tied to poles situated at each end of the water net.



**Fig. 2. Spawning pond**

152  
153 In the prepared water net, the next step involves submerging the *kakaban*. The *kakaban* serves as a tool for securing  
154 eggs during the spawning process. Various materials are used by cultivators to create *kakaban*, such as coconut fiber,  
155 palm fiber, *paranets*, or even water nets. At the Cibiru Fish Fry Center, rectangular palm fiber is used to construct the  
156 *kakaban*. It is clamped with two bamboo blades and weighted with stones. A total of 10 *kakabans* and 2 water nets are  
157 utilized. The *kakaban* and water net serve the crucial function of providing a suitable environment for tilapia eggs to attach  
158 and develop.  
159  
160  
161  
162



163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178 **Fig. 3. Water Net instalation**  
179



180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192 **Fig. 4. Substrate for egg (*kakaban*) instalation**  
193  
194

### 195 **3.2.2 Parent Selection**

196 At the Cibiru Fish Fry Center, meticulous parent selection takes place in the broodstock rearing pond. The selection  
197 process is primarily based on sex determination, which is achieved through the stripping method applied to the abdomen.  
198 When male tilapia are stripped, they release a thick milky white sperm, while female tilapia exhibit abdominal swelling  
199 towards the urogenital region and emit a yellow liquid when stripped.

200 Parent tilapia at the Cibiru Fish Fry Center is deemed ready for spawning when they reach two years of gonad maturity,  
201 ensuring that their reproductive organs are fully developed. The maturation process of the main gonads is influenced by  
202 both internal and external factors. Internal factors encompass the fish species and hormones, while external factors  
203 encompass temperature, food availability, stocking density, light intensity, and more.

204 Among the external factors, particular attention is given to the influence of feed and the environment on the maturation  
205 of parent gonads [9]. Reproduction in fish is intricately regulated by environmental factors that impact the function of the  
206 endocrine glands, which are responsible for producing reproductive hormones crucial for gonad development,  
207 gametogenesis, and the reproductive cycle [10]. The readiness of the parents for spawning is indicated by the release of  
208 white liquid (sperm) upon the gentle massage of the lower abdomen towards the anus [11]. This process is influenced by  
209 various environmental stimuli, including temperature, light, and weather, which are received by receptors and  
210 subsequently transmitted to the nervous system. The hypothalamus then releases hormones that stimulate the pituitary  
211 gland, thereby controlling the development and maturity of the gonads for successful spawning [11].  
212



Fig. 5. Preparation of tilapia parent sorting

Characteristics of mature female tilapia include slow movements, an enlarged or distended abdomen towards the back, a soft texture when touched, a slight protrusion or swelling of the anal canal, and the release of a reddish-yellow liquid when gently massaged towards the anus. On the other hand, male parents exhibit agile movements, slim bands, and emit white sperm fluid when massaged toward the anus [12]. In terms of weight ratio, the broodstock used consists of 6 kg of males and 2 kg of females, maintaining a ratio of 3:1. This ratio is chosen to ensure maximum fertilization of the female's eggs by male tilapia sperm, thus optimizing the reproductive success of the breeding process.

### 3.2.3 Spawning

The reproductive activity of aquatic animals varies depending on environmental conditions [10]. According to [13], fish spawning techniques in fish farming can be categorized into three methods:

- 1) Natural fish spawning: This method involves fish spawning without human intervention, occurring naturally without hormone stimulation.
- 2) Semi-artificial spawning: In this method, hormone stimulation is provided to accelerate gonadal maturity, but the actual ovulation process occurs naturally in ponds.
- 3) Artificial fish spawning: This technique involves hormone stimulation to accelerate gonadal maturity, and the ovulation process is artificially induced through stripping or sequencing techniques.

Each method offers distinct advantages and can be chosen based on specific breeding goals and environmental factors.



Fig. 6. Tilapia spawning ponds

Tilapia spawning at the Cibiru Fish Fry Center is conducted naturally, without human intervention or hormone stimulation. Male and female parents are placed together in a pond, utilizing *kakaban* and water net as substrates for the eggs to attach to. According to [14], the optimal time to release the broodstock is in the morning or evening, when the water temperature is relatively low. At the Cibiru Fish Fry Center, tilapia spawning takes place at night. The female parent releases eggs onto the *kakaban* and water net before midnight, followed by the male parent secreting white sperm. Fertilization calculations are carried out 6 hours after the spawning process. Based on observations, fertilized eggs appear transparent with clear contents, while unfertilized eggs appear pale, in line with [15] statement that fertilized eggs display a brown color, whereas unfertilized eggs appear opaque and pale.

After spawning, further observations are made in the spawning ponds, where tilapia eggs are observed adhering to the *kakaban* and water net. Hatched eggs develop into larvae, while unhatched eggs remain white, indicating their demise. Factors such as imperfect fertilization and eggs sticking together or overlapping in the nets, causing disrupted oxygen circulation, can contribute to egg mortality [16]. The success of hatching tilapia eggs is influenced by the maturity of brood fish gonads and water quality.

274 Spawning is conducted three times a year, resulting in approximately 30,000 larvae per spawning. Each female tilapia  
275 weighing 450 grams exhibits high fecundity, producing around 550 eggs per head. This is supported by [17], which states  
276 that tilapia broodstock weighing 500 grams has a fecundity of 200 eggs per 100 grams. Research conducted by  
277 Widyastuti et al. (2008) on selected and non-selected tilapia using artificial spawning revealed a fecundity range of 620–  
278 1,175 eggs per kilogram. Approximately 350 larvae successfully hatch, resulting in an 80% hatching rate (HR). According  
279 to [18], the normal hatching percentage for fish ranges from 50% to 80%.

280 After hatching, the water net and *kakaban* are removed and cleaned. After one month, the parent tilapia is transferred  
281 from the spawning pond to the brood-rearing pond. The advantage of the tilapia spawning technique at the Cibiru Fish Fry  
282 Center is the integration of spawning and larval rearing in a single pond, eliminating the need for the "*anco*" (life net)  
283 system used in other fish fry centers. This approach offers space efficiency by utilizing a limited hall area, optimizing time  
284 usage, and requiring fewer human resources.

### 285 3.2.4 Larvae Management

287 Tilapia larvae handling takes place in the same ponds as the spawning ponds at the Cibiru Fish Fry Center. The  
288 hatching process of tilapia eggs typically lasts for approximately 48 hours. Hatched eggs develop into larvae, while  
289 unhatched eggs remain white, indicating that they did not survive, as mentioned by [12].

290 Fish larvae represent a critical phase in fish farming due to their vulnerability to environmental changes and limited  
291 resistance. Once hatched, tilapia larvae will attach themselves to the *kakaban*. Therefore, the *kakaban* is removed 2–3  
292 days after the eggs have hatched.

293 At 2–3 days old, the larvae still possess food reserves in the form of egg yolk sacs, which can sustain them for around  
294 3–4 days. Another essential aspect of larval management is the application of fertilizer. The Cibiru Fish Fry Center utilizes  
295 a combination of goat manure and rice husk as fertilizers. This mixture is prepared by creating small holes on each  
296 surface of the sacks containing eight sacks of fertilizer, each weighing 50 kg. The purpose of this fertilizer application is to  
297 promote the growth of natural food sources required by the larvae once they have depleted their yolk reserves.

### 298 3.3 Feeding Management

299 The quality and quantity of feed play a crucial role in hatchery processes, broodstock maintenance, and fish growth.  
300 Effective feeding management significantly impacts the growth and overall condition of fish, especially in achieving  
301 optimal gonadal maturity. One approach to ensuring proper nutrition is by estimating protein requirements and protein-  
302 energy ratios, as suggested by [19]. This aligns with the notion emphasized by [9], highlighting the significant influence of  
303 feed and environmental factors on broodstock gonadal maturity. Providing high-quality feed in sufficient quantities is  
304 essential for enhancing the quality of the broodstock. It is important to note that feed quality has a significant impact on  
305 both male and female gonadal maturity, making the selection of appropriate feed crucial to the process.

306 The availability of adequate feed in terms of quality and quantity greatly influences tilapia cultivation, as emphasized by  
307 [20]. Feed accounts for approximately 60 ± 70% of the production costs in tilapia aquaculture, underscoring the need for  
308 effective and efficient feed management. The good feed should fulfill the nutritional requirements of the fish. Enhanced  
309 tilapia growth can be achieved through improved utilization of feed protein, leading to greater deposition of body protein  
310 [21]. The absorbability of nutrients by the fish's body is influenced by factors such as feed composition and quantity [22].

311 At the Cibiru Fish Fry Center, artificial feed (pellets) is provided to the parent fish and tilapia larvae twice a day, in the  
312 morning at 08:30 AM and in the afternoon at 03:00 PM. [23] suggests that feeding tilapia broodstock should be carried out  
313 in the morning and evening, with a feeding rate (FR) of 3%. At the center, feeding is done along the edges of the pond,  
314 establishing a habit for the fish to gather near the edge, which facilitates harvesting later on.

315 For the maintenance of tilapia broodstock, sinking pellet-type artificial feed is used, specifically the "Cargill" brand. The  
316 feed provided to tilapia larvae is the "Feng-Li Standard" brand in powder form. The availability of feed at the Cibiru Fish  
317 Fry Center is determined by the Bandung Food and Agriculture Service. The "Feng-Li" feed has a refined protein content  
318 of 40%. Cargill fish feed contains a minimum of 26% crude protein, 5% fat, 4% maximum crude fiber, 10% maximum ash,  
319 12% maximum moisture content, and 100 ppm vitamin C.

320 It is worth noting that the feed provided at the Cibiru Fish Fry Center is not specifically formulated for tilapia species.  
321 The catfish feed used is designed for sinking and tailored to catfish needs. On the other hand, the "Feng-Li Standard"  
322 feed is formulated for intensive tiger prawn cultivation. The "Cargill" brand feed, with a protein content of 26%, does not  
323 fully meet the protein requirements of tilapia broodstock. The digestibility values, which indicate the percentage of  
324 nutrients absorbed by the fish's digestive tract, closely relate to feed conversion and efficiency.

325 Nutrient absorption by the fish's body is influenced by various factors, including feed quality and quantity consumed.  
326 Therefore, the suitability of the feed type can directly impact tilapia growth. The utilization of nutrients by tilapia affects the  
327 supply of protein and non-protein energy in the body. Increased energy availability enhances the fish's ability to convert  
328 and store energy in the form of protein and fat.

329 Feeding should be adjusted to the fish's needs, employing a gradual feeding system until the fish no longer consume  
330 the feed provided. This approach reduces the amount of unconsumed feed, minimizing contamination of the rearing  
331 environment. Nutrient intake from additional or artificial feed significantly improves fish growth due to the more complete  
332 and sufficient nutrients entering the fish's body.

333 Ad-libitum feeding was implemented, which involved providing feed in small increments until the fish exhibited signs of  
334

335 satiety and no longer responded to the feed. This feeding method aimed to minimize the accumulation of leftover feed at  
336 the bottom of the rearing tank, as such remnants can lead to the production of ammonia. Excessive ammonia levels can  
337 negatively impact water quality and often lead to fish mortality. For a visual representation of the feeding process at the  
338 Cibiru Fish Fry Center, please refer to Figure 7.



340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357 **Fig. 7. Feeding stage**

#### 358 359 360 **4. CONCLUSION**

361 Through a comprehensive analysis of observations, literature studies, and practical experience, it can be inferred that tilapia  
362 spawning at the Cibiru Fish Fry Center is conducted naturally, resulting in remarkable productivity. The spawning process occurs  
363 three times a year, yielding a substantial number of larvae in each cycle. Notably, the Cibiru Fish Fry Center stands out for its  
364 advantageous tilapia spawning technique, wherein both spawning and larval rearing take place within a single pond, ensuring  
365 heightened efficiency throughout the entire process.

#### 366 367 **COMPETING INTERESTS**

368  
369 "Authors have declared that no competing interests exist."  
370  
371

#### 372 373 **AUTHORS' CONTRIBUTIONS**

374 Author 1' designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the  
375 manuscript, and managed the analyses of the study. 'Author 2' managed the literature searches and writing. All authors  
376 read and approved the final manuscript."  
377  
378

#### 379 380 **REFERENCES**

- 381 1. Direktorat Jenderal Perikanan Budidaya. Statistik Perikanan Budidaya. Jakarta: Direktorat Jenderal Perikanan Budidaya,  
382 Kementerian Kelautan dan Perikanan; 2017.
- 383 2. Kementerian Kelautan dan Perikanan. Peluang Usaha dan Investasi Nila. Jakarta: Direktorat Jenderal Penguatan Daya Saing  
384 Produk Kelautan dan Perikanan; 2019.
- 385 3. Arie, U dan D. Dejee. Panduan Lengkap Benih Ikan Konsumsi. Jakarta: Penebar Swadaya; 2013
- 386 4. Ghufuran M, Kordi H. Panduan Lengkap Memelihara Ikan Air Tawar di Kolam Terpal. Yogyakarta: Penerbit Andi; 2010.
- 387 5. Kurniaji A, Nuryati S, Murtini S, Alimuddin. Maternal immunity response and larval growth of anty cyhv-3 dna vaccinated  
388 common carp (*Cyprinus carpio*) at different pre-spawning time. Pak. J. Biotechnol. 2018;15(3): 689-698.
- 389 6. Ansyari S, Slamet S. The gonad maturity and fecundity of toman fish (*Channami cropeltus*) during rainyseason to dry season  
390 periode in Rawa Danau Panggang waters, Hulu Sungai Utara Regency, South Kalimantan, Indonesia. IOP Conference  
391 Series: Earth and Environmental Science. 2019.
- 392 7. Amri, Khairuman. Budidaya Ikan Nila Secara Intensif. Jakarta: Agromedia; 2003.
- 393 8. Kordi MGH. Budidaya Ikan Nila di Kolam Terpal. Yogyakarta: Lili Publisher; 2013.

- 394 9. Sitiady S. Pengaruh Pemberian Vitamin E dengan Dosis yang Berbeda Terhadap Kematangan Gonad Ikan Selais (*Ompok*  
395 *hypophthalmus*). Riau: Fakultas Perikanan dan Ilmu Kelautan Universitas Riau. 2008. (*bachelor thesis*)
- 396 10. Fujaya Y. Fisiologi Ikan Dasar Pengembangan Teknik Perikanan. Cetakan pertama. Jakarta: Rineka Putra; 2004.
- 397 11. Mashaii N, Rajabipour F, Jafari M, Mohammadi M, Sarsangi H, Zadeh HH, Matinfar A. Some Spawning Performance  
398 Parameters of Cultured Nile Tilapia *Oreochromis niloticus*, under Brackish Water. *Aquatic Science and Technology*. 2019;  
399 7(1):8-22.
- 400 12. Saputra SD. Aplikasi Sistem Resirkulasi Air Terkendali (SRAT) pada Budidaya Ikan Mas (*Cyprinus carpio*). Bogor: Fakultas  
401 Teknologi Pertanian. Institut Pertanian Bogor; 2011. (*bachelor thesis*)
- 402 13. Susanto H. *Teknik Kawin Suntik Ikan Ekonomis*. Jakarta: Penebar Swadaya; 2011.
- 403 14. Ismail A, Khumaidi. Teknik Pembenihan Ikan Mas (*Cyprinus carpio* L.) di Balai Benih Ikan Tenggara Bondowoso. *Jurnal*  
404 *Ilmu Perikanan*. 2016;7(1): 32.
- 405 15. Suryani SAMP, Abdulah FC, Faudzi NM, Sadguna DN, Arya IW, Edi DGS, Pratama GAS. Improving the Quality of Tilapia  
406 Fish Seeds (*Oreochromis Niloticus*) with Selection Methods in Mina Ayu Group of Tabanan Regency Bali Province. *Asian*  
407 *Journal of Applied Research for Community Development and Empowerment*. 2022; 6(1).
- 408 16. Setyono B. Pengaruh Perbedaan Konsentrasi Bahan pada Pengencer Sperma Ikan “Skim Kuning Telur” terhadap Laju  
409 Fertilisasi, Laju Penetasan dan Sintasan Ikan Mas (*Cyprinus carpio* L.) Malang: Fakultas Pertanian dan Peternakan.  
410 Universitas Muhammadiyah Malang; 2009. (*bachelor thesis*)
- 411 17. SNI. Produksi Benih Ikan Nila Hitam (*Oreochromis niloticus*) Kelas Benih Sebar; 2009
- 412 18. Putri FP, Dewi NN. Growth monitoring of koi fish (*Cyprinus carpio*) in natural hatchery techniques in Umbulan, Pasuruan, East  
413 Java. *IOP Conference Series: Earth and Environmental Science*. 2019.
- 414 19. Pramono TB, Sanjayasari D, Soedibya PHT. Optimasi Pakan dengan Level Protein dan Energi Protein untuk Pertumbuhan  
415 Calon Induk Ikan Senggaringan (*Mystus nigriceps*). *Terubuk*. 2007; 2:152-157.
- 416 20. Handajani. Penyusun Pakan Ikan terhadap Pertumbuhan dan Daya Cerna Ikan Nila GIFT (*Oreochromis* sp.). Malang Fapetrik  
417 Universitas Muhammadiyah Malang. Malang; 2004. (*grant report*)
- 418 21. Phipilai S, Ronald D Wardlaw FB. Relation of Two In Vitro Assays in Protein Efficiency Ratio Determination on Selected  
419 Agricultural by  $\pm$  Products. *Journal Science Technology*. 2006;26: 81-87.
- 420 22. Marzuqi M, Anjusary DN. Kecernaan Nutrien Pakan dengan Kadar Protein dan Lemak Berbeda pada Juvenil Ikan Kerapu  
421 Pasir (*Epinephelus corallicola*). *Jurnal Ilmu dan Teknologi Kelautan Tropis*; 2013; 5(2):311-323.
- 422 23. Nurezka K, Junianto. Feeding Management for Tilapia (*Oreochromis niloticus*) Seeds and Parent At Cibiru Fish Seed Center,  
423 Bandung City, West Java. *IJARIE*. 2022; 8(6).
- 424