

Original Research Article

NUTRIENT MANAGEMENT PRACTICES ON GROWTH AND YIELD OF FINGER MILLET INFLUENCED BY DIFFERENT FACTORS

ABSTRACT

A field experiment was conducted during Rabi season of 2022 in at South farm of Karunya Institute of Technology and Sciences, Coimbatore. This study was conducted to assess the growth and yield of finger millet using nutrient management practices. The treatment consists of seven different parameters with control. Results revealed that growth parameters, yield attributes, and yield of finger millet were significantly influenced by different treatments of nutrient management. In finger millet, plant height at 90 DAS, LAI at 60 DAS, and dry matter accumulation per m² at harvest were maximum with 125% seed rate and 100% N through vermicompost. Yield attributes namely, effective tillers per m², number of fingers per ear, ear weight, test weight, number of grains per ear, grain yield and straw yield of finger millet were also recorded maximum in the same treatment.

Keywords: Nutrient management; Vermicompost; growth attributes; Yield attributes; Finger millet.

INTRODUCTION

Finger millet, commonly known as ragi, is grown extensively in various regions of India. It is used as a staple food that supplies a major portion of calories and protein for people of low-income groups. Finger millet is generally taken in uplands in Jharkhand where they perform poorly due to low soil fertility as well as poor plant stand under direct sowing conditions. In recent decades, emphasis has been shifted from individual crop to cropping systems because responses in component crops are influenced by the nutrient application to preceding crops by leaving a substantial effect on the succeeding crop as carry-over benefit. (1)

In recent decades, emphasis has been shifted from individual crops to cropping systems because responses in component crops are influenced by the nutrient application to preceding crops by leaving a substantial effect on the succeeding crop as carry-over benefit. Also, taking two crops in a sequence, or intensive cropping in place of mono-cropping in uplands like the inclusion of pulses in crop sequence is agronomically very significant. It has been suggested that there is no need to apply fertilizers if moderate nutrient-requiring crops like pea succeeds. A deleterious effect of chemical

fertilizers in agriculture has led to adopt organic crop production as an alternative method which also maintains soil health and improves the overall ecological balance of the production system. Thus, adopting combination of proper plant population and organic nutrient management can lead in to better grain production. Information on seed rate in finger millet under organic nutrient management and its residual effect on the second crop in the sequence is very meager. (2)

Therefore in this study, the use of locally available agro-inputs in agriculture by avoiding or minimizing the use of synthetic agrochemicals appears to be one of the probable possible options to sustain the agricultural productivity. Various vermicompost nutrient sources are available which contain good amounts of major significant plant nutrients to produce comparable yields. At the same time, the food habits of the consumers are changing rapidly. Especially in the developed countries people have become more health conscious. Hence the demand for organic food products is on the rise. (3)

MATERIALS AND METHODS

Experimental site

During the Rabi season of 2022, the experiment was conducted in the South Farm, School of Agricultural Sciences, Division of Agronomy, Karunya Institute of Technology and Sciences, Coimbatore. 10.9362° N latitude and 76.744° E longitude are the farm's coordinates. The region experiences a typical warm, muggy environment. (4)

Soil characteristics

Prior to the experiment, random soil samples were taken from the experimental field at 5 separate places, ranging in depth from 0 to 15 cm. All of the soil samples were combined to create a typical homogenous composite sample, which was then examined to ascertain the mechanical and physicochemical characteristics of the soil. (5)

Experimental design

The experiment was laid out in randomised block design with ten treatments and replicated three times. The treatments were allotted at random to plots within each replication.

List 1: Various treatments and their doses used for the study.

S No.	Treatment	Dose(kg/ha)	Legends
1.	100% (recommended dose of fertilizer)	40:20:20(N:P:K)	T ₁

2.	75% of RDF +25% of FYM/ha (2 tonn)	30:15:15(N:P:K) 2 t FYM	T ₂
3,	50% of RDF +50% of FYM/ha(4tonn)	20:10:10(N:P:K) 4 t FYM	T ₃
4.	25% of RDF +75% of FYM/ha (6 tonn)	10:05:05(N:P:K) 6 t FYM	T ₄
5.	75% of RDF+ 25% of vermicompost/ha (0.5 tonn)	30:15:15(N:P:K) 0.5 t Vermicompost	T ₅
6.	50% of RDF+ 50% of vermicompost/ha (1tonn)	20:10:10(N:P:K) 1 t Vermicompost	T ₆
7.	25% of RDF+ 75% of vermicompost/ha (1.5 tonn)	10:05:05(N:P:K) 1.5 t Vermicompost	T ₇
8.	Control	00-00-00	T ₈

Growth attributes

Five plants are selected at random from the net plot area of each treatment and tagged. The following parameters are recorded in those tagged plants at different days.

3.5.1.1. Plant height

Plant height was measured at 30, 60 DAS and at harvest stage in the five tagged plants from the ground level to tip of the plant and the value is expressed in cm. (6)

3.5.1.2. Number of tillers plant⁻¹

Number of tillers were manually counted at 30, 60 DAS and at harvest stage in the five tagged plants. (7)

3.5.1.3. Dry matter production

Five plants selected at random at all stages from each plot outside the net plot but within the border rows were cut close to the ground level and the samples were collected. These samples were shades dried and then oven dried at 80°C for 72 hours. The dry matter production was computed per unit area and expressed in kg ha⁻¹. (8)

Yield attributes of crop

3.5.3.1. Number of productive tillers hill⁻¹

Five plants in each net plot were selected at random and the number of productive tillers per plant were counted and averaged. (9)

3.5.3.2. Number of fingers earhead⁻¹

Five plants in each plot were selected random and number of fingers per ear head were counted and averaged. (10)

3.5.3.3. Number of tillers m⁻²

The number of tillers were counted from randomly tagged plants in net plot area and averaged to compute number of tillers m⁻². (11)

3.5.3.4. Thousand grain weight

Three composite samples of each 1000 grains were drawn from the net plot yield of each treatment and weight of these samples was recorded, averaged and expressed as 1000 grain weight in grams. (12)

3.5.3.5. Grain yield

The matured ear heads are collected from the net plot by harvesting followed by threshing. Finally the grains were collected and later it was cleaned, sun dried, weighed and grain yield was calculated and expressed in kg ha⁻¹. (13)

3.5.3.6. Straw yield

After harvest of the ear head, the whole plants inside the net plot are cut above the ground level. Later it was sun dried for three days and weighed. The straw yield was calculated and expressed in kg ha⁻¹. (14)

RESULTS

Table 1 Effect of integrated nutrient management on plant height (cm) at various growth stages of Finger Millet

Treatments		Plant height (cm)		
		60 DAT	90 DAT	At Harvest
T ₁	100% (recommended dose of fertilizer)	36.19	55.32	74.43
T ₂	75% of RDF +25% of FYM/ha (2 tonn)	36.03	54.42	70.17
T ₃	50% of RDF +50% of FYM/ha(4tonn)	37.12	56.74	76.28
T ₄	25% of RDF +75% of FYM/ha (6 tonn)	37.01	56.03	71.32
T ₅	75% of RDF+ 25% of vermicompost/ha (0.5 tonn)	37.43	57.98	72.19
T ₆	50% of RDF+ 50% of vermicompost/ha (1tonn)	38.17	61.02	78.34
T ₇	25% of RDF+ 75% of vermicompost/ha (1.5 tonn)	36.16	58.12	73.38
T ₈	Control	35.18	53.43	70.55
SEm±		0.33	0.85	1.01
C.D. (5%)		NS	2.32	2.78

The data on plant height (cm) at different stages of growth are given in Table no.1 and the analysis of variance in appendix 1.

At 60 DAT, T₆ (38.17) recorded the highest plant height and there was no significant difference between treatments. The lowest plant height (35.18) was observed in T₈.

At 90 DAT, the plant height was significantly influenced by the treatments, maximum plant height (61.02) was observed in T₆ and it was at par with T₄, T₅, T₇ and T₈ and was significantly superior over T₁, T₂, T₃ and T₈. The lowest plant height (53.43) was observed in T₈ and was at par with T₁, T₃, T₄ and T₇.

At Harvest, also maximum plant height was observed in T₆ (78.34) and it was at par with T₄, T₅, T₇ and T₈ and was significantly superior over T₁, T₂, T₃ and T₇. The lowest plant height (70.55) was observed in T₈ and was at par with T₁, T₃, T₄ and T₅. In general T₆ produced taller plants than other treatments at all times. (15)

Table 2 Effect of integrated nutrient management on numbers of tillers (m⁻²) at various growth stages of Finger Millet

Treatments		Plant height (cm)		
		60 DAT	90 DAT	At Harvest
T ₁	100% (recommended dose of fertilizer)	34.23	56.25	68.73
T ₂	75% of RDF + 25% of FYM/ha (2 tonn)	30.11	52.39	64.38
T ₃	50% of RDF + 50% of FYM/ha(4tonn)	35.28	57.63	69.29
T ₄	25% of RDF + 75% of FYM/ha (6 tonn)	31.82	53.28	65.34
T ₅	75% of RDF + 25% of vermicompost/ha (0.5 tonn)	32.38	54.32	66.18
T ₆	50% of RDF + 50% of vermicompost/ha (1tonn)	36.46	58.98	70.87
T ₇	25% of RDF + 75% of vermicompost/ha (1.5 tonn)	33.28	55.43	67.22
T ₈	Control	29.32	50.32	63.23
SEm ±		0.86	1.00	0.92
C.D. (5%)		NS	2.12	1.89

The data on numbers of tillers at different stages of growth are given in Table no.2 and the analysis of variance in appendix 1.

At 60 DAT, T₆ (36.46) recorded the highest numbers of tillers and there was no significant difference between treatments. The lowest numbers of tillers (29.32) was observed in T₈.

At 90 DAT, the numbers of tillers was significantly influenced by the treatments, maximum numbers of tillers (58.98) was observed in T₆ and it was at par with T₄, T₅, T₇ and T₈ and was

significantly superior over T₁, T₂, T₃ and T₄. The lowest numbers of tillers (50.32) was observed in T₈ and was at par with T₁, T₃, T₄ and T₅.

At Harvest, also maximum numbers of tillers was observed in T₆ (70.87) and it was at par with T₄, T₅, T₇ and T₈ and was significantly superior over T₁, T₂, T₃ and T₇. The lowest numbers of tillers (63.23) was observed in T₈ and was at par with T₁, T₃, T₄ and T₅. In general T₆ produced more numbers of tillers then other treatments at all times. (16)

Table 3 Effect of integrated nutrient management on Days to first flowering of Finger Millet

Treatments		Days to first flowering
T ₁	100% (recommended dose of fertilizer)	43.22
T ₂	75% of RDF + 25% of FYM/ha (2 tonn)	40.21
T ₃	50% of RDF + 50% of FYM/ha(4tonn)	44.32
T ₄	25% of RDF + 75% of FYM/ha (6 tonn)	41.01
T ₅	75% of RDF + 25% of vermicompost/ha (0.5 tonn)	41.52
T ₆	50% of RDF + 50% of vermicompost/ha (1tonn)	45.55
T ₇	25% of RDF + 75% of vermicompost/ha (1.5 tonn)	42.48
T ₈	Control	39.21
S.E. (m±)		0.75
CD (p=0.05)		1.73

The data on days of first flowering at different stages of growth are given in Table no.3 and the analysis of variance in appendix 1.

The highest Days to first flowering was occurred in T₆ (45.55) and there was no significant difference between treatments. The lowest Days to first flowering was occurred in (39.21) was observed in T₈.

Table 4 Effect of integrated nutrient management on Days to 50% flowering of Finger Millet

Treatments		Days to 50% flowering
T ₁	100% (recommended dose of fertilizer)	73.27
T ₂	75% of RDF + 25% of FYM/ha (2 tonn)	70.21
T ₃	50% of RDF + 50% of FYM/ha(4tonn)	74.32

T ₄	25% of RDF + 75% of FYM/ha (6 tonn)	71.02
T ₅	75% of RDF + 25% of vermicompost/ha (0.5 tonn)	71.27
T ₆	50% of RDF + 50% of vermicompost/ha (1tonn)	75.45
T ₇	25% of RDF + 75% of vermicompost/ha (1.5 tonn)	72.33
T ₈	Control	69.46
S.E. (m±)		0.73
CD (p=0.05)		2.12

The data on Days to 50% flowering at different stages of growth are given in Table no.3 and the analysis of variance in appendix 1.

The highest Days to 50% flowering was occurred in T₆ (75.45) and there was no significant difference between treatments. The lowest Days to 50% flowering was occurred in (69.46) was observed in T₈. (17)

Table 5 Effect of integrated nutrient management on Total Dry matter production (kg/ha) at various growth stages of Finger Millet

Treatments		Total Dry matter production (kg/ha)		
		60 DAT	90 DAT	At Harvest
T ₁	100% (recommended dose of fertilizer)	39.28	61.98	78.39
T ₂	75% of RDF +25% of FYM/ha (2 tonn)	34.98	57.47	74.32
T ₃	50% of RDF +50% of FYM/ha(4tonn)	40.26	62.87	79.03
T ₄	25% of RDF +75% of FYM/ha (6 tonn)	36.52	58.72	75.38
T ₅	75% of RDF+ 25% of vermicompost/ha (0.5 tonn)	37.33	59.83	76.63
T ₆	50% of RDF+ 50% of vermicompost/ha (1tonn)	41.25	63.23	81.26
T ₇	25% of RDF+ 75% of vermicompost/ha (1.5 tonn)	38.22	60.12	77.38
T ₈	Control	32.11	55.23	72.32
SEm±		1.05	0.97	0.99
C.D. (5%)		NS	2.12	2.56

The data on total Dry matter production at different stages of growth are given in Table no.5 and the analysis of variance in appendix 1.

At 60 DAT, T₆ (41.25) recorded the highest total Dry matter production and there was no significant difference between treatments. The lowest total Dry matter production (32.11) was observed in T₈.

At 90 DAT, the total Dry matter production was significantly influenced by the treatments, maximum total Dry matter production (63.23) was observed in T₆ and it was at par with T₄, T₅, T₇ and T₈ and was significantly superior over T₁, T₂, T₃ and T₅. The lowest total Dry matter production (55.23) was observed in T₈ and was at par with T₁, T₃, T₄ and T₅.

At Harvest, also maximum total Dry matter production was observed in T₆ (81.26) and it was at par with T₄, T₅, T₇ and T₈ and was significantly superior over T₁, T₂, T₃ and T₅. The lowest total Dry matter production (72.32) was observed in T₈ and was at par with T₁, T₃, T₄ and T₅. In general T₆ produced more Dry matter production then other treatments at all times. (18)

YIELD PARAMETERS

Table 6 Effect of integrated nutrient management on Yield Parameters (No. of fingers/head, No. of productive tillers/plant, No. of ear head/plant) of Finger Millet

Treatments		No. of fingers/head	No. of productive tillers/plant	No. of ear head/plant
T ₁	100% (recommended dose of fertilizer)	10.42	117.88	8.75
T ₂	75% of RDF +25% of FYM/ha (2 tonn)	6.43	112.22	4.19
T ₃	50% of RDF +50% of FYM/ha (4tonn)	11.21	118.28	9.23
T ₄	25% of RDF +75% of FYM/ha (6 tonn)	7.62	113.27	5.53
T ₅	75% of RDF+ 25% of vermicompost/ha (0.5 tonn)	8.27	115.12	6.21
T ₆	50% of RDF+ 50% of vermicompost/ha (1tonn)	12.56	120.27	10.27
T ₇	25% of RDF+ 75% of vermicompost/ha (1.5 tonn)	9.33	116.23	7.55
T ₈	Control	5.43	110.43	2.77
SEm±		0.85	1.18	0.92
C.D. (5%)		NS	3.23	1.58

The various yield parameters majorly contributing in the treatments can be determined, No. of fingers/head in yield parameters contributing T₆ (12.56) recorded the highest and there was no significant difference between treatments. The lowest (5.43) was observed in T₈.

No. of productive tillers/plant in yield parameters contributing T₆ (120.27) recorded the highest and there was no significant difference between treatments. The lowest (110.43) was observed in T₈.

No. of ear head/plant in yield parameters contributing T₆ (10.27) recorded the highest and there was no significant difference between treatments. The lowest (7.55) was observed in T₈.

Table 7 Effect of integrated nutrient management on Yield Parameters (100 grain weight (g), Grain yield (kg/ha), Straw yield (kg/ha)) of Finger Millet

Treatments		1000 grain weight (g)	Grain yield (kg/ha)	Straw yield (kg/ha)
T ₁	100% (recommended dose of fertilizer)	3.64	2478	5532
T ₂	75% of RDF +25% of FYM/ha (2 tonn)	3.28	2037	5122
T ₃	50% of RDF +50% of FYM/ha(4tonn)	3.75	2543	5635
T ₄	25% of RDF +75% of FYM/ha (6 tonn)	3.31	2127	5213
T ₅	75% of RDF+ 25% of vermicompost/ha (0.5 tonn)	3.44	2283	5321
T ₆	50% of RDF+ 50% of vermicompost/ha (1tonn)	3.87	2650	5753
T ₇	25% of RDF+ 75% of vermicompost/ha (1.5 tonn)	3.52	2387	5483
T ₈	Control	3.16	1990	5023
SEm±		0.08	86.10	90.93
CD (p=0.05)		NS	112.32	125.27

The various yield parameters majorly contributing in the treatments can be determined, 1000 grain weight (g) in yield parameters contributing T₆ (3.87) recorded the highest and there was no significant difference between treatments. The lowest (3.16) was observed in T₈.

Grain yield (kg/ha) in yield parameters contributing T₆(2650) recorded the highest and there was no significant difference between treatments. The lowest (1990) was observed in T₈.

Straw yield (kg/ha) in yield parameters contributing T₆(5753) recorded the highest and there was no significant difference between treatments. The lowest (5023) was observed in T₈. (19)

Conclusion

Among the nutrient management practices 50% of recommended dose of fertilizer with 50% of vermicompost was best in yield and yield attributing parameter. As per the uptake of nutrients and economic point of view, the application of 50% RDF with 50% vermicompost was superior as compare to all treatments. The present study concluding that 50 % NPK through inorganic and 50 % through vermicompost may be suggest to take good yield, economical and good soil health advantages of finger millet crop under rainfed condition.

REFERENCES

1. Agriculture statistics at a glance. Government of India, Ministry of Agriculture and farmer's welfare, Department of Agriculture, Directorate of Economics and Statistics, 2017.

2. Ananda MR, Sharanappa, Kalyana Murthy KN. Growth, yield and quality of groundnut as influenced by organic nutrient management in groundnut (*Arachis hypogaea* L.) finger millet (*Eleusine coracana* L.) cropping system. Mysore Journal of Agricultural Sciences. 2017 51(2):385-391.
3. Basavaraju TB, Purushotham S. Integrated nutrient management in rainfed ragi (*Eleusine coracana* L. Gaertn.). Mysore Journal of Agricultural Sciences. 2009; 43(2):366-368.
4. Basavaraj Naik T, Kumar Naik AH, Suresh Naik KP. Nutrient management practices for organic cultivation of finger millet (*Eleusine coracana* L.) under southern transitional zone of Karnataka, India. International Journal of Current Microbiological Applied Sciences. 2017; 6(11):3371-3376.
5. Giraddi RS. Effect of stocking rate of *Eudrilus eugeniae* on vermicompost production. Karnataka Journal of Agricultural sciences. 2008; 21(1):49-51.
6. Giribabu B, Luther MM, Chandra Sekhar K, Sankara Rao V. Effect of nutrient management system on productivity of finger millet (*Eleusine coracana* L. Gaertn) cultivars under sandy soils. The Andhra Agricultural Journal. 2010; 57(1):4-6
7. Narayan Hebbal, Ramachandrappa BK, Thimmegouda MN. Yield and economics of finger millet with establishment methods under different Planting geometry and nutrient Source. Indian Journal of Dryland Research and Development. 2018; 33(1):54-58.
8. Pallavi CH. Nutrient management in finger millet (*Eleusine coracana* L.) under Melia azedarach based agri- silvisystem. M.Sc. (Ag) Thesis. Acharya N.G Ranga Agricultural University, Hyderabad, India, 2014.
9. Prakasha G, Kalyana murthy KN, Rohani N, Meti, Jagadish Prathima AS. Nutrient uptake and economics offingermillet (*Eleusine coracana* (L.) Gaertn) under guni method of planting in eastern dryzone of Karnataka. International Journal of Pure and Applied Bio Science. 2017; 596):144-151.
10. Prakasha G, Kalyana murthy KN, Rohani N, Prathima AS Meti. Effect of spacing and nutrient levels on growth and attributes and yield of finger millet (*Eleusine coracana* (L.) Gaertn] under guni method of planting in red sandy loamy soil of Karnataka, India. International Journal of Current Microbiology and Applied Sciences. 2018; 7(5):1337-1343.

11. Sudheendra Saunsi, Reddy VC, mallikarjuna Rajesh Rawal. Influence of enriched biodigester liquid manure on growth and yield of finger millet. *The Bio-scan*. 2014; 9(2):613-616.
12. Anonymous, 2017. Annual Report, All India coordinated small millet improvement programme on finger millet. ICAR, pp.33-37.
13. Dhanapal, G.N., Sanjay, M.T., Hareesh, G.R. and Patil, V.B. 2015. Weed and fertility management effects on grain yield and economics of finger millet following groundnut. *Indian J. Weed Sci.*, 47(2): 139-143.
14. Gill, H.S. and Vijayakumar. 1969. Weed index- a new method for reporting weed control trials. *Indian J. Agron.*, 14(1): 96-98.
15. Gomez, K.A. and Gomez, A.A. 1984. *Statistical procedures for agricultural research*. (2nd Edn.) Chichesler, UK: John Wiley and Sons.
16. Kujur, S., Singh, V.K., Gupta, D.K., Tandon, A., Ekka, V. and Agrawal, H.P. 2018. Influence of weed management practices on weeds, yield and economics of finger millet (*Eleusine coracana* L. Gaertn). *Int. J. Bioresour. Stress Mgmt.*,9(2): 209- 213.
17. Lall, M. and Yadav, L.M.S. 1982. Critical time of weed removal in finger millet. *Indian J. Weed Sci.*,14: 85-88.
18. Malemnganbi, T. and Lungdim, J. 2019. Influence of different doses of pyrazosulfuron-ethyl and establishment methods on the yield of lowland rice (*Oryza sativa* L.). *J. Crop and Weed*,15(1): 209-212.
19. Mani, V.S., Pandita, M.L., Gautam, K.C. and Bhagwandas. 1973. Weed killing chemicals in potato cultivation. *Indian Farm.*, 23: 7-13.