

ASSESSMENT OF EFFECT OF DIFFERENT PACKAGING MATERIALS ON MICROBIAL QUALITY OF LOCUST BEANS DURING STORAGE

ABSTRACT

This study assesses the effect of different packaging materials on the microbial quality of Africa Locust beans during storage. Locust beans (Dawadawa) also known as iru, among the Yorubas in South-west Nigeria, is a popular condiment used as taste and flavour enhancer in soup and dishes in Africa. Locust bean is traditionally produced from locust beans seed (*Parkia biglobosa*) and preserved using different packaging materials before use in order to prolong its shelf life. The Microbiological examination of the produced locust beans was carried out on samples wrapped with different packaging materials such as plastic container, nylon and dry banana leaves. The suspected organisms isolated include *Salmonella spp*, *Staphylococcus aureus*, *Streptococcus Lactis*, *Pseudomonas aeruginosa*, *Lactobacillus plantarium*, *Bacillus cereus*, *Bacillus subtilis*, *Leuconostocsp*, *Escherichia coli* and *Micrococcus sp*. It was observed from this study that locust bean packaged with banana leaves had the highest microbial load, few of which are of public health concern while plastic container had least microbial load. It is therefore recommended that locust beans should be produced and packaged with sealed plastic containers to avoid contamination by pathogenic organisms.

Comment [U1]: Are a popular

Comment [U2]: A taste and flavor

Comment [U3]: containers

Comment [U4]: ,and

Comment [U5]: lactis

Comment [U6]: beans

Comment [U7]: containers

Comment [U8]: the least

Comment [U9]: include keywords

INTRODUCTION

African locust beans (*Parkia biglobosa*) also known as iru, among the Yorubas in South-west Nigeria, is a popular condiment used as taste and flavour enhancer in soup and dishes in Africa.

Comment [U10]: are a popular

Comment [U11]: a taste and flavor

Dawadawais traditionally produced from locust beans (*Parkia biglobosa*) seeds (Farindeet *al.*, 2014). Odunfa (2010) stated that fermented locust bean seed is commonly consumed in Ghana, Nigeria, Sierra Leone and Togo. In Nigeria it is called iru in Yoruba, dawadawa in Hausa and ogiri `igala in Igbo. It is also referred to as “kinda” in Sierra Leone and kpalugu in Ghana. Preservation and preservatives are designed to inhibit/control the activities of spoilage causing organisms in food, a process also referred to as sanitization. African locust bean is a nutritious source of food since it is rich in protein and some beneficial health components. It serves as a cheap source of protein for people whose source of protein is low due to high cost of animal protein sources (Jay, 2010).

Comment [U12]: has traditionally

Comment [U13]: ,and

Comment [U14]: Spoilage - causing

Comment [U15]: The high

Food packaging is an integral part of food processing, which entails the use of some materials in the wrapping of foods (Ayo-Lawa *et al.*, 2016). The success of most preservation methods depends on appropriate packaging. However, faulty packaging will undo all that a good processor has attempted to accomplish by the most meticulous manufacturing process (Ihemejeet *al.*,2022).

Comment [U16]: through

Spoilage causing organisms due to their growth and metabolic activities produce by-products, which change the texture, taste, flavor and the aroma of the food (Jay, 2010). The use of packaging and preservation of locust beans is to make the product shelf life stay fresher and longer. Dehydration, salting and packaging of home-produced food items such as locust beans (iru) in simple polyethylene bags is desirable to extend their shelf life. (Farindeet *al.*, 2014).

Comment [U17]: Spoilage-causing

Comment [U18]: ,and

Comment [U19]: ,and

Comment [U20]: are

In Nigeria, African locust bean (*Parkia biglobosa*) tree grows widely throughout the savanna (Akoma *et al.*, 2001). A mature pod contains yellow, dry and powdery pulp in which dark brown seeds are embedded. The pulp is licked for its sweet taste but only to a small extent.

Comment [U21]: the African

Comment [U22]: ,and

The pulp is usually washed away when the seeds are processed into **condiment** called dawadawa or iru. Dawadawa is a source of protein intake among the low-income groups and rural populations of West Africa. While the seed has been extensively studied (Addy *et al.*, 1995), little has been done on the utilization of the pulp. In West Africa, the pulp is prepared as flour and used in soups and stews or eaten with cereals as porridge (Omauvbeet *et al.*, 2004).

Comment [U23]: A condiment

Plants are known to contain high amounts of essential nutrients, vitamins, minerals, fatty acids and fibre (Gafar and Itodo, 2011). Plants also contain other chemical compounds such as saponins, tannins, oxalates, phytates, trypsin inhibitors **and** cyanogenic glycosides, which are known as secondary metabolites and are biologically active (Soetan and Oyewole, 2009). The seeds on fermentation are used in cooking stew **and** soup. It has been reported that the **husks** and pods are good for livestock (Obizoba, 1998). The roots, barks, leaves, stems, flowers, fruits **and** seeds of *P. biglobosa* are all used medicinally to treat a range of ailments, including diarrhea ulcers, pneumonia, burns, coughs, etc (Sacande and Clethero, 2007).

Comment [U24]: and

Comment [U25]: and

Comment [U26]: That husk

Comment [U27]: and

Different packaging materials are used for wrapping locust beans in **order** to extend **its** shelf life and the microbial load of these packaging materials **are** mostly not assessed due to ignorance and little information on its effect on the product. It is therefore essential to assess the effect of different packaging materials used for the storage and preservation of the locust beans. This will ascertain the best packaging materials with the lowest microbial load suitable for **its** preservation. This will give information on how to improve the shelf life of locust beans and encourage its acceptability

Comment [U28]: Beans to extend

Comment [U29]: their

Comment [U30]: is

Comment [U31]: their

MATERIALS AND METHODS

Sample Collection and Preparation

Parkia biglobosa (Africa locust bean) seed used for this research were purchased from Oja-Oba market in Ado-Ekiti, Ekiti-State, Nigeria.

Comment [U32]: seeds

Sample Preparation

The production and fermentation process were carried out in the department of Food Technology laboratory, School of science and computer studies. Federal polytechnic Ado-Ekiti, Ekiti-State Nigeria

Comment [U33]: was

Comment [U34]: Department of Food Technology Laboratory

Production of fermented *Parkia biglobosa* (Africa locust bean) “iru” using traditional method.

Parkia biglobosa (Africa locust bean) seed was prepared according to the method of (Auduet *al.*, 2004). *Parkia biglobosa* (Africa locust bean) seed were sorted and washed to remove the yellowish pulp from the seeds. The locust bean seed was cooked with pressure pot. Put on electric cooker for 7 hours, with water being added frequently to it, until the seed becomes softer, then the seed was allowed to cool down and it was pounded by using mortar and pestle, the cotyledon was separated from its coat. Inside a bowl of water by using hand and sieve then the seed was washed and cleaned thoroughly. It was then re-cooked for another 1hour 30minutes with pressure pot, until the seed becomes softer. Then the water was drained off by using sieve. The locust bean was spread into a big sieve wrapped with multiple cloth to maintain a hot atmosphere to be fermented for 72hours. After fermentation the locust bean “iru” was packaged in three different materials (Banana leaf, Plastic rubber and nylon) and preserved for 7 days.

Comment [U35]: seeds

Comment [U36]: a pressure

Comment [U37]: eliminated by

Comment [U38]: a hand

Comment [U39]: 1 hour 30 minutes

Comment [U40]: A sieve

Media Preparation

The media (Nutrient agar, MacConkey agar, Salmonella shigella agar and Eosin Methylene blue agar) used for this research were prepared according to the manufacturer's

instruction. The prepared media were poured into conical flasks and the flasks were covered with cotton wool and Aluminum foil. The media were sterilized and autoclaved at 121⁰c for 15minutes and used for microbial analysis (Temple, *et al.*, 2006).

Comment [U41]: instructions

Comment [U42]: 15 minutes

Microbiological Analysis of Fermented Sample

One gram of the sample (Fermented *Parkia biglobosa* (Africa locust bean) were taken respectively and marched using mortar and pestle for proper homogenization. One gram of sample was diluted serially in seven folds dilution blanks and properly mixed with sterile glass rod. 0.1ml of diluted sample was introduced into sterile plates and agar. The media was poured at 45⁰C. The plates were rotated gently to disperse the inoculum in medium and allowed solidify. Then the plates were incubated at 37⁰C and the resulting colonies of microbes were sub-cultured to get the colonies in their pure form and to lower the density of cells. The culture was maintained as slants on both nutrient Agars. The slants were kept and stored for further use.

Comment [U43]: Africans

Comment [U44]: was

Comment [U45]: fold

Comment [U46]: the sample

Comment [U47]: the medium

Comment [U48]: to solidify

Examination of Culture Plates of Isolated

Macroscopic: the colonial appearance of the organism was noted such as colour, shape and size after 24hours of incubation.

Comment [U49]: color, shape, and shape and size after 24 hours

Characterization and identification of sample

Gram Staining reaction was carried out by emulsifying one isolate 24hour old colony in a drop of water placed at the center of a clean grease tree slide passed through a Bunsen burner. The heat fixed smear was flooded with crystal violet for 60seconds, after which the stain was poured off the slide and rinsed with running tap water. The smear was flooded with iodine.

Comment [U50]: The Gram

Comment [U51]: 24 hours

Comment [U52]: Heat-fixed

Comment [U53]: 60 seconds

Identification of Bacteria Isolates

The bacterial isolates were identified by using morphological and biochemical tests such as coagulase test, catalase test, oxidase test, indole test, urea production, nitrate reductions, motility test, spore test, hydrogen sulfide production test, oxidase production, citrate utilization test, nitrate reduction test, VogesProskauer and Methyl red test. (Fawole and Oso, 2004) and Olutiola *et al.* (2000)

Comment [U54]: Include a section of ethical aspects and disclosure of conflict of interest

RESULTS

The results of the biochemical characterization of the microorganisms isolated from locust bean wrapped with different packaging materials such as plastic container, nylon and dry banana leaves are shown in the tables below.

Comment [U55]: beans

Table 1: Biochemical characteristics of isolates from African Locust beans package with plastic container

Isolates	Grams Rxn	Shape	Catalase	Indole	Vp	Oxidase	Methyl red	Citrate	Probable organisms
1	-	rod	+	+	-	-	+	+	<i>Salmonella spp.</i>
2	+	Cocci in cluster	+	+	-	+	+	+	<i>Staphylococcus Aureus</i>
3	-	rod	+	-	-	+	+	+	<i>Pseudomonas aeruginosa</i>
4	+	rod	+	-	+	-	-	+	<i>Bacillus Cereus</i>

Comment [U56]: spp.

Comment [U57]: aureus

5	+	rod	+	-	+	+	-	+	<i>Bacillus subtilis</i>
---	---	-----	---	---	---	---	---	---	--------------------------

Key:

+ means positive

- Means negative

Table2: Biochemical characteristics of isolates from African Locust beans packaged with nylon

Isolates	Grams Rxn	Shape	Catalase	Indole	Vp	Oxidase	Methyl red	Citrate	Probable organisms
1	-	rod	+	+	-	-	+	+	<i>Salmonella spp</i>
2	+	Cocci in chains	-	-	-	+	+	-	<i>Streptococcus Lactis</i>
3	-	rod	+	-	-	+	+	+	<i>Pseudomonas aeruginosa</i>
4	+	Bacilli	-	-	-	-	-	-	<i>Lactobacillus plantarium</i>
5	+	rod	+	-	+	-	-	+	<i>Bacillus</i>

Cereus

6	+	rod	+	-	+	+	-	+	<i>Bacillus subtilis</i>
7	-	cocco bacilli	-	+	+	-	-	-	<i>Leuconostocsp</i>
8	+	Spheri cal	+	-	+	+	-	+	<i>Micrococcus sp</i>

Key:

+ means positive

- Means negative

Table 3: Biochemical characteristics of isolates from African Locust beans from Leaves

Isolates	Grams Rxn	Shape	Catalase	Indole	Vp	Oxidase	Methyl red	Citrate	Probable organisms
1	-	rod	+	+	-	-	+	+	<i>Salmonella spp</i>
2	+	Cocci in cluster	+	+	-	+	+	+	<i>Staphylococcus Aureus</i>
3	+	Cocci	-	-	-	+	+	-	<i>Streptococcus</i>

		in							<i>Lactis</i>
		chains							
4	-	rod	+	-	-	+	+	+	<i>Pseudomonas aeruginosa</i>
5	+	Bacilli	-	-	-	-	-	-	<i>Lactobacillus plantarium</i>
6	+	rod	+	-	+	-	-	+	<i>Bacillus Cereus</i>
7	+	rod	+	-	+	+	-	+	<i>Bacillus subtilis</i>
8	-	cocco bacilli	-	+	+	-	-	-	<i>Leuconostocsp</i>
9	-	rod	+	+	-	-	+	-	<i>Escherichia coli</i>
10	+	Spheri cal	+	-	+	+	-	+	<i>Micrococcus sp</i>

Key:

+ means positive

- Means negative

DISCUSSION

The suspected isolates recorded in table 1 includes *Salmonella* spp, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Bacillus cereus* and *Bacillus subtilis*. *Salmonella* species has been reported to be transmitted through contaminated water and this bacterium is suspected carrier found in food handlers (Bukeret *et al.*, 2012). *Bacillus* species were the predominant microorganisms in this sample. The result of this study justifies the assertion of Ogbuonye (2018) that *Bacillus* species was the predominant bacteria involved in the fermentation of ogiri which is also a fermented soup condiment. *Pseudomonas* isolated in this sample is an aerobic organism and thrives well in the presence of oxygen and water (Ukaoma *et al.*, 2018). The locust bean seed usually comes with a substantial microbial local comprising spores of aerobic spore forming bacterial.

Comment [U58]: Table 1

Comment [U59]: have

Comment [U60]: is a suspected

Comment [U61]: spore-forming

The result on table 2 shows the characteristics of isolates from those packaged with nylon. Suspected organisms include; *Salmonella spp*, *Streptococcus lactis*, *Pseudomonas aeruginosa*, *Lactobacillus plantarium*, *Bacillus cereus*, *Bacillus subtilis*, *Leuconostoc sp* and *Micrococcus sp*. Eight isolates were isolated from the sample. *Bacillus spp*. which is a predominant organism obtained in this study and *Leuconostoc sp*. are capable of increasing the protein and fat contents of locust beans. This is similar to the study of Ogbadu and Okagbue (1988) who observed that various *Bacillus* species were responsible for the fermentation of African Locust Bean seeds. The result obtained from this study is similar to the work of IHEMEJE *et al.*, 2022 who also isolated *Lactobacillus*, *Pseudomonas* and *bacillus* from fermented melon seed.

Comment [U62]: ,and

The result from table 3 shows the biochemical characteristics of isolates from African Locust beans packaged with dry banana leaves. The suspected isolates are *Salmonella spp*, *Staphylococcus aureus*, *Streptococcus Lactis*, *Pseudomonas aeruginosa*, *Lactobacillus planetarium*, *Bacillus cereus*, *Bacillus subtilis*, *Leuconostoc sp*, *Escherichia coli* and

Comment [U63]: Table

Comment [U64]: lactis

Micrococcus sp. The microorganisms isolated from locust beans packaged with dry banana leaves might be due to the contamination from leaves used as packaging materials. The leaves are usually picked up from the ground and on the farm land and are not normally washed thereby introducing different microorganisms from soil and insect droppings. Studies by (Antia and Ibrahim 2000) and (Odunfa1983) has shown that several microorganisms are associated with locust bean fermentation. He also noted that the most abundant and the major dominant agent of fermentation after 24hours was *Bacillus subtilis*. They also noted the presence of *Leuconostocmesenteroides* and *Staphylococcus* species after fermentation. Similar report also showed that *Bacillus subtilis*, *Leuconostocmesenteroids*and *Staphylococcus* sp. were associated with the fermentation of African locust bean seeds to produce “iru” condiment (Omafuvbeet *al.*, 2002). However, the presence of *Escherichia coli* and *Staphylococcus aureus* are of public health concern and can lead to foodborne diseases. The microbes may be introduced into the products through the contaminated hands of handlers, unsterilized utensils, other materials being used for the processing and even from the air from the environment during the preparation of the product(Nwaguet *al* 2011; Tope, 2013). Achi (2005)

In conclusion, the results from this study **shows** that the use of plastic container as a packaging material should be **encourage** when compare to nylon and leaves for packaging of locust beans before consumption. Locust beans packaged with dry banana leaves harbors a lot of microorganisms that are **able** to cause foodborne diseases which are therefore of public health concern. It is therefore recommended that locust beans produced must be well preserved with the best packaging materials in order to prevent the entry and contamination by microorganisms

Comment [U65]: show

Comment [U66]: be encourage

Comment [U67]: that can cause

REFERENCES

Achi O K (2005) Traditional fermented protein condiments in Nigeria. African Journal of Biotechnology, 4(13):1612-1621.

Adedokun A. A. (2006): Contributions of Locust Bean (*Parkia biglobosa*. Jacq. Berth) Seeds Production and Marketing to the Household Economy of Kajola Local Government Area, Oyo State. B. Sc. Project Report Submitted to Department of Forest Resources Management, University of Ibadan (unpublished). Pp. 13-14.

Comment [U68]: adjust bibliography in Vancouver format

Akoma, O.; Olawepo, O. and Ogunrinde, B.A. 2002. The Production of „Tsamiya“ Wine from Tamarindus Indica. J. Chem. Soc. Nig. 27 (1): 17-19.

Alabi, D. A; Akinsuhre, O. R; and Sanyaolu, M. A. (2005): Quantitative Determination of Chemical and Nutritional Composition of *Parkia biglobosa*(Jacq.)Berth. African Journal of Biotechnology 4 (8) 812815. www.univ.academicjournals.org/AJB. Pp. 4-6.

Antai, S.P and Ibrahim, M.H. (1988) Microorganisms associated with African locust bean (*Parkia filicoided*) fermentation for ‘dawadawa’production. J. Applied Bacteriology 61:145-148.

Audu, I.; Oloso, A.; and Umar, B. 2004. Development of a Concentric Cylinder Locust Beans Dehuller. Agricultural Engineering International: the CIGR Journal of Scientific Research and Development. Manuscript PM 04 003 .VI: 2004. 11pp.

Ayo-Lawal, R. A., Osoniyi, R. O., Rivera, G. and Elufisan, T. O. (2016). Fermented Condiments from Melon (*Citullus vulgaris*) and African Locust Bean (*Parkia biglobosa*) seeds-A review. Nigerian Food Journal, 34(2):94-104.

- Bukar, A (2012). Preservative properties of extracts of *Parkia biglobosa* (JACQ) Benth., *Anogeisusleiocarpus* L. and *Moringa oleifera* lam. On some minimally and fully processed foods. PhD. Thesis. Department of Microbiology Bayero University Kano
- Farinde, E. O., Adeniran, H. A. and Abiose, S. H. (2014). Comparative Microbial Assessment of Fermented Lima Bean (*Phaseolus lunatus*) and Locust Bean (*Parkia biglobosa*) in Production of Daddawa. *British Microbiology Research Journal*, 4(7):772-784
- Fawole, M.O. and B.A. Oso, 2004. Characterization of Bacteria: Laboratory Manual of Microbiology. 4th Edn., Spectrum Book Ltd., Ibadan, Nigeria, pp: 24-33
- Gafar MK, Itodo AU (2011). Proximate and mineral composition of hairy indigo leaves. *Electronic Journal of Environmental, Agricultural and Food Chemistry* 10(3): 2007-2018.
- Ihemeje, A, Uneanya C.G, Asuzu C.C and Nnamocha, T.O (2022). Effect of Different Packaging Material on the Quality of Fermented Melon Seeds. *European Journal of Science, Innovation and Technology*. 2(2): 115-129
- Jay, M.J. (2000): *Modern Food Microbiology*. 4Th ed. Chapman and Hall Inc. New York. Pp. 187 3 195.
- Kourouma, K; Jean C.G; Achille, E.A and Clement,A.(2011): Ethnic differences in use values and use patterns of *Parkia biglobosa* in Northern Benin. Pp. 3. Publisher: *Journal of Ethnobiology and Ethnomedicine*. Cotonou, Benin Republic.
- Nwagu TNT and Orji MU, Nwobodo I and Nwobodo HA 2011 Mixed microbial flora as starter for the production of ugba from African oil bean seed. *Asian Journal of Biological Research*, 4 (1), pp. 62-69.

- Obizoba IC (1998). Fermentation of African locust bean. Text on nutritional quality of plant fruits. (Edn) Osagie, Eka (2000). Post-harvest Res. Unit. Dept. Biochem. Uniben, Nigeria. Pp. 160-198. 8.
- Odunfa VSA (1983). Carbohydrate changes in fermenting locust bean (*Parkia filicoidea*) during preparation. Qual Pl Hum Nutr 32:45-52
- Odunfa, S.A (1981): Micro-organism Association with Fermented African Locust Bean during Iru preparation 3.Plant Food 25. Pp. 245-250.
- Ogbadu L, Okagbue R (1988). Bacterial fermentation of soybeans for 'Daddawa' production. Journal of Applied Bacteriology 65:353-356.
- Ogbuonye, E. (2018). Chemical and Microbial Evaluation of 'Ogiri' (A Locally Fermented Food Condiment) Produced from Kersfing Groundnut Seeds. Sci. Arena Pub. Spec. J. Bio. Sci., 4(2), 7-13.
- Olutiola, P.O., O. Famurewa and H.G. Sonntag, (2000). Introduction to General Microbiology: A Practical Approach. 2nd Edn., Bolabay Publications, Ikeja, Nigeria
- Omafuvbe BO, Falade OS, Osuntogun BA (2004). Chemical and biochemical changes in African locust bean (*Parkia biglobosa*) and melon (*Citrullus vulgaris*) seeds during fermentation to condiments. Pak J Nutr 3:140-145.
- Omafuvbe, B.O.; Falade, O.S.; Osuntogun B.A. and Adewusi S.R.A.(2004). Chemical and Biochemical Changes in African Locust Bean (*Parkia Biglobosa*) and Melon (*Citrullus Vulgaris*) Seeds during Fermentation to Condiments. Pakistan Journal of Nutrition 3(3): 140-145.

- Sacande M, Clethero C (2007) *Parkia biglobosa* (Jacq) G. Don. Millenium seed bank Project
Kew. Seed leaflet No. 124
- Sadiku, O. A (2010): Processing Methods Influence the Quality of Fermented African Locust
Bean (Iru/ogiri/dawadawa) *Parkia biglobosa* Pp. 1. Publisher: Journal of Applied
Sciences Research, Ibadan, Nigeria
- Soetan KO, Oyewole OE (2009). The need for adequate processing to reduce the anti-nutritional
factors in animal feeds: A review. *Afr J Food Sci* 3(9): 223-232
- Tope A K 2013 Microbial and nutrient studies of fermented cooked lima bean (*Phaseolus
lunatus*) seeds. *Global Journal of Biology, Agriculture and Health Sciences*, 2(2): 94-101.
- Ukaoma, A., Nwachukwu, M., Adjeroh, L., Ezeanowai, C., Iwu, I., &Duru, M. (2018).
Comparative Study on Microbial Loads of Fermented Castor Oil Seeds (*Ricinus
communis*) and Melon Seeds (*Citrullus vulgaris*). *Intl. J. Sci. Res.*, 8(8), 83-89.