

# Association study and direct - indirect effects of characters for yield in rice (*Oryza sativa* L.)

## Abstract

The results were indicated that the seed yield per plant showed significant positive correlation with most of the traits like days to 50% flowering, plant height, filled grains per panicle, number of grains per panicle, spikelet fertility, flag leaf width, numbers of tillers per plant, number of effective tillers per plant, panicle length and 100 seed weight, representing importance of these traits for development of high yielding varieties whereas the trait spikelet sterility has been reported negative and significant association with yield trait and unfilled grains per panicle and flag leaf length reported negative non-significant correlation with yield. The of results of path coefficient analysis showed that the direct positive effect on yield were recorded by the traits, effective tillers per plant, 100 seed weight, Number of grains per panicle and most of the traits.

**Keywords:** Correlation, path coefficient analysis, rice (*Oryza sativa* L.).

## Introduction

Rice (*Oryza sativa*) is the most important cereal crop in the developing world and is the staple food of over half the world's population. The genus *Oryza* originated in the Gondwanaland continents (Chang, 1976). India ranks first in rice area and second in rice production, after China. Asian countries produces 89 % of world's rice and 50% of the people of world depends on rice as major food (FAO, 2016). Since Vedic period, rice has influenced the religions, traditions, culture and life style of a large number of people on earth. The cultivation of rice is practiced under more diverse conditions than any other food crop, ranging from irrigated (lowland), rainfed, upland, to deep water logged conditions. Rice however, compares favorably with other cereals in amino acids content. The biological value of its protein is high. The fat content of rice is low (2.0 to 2.5 per cent) and much of the fat is lost during milling. Rice contains a low percentage of calcium. Rice grain contains as much B group vitamins as wheat. Milled rice loses valuable proteins, vitamins and minerals in the milling process during which

Comment [M1]: Delete

Comment [M2]: a significant

Comment [M3]: the importance

Comment [M4]: great

Comment [M5]: . In contrast, the

Comment [M6]: a negative

Comment [M7]: Path coefficient analysis results showed that the direct positive effect on yield was recorded by the traits, effective tillers per plant, 100 seed weight, Number of grains per panicle, and most of the traits.

Comment [M8]: Italic

Comment [M9]: Italic

Comment [M10]: Delete

Comment [M11]: produce 89 % of the world's

Comment [M12]: the world depend

Comment [M13]: the Vedic period

Comment [M14]: culture, and lifestyle

Comment [M15]: Earth

Comment [M16]: rainfed, and upland, to deep waterlogged conditions. The rice, however,

Comment [M17]: amino acid

Comment [M18]: percent

Comment [M19]: many B-group

the embryo and the aleuronic layer are removed. Much of the loss of nutrients can be avoided through parboiling process. (Anonymous, 2013a). Information on association of characters, direct and indirect effects contributed by each character towards yield will be an added advantage in aiding the selection process. Correlation and path analysis establish the extent of association between yield and its components and also bring out relative importance of their direct and indirect effects, thus giving an obvious understanding of their association with grain yield. Ultimately, this kind of analysis could help the breeder to design his selection strategies to improve grain yield. (Yadav *et al.*, 2018).

Comment [M20]: the parboiling

Comment [M21]: Delete

Comment [M22]: the association of characters, and direct

Comment [M23]: the relative

Comment [M24]: Please add the aim of conducting the research at the end of the introduction.

## Materials and Methods

The experimental material consists of 120 rice germplasm accessions along with five checks *viz.* Poomima, Mahamaya, Swarna, Indira Sugandhit Dhan-1 and Indira Aerobic Dhan-1. The criteria of selection of checks were the habits of heat tolerance and their yielding ability under diverse climatic conditions. The germplasm accessions were screened for heat tolerance and ratooning under natural conditions. The 120 rice germplasm and five checks used in the present investigation were obtained from germplasm unit of IGKV, Raipur. The simple correlations ( $r$ ) between different characters at phenotypic ( $p$ ) and genotypic ( $g$ ) levels were worked out as suggested by Searle (1961). Path coefficient analysis was carried out according to Dewey and Lu (1959).

Comment [M25]: for the selection of checks were the habits

Comment [M26]: the germplasm

## Result and Discussion

In general, genotypic correlations were higher than phenotypic ones in magnitude for all the characters. The characters which showed negative correlation at genotypic level also showed negative correlation at phenotypic level. The correlation co-efficient results showing that the traits *viz.* days to 50% flowering, plant height, filled grains per panicle, number of grains per panicle, spikelet fertility, flag leaf width, numbers of tillers per plant, number of effective tillers per plant, panicle length and 100 seed weight has significant positive association with seed yield per plant representing importance of these traits for development of high yielding varieties in rice. This type of significant positive association of traits with yield is also reported by some researchers in last decades (Rasheed *et al.*, 2002, Girish *et al.*, 2006, Rokonuzzman *et al.*, 2008). The trait spikelet sterility has been reported negative and significant association with yield

Comment [M27]: a negative correlation at the genotypic level also showed a negative correlation at the phenotypic level. The correlation coefficient results show that

Comment [M28]: a significant positive association with seed yield per plant representing the importance of these traits for development of great

Comment [M29]: the last

trait, whereas the traits unfilled grains per panicle and flag leaf length reported negative non-significant correlation with yield which is similar with findings of (Ullah *et al.*, 2011). On the basis of present findings the plant breeder can be encourage for development of high yielding cultivars by making the selection of higher number of filled grain per panicle with a reasonable balance for moderate plant height, moderate days to 50% flowering, higher number of grains per panicle and higher number of effective tillers per plant. Similar findings were reported by Akinwale *et al.*, (2020), Jeke *et al.*, (2021) and Manivelan *et al.*, (2022).

The of results of path coefficient analysis depicted that the traits viz. spikelet fertility (11.02), spikelet sterility (11.00), number of grains per panicle (1.75), number of effective tillers per plant (0.45), 100 seed weight (0.38), days to 50% flowering (0.30), flag leaf length (0.05), plant height (0.47), flag leaf width (0.41) and panicle length (0.21) is the main contributors to yield because these traits has direct positive effect on yield. In the past studies also found that the spikelet fertility had positive direct effect on yield followed by the number of grains per panicle (Yolanda and Das, 1995). The traits such as number of effective tillers per plant, days to 50% flowering and plant height has positive direct effect on yield (Akter *et al.*, 2010). On the basis of path coefficient analysis the residual effect was 0.50 representing the contribution of component traits on yield. Hence, we can say that the contribution of component traits (fourteen traits studied) is 50% on yield and the 50% contribution came from the traits which are not studied in path coefficient analysis. Similar findings were reported by Williams *et al.*, (2021) and Reetisana *et al.*, (2022).

**Comment [M30]:** a negative non-significant correlation with a yield which is similar to the findings of Ullah *et al.*, (2011).

**Comment [M31]:** Based on the present findings, the plant breeder can be encouraged to development of high yielding cultivars

**Comment [M32]:** a higher number of filled grains

**Comment [M33]:** a higher

**Comment [M34]:** 2011

**Comment [M35]:** Delete

**Comment [M36]:** are

**Comment [M37]:** have a direct

**Comment [M38]:** that spikelet fertility had a positive direct effect

**Comment [M39]:** the number

**Comment [M40]:** flowering, and plant height have a positive direct effect

**Comment [M41]:** Based on

**Comment [M42]:** analysis,

**Comment [M43]:** fourteen

**Comment [M44]:** Please add conclusions and recommendations at the end of the results and discussion.

## References

- Akinwale, M., Gregorio, G., Nwilene, F., Akinyele, B., Ogunbayo, S. A. and Odiyi, A. C. 2011. Heritability and correlation coefficient analysis for yield and its components in rice (*Oryza sativa* L.). *African J. of Plant Sci.*, 5: 207–212.
- Akter, T., Ivy, N. A., Rasul, M. G. and Mian, M. A. K. 2010. Variability and character association of reproductive traits in exotic rice germplasm. *Bangladesh J. Pl. Breed. Genet.*, 23(1): 39-43.
- Anonymous. 2013. Statistical data base on agriculture. [http:// apps. FAO.org](http://apps.FAO.org).
- Chang, T. T. 1976. The origin, evolution, cultivation, dissemination and diversification of Asian and African rice. *Euphytica*, 25:435-441.
- Dewey, D.R. and Lu, K.H.A. 1959. Correlation and path coefficient analysis of component in crested wheat grass seed production. *Agron. J.*, 5: 515-518.
- Girish, T. N., Gireesha, T. M., Vaishali, M. G., Hanamareddy, B. G. and Hittalmani, S. 2006. Response of a new IR 50/Moroberekan recombinant inbred population of rice (*Oryza sativa* L.) from an *indica* × *japonica* cross for growth and yield traits under aerobic conditions. *J. Euphytica*, 152(2): 149-161.
- Jeke, E., Mzengeza, T., Kyung, H. K. and Imani, C. 2021. Correlation and Path Coefficient Analysis of Yield and Component Traits of KAFACI doubled haploid Rice (*Oryza sativa* L.) Genotypes in Malawi. *Int. J. Agri. Tech.*, 1(2): 1-9.
- Manivelan, K., Juliet, H. S., Suresh, R., Theradimani, M., Renuka, R. and Gnanamalar, R. P. 2022. Inherent variability, correlation and path analysis in lowland rice (*Oryza sativa* L.). *Biological Forum – An Intl. J.*, 14(2): 771-778.
- Rasheed, M. S., Sadaqat, H. A. and Babar, H. A. 2002. Correlation and path coefficient analysis for yield and its components in rice. *Asian J. Pl. Sci.*, 1(3): 241-244.
- Reetisana, N., Sastry, E. V. D., Renuka, T., Julia, T. and Pyngrope, A. H. 2022. Correlation and path coefficient analysis in screening of submergence tolerance in rice (*Oryza sativa* L.) genotypes of Manipur. *Biological Forum – An Interl. J.*, 14(2): 1130-1135.

Rokonuzzaman, M., Zahangir, M. S. and Hussain, M. D. I. 2008. Genotype variability of components and their effects on the rice yield: Correlation and path analysis study. *Ital. J. Agron.*, 2: 131-134.

**Comment [M45]:** This scientific reference is not cited in the manuscript text.

Singh, R., Yadav, V., Mishra, D.N. and Yadav, A.2018. Correlation and Path Analysis Studies in Rice (*Oryza sativa L.*). 2018. Journal of Pharmacognosy and Phytochemistry. SP1: 2084-2090.

**Comment [M46]:** This scientific reference is not cited in the manuscript text.

Ullah, M. Z., Bashar, M. K., Bhuiyan, M. S. R., Khalequzzamana, M. and Hasan, M. J. 2011. Interrelationship and cause-effect analysis among morpho-physiological traits in birain rice of Bangladesh. *Int. J. Plant Breed. Genet.*,5: 246-254.

Williams, K., Mishra, Aman., Verma, A., Suresh, B. G. and Lavanya, G. R. 2021. Genetic Variability and Correlation Studies for Yield and Yield Related Traits in Rice (*Oryza sativa L.*) Genotypes. *Int. J. Curr. Microbiol. App. Sci.*, 10(01): 752-764.

Yolanda, J. L. and Das. L. D. V. 1995. Correlation and path analysis in rice (*Oryza sativa L.*) *Madras Agri. J.*, 82: 576-578.

UNDER PEER REVIEW

**Table 1: Correlation coefficient between yield and yield contributing traits**

Traits	D50	PH	FG	UG	NG	SF	SS	FLL	FLW	NT	NE	PL	100S	SY
D50	1.000													
PH	0.520**	1.000												
FG	0.610**	0.430**	1.000											
UG	0.130*	0.389**	0.266**	1.000										
NG	0.547**	0.508**	0.918**	0.626**	1.000									
SF	0.099 <sup>NS</sup>	-	0.239**	-	-0.137*	1.000								
SS	-0.099 <sup>NS</sup>	0.175**	-	0.806**	0.137*	-	1.000							
FLL	-	0.011 <sup>NS</sup>	-	-0.004 <sup>NS</sup>	-	-0.112 <sup>NS</sup>	0.112 <sup>NS</sup>	1.000						
FLW	0.266**	0.437**	0.201**	0.215**	0.276 <sup>NS</sup>	-0.072 <sup>NS</sup>	0.072 <sup>NS</sup>	0.083 <sup>NS</sup>	1.000					
NT	-0.008 <sup>NS</sup>	-	0.007 <sup>NS</sup>	-0.134*	-0.049 <sup>NS</sup>	0.094 <sup>NS</sup>	-0.094 <sup>NS</sup>	-	-	1.000				
NE	-0.035 <sup>NS</sup>	0.214**	-0.024 <sup>NS</sup>	-	-0.092 <sup>NS</sup>	0.130*	-0.130*	-	-	0.978**	1.000			
PL	0.118 <sup>NS</sup>	0.269**	0.359**	0.406**	0.457**	-	0.167**	-0.031 <sup>NS</sup>	0.349**	-0.264**	-0.295**	1.000		
100S	-0.153*	-	-	-	-	0.168**	-	0.204**	-0.090 <sup>NS</sup>	-0.070 <sup>NS</sup>	-0.038 <sup>NS</sup>	-0.254**	1.000	
SY	0.460**	0.249**	0.208**	0.442**	0.350**	0.304**	0.304**	-0.063 <sup>NS</sup>	0.178**	0.172**	0.174**	0.191**	0.239**	1.000

Here, D50 = days to 50% flowering, PH = plant height, FG = filled grains per panicle, UG = unfilled grains per panicle, NG = number of grains per panicle, SF = spikelet fertility, SS = spikelet sterility, FLL = flag leaf length, FLW = flag leaf width, NT = numbers of tillers per plant, NE = number of effective tillers per plant, PL = panicle length, 100S = 100 seed weight and SY = seed yield per plant.

**Table 2: Path coefficient of yield contributing traits on seed yield of rice germplasm accessions**

Traits	D50	PH	FG	UG	NG	SF	SS	FLL	FLW	NT	NE	PL	100S	Correlation With SY
<b>D50</b>	<b>0.30759</b>	0.02453	-	-	0.95941	1.09701	-1.09346	-	0.00946	0.00134	-	0.02564	-	<b>0.460</b>
<b>PH</b>	0.16004	<b>0.04714</b>	-	-	0.89102	-1.92874	1.92287	0.00059	0.01809	0.03568	-	0.13170	-	<b>0.280</b>
<b>FG</b>	0.18754	0.02029	<b>1.11800</b>	-	1.61088	2.64040	-2.63596	-	0.00961	-	-	0.07772	-	<b>0.484</b>
<b>UG</b>	0.04012	0.01833	0.29701	<b>0.77203</b>	1.09778	-8.89089	8.87327	0.00024	0.00891	0.02243	-	0.08777	-	<b>-0.062</b>
<b>NG</b>	0.16822	0.02394	1.02658	0.48310	<b>1.75433</b>	-1.51610	1.51245	-	0.01144	0.00821	-	0.09894	-	<b>0.367</b>
<b>SF</b>	0.03060	0.00825	0.26770	0.62246	0.24120	<b>11.02726</b>	11.00658	0.00607	0.00297	0.01568	-	0.03625	-	<b>0.271</b>
<b>SS</b>	-	0.00824	0.26775	-	0.24107	-	<b>11.00661</b>	0.00606	0.00297	0.01568	-	0.03624	-	<b>-0.271</b>
<b>FLL</b>	-	0.00051	0.22504	0.00343	-	-1.23349	1.22904	<b>0.05431</b>	0.00344	0.02781	-	0.07433	-	<b>-0.063</b>
<b>FLW</b>	0.07019	0.02058	-	-	0.48411	-0.79060	0.78860	0.00451	<b>0.04145</b>	0.03168	-	0.07561	-	<b>0.178</b>
<b>NT</b>	-	-	0.25930	0.16593	-	1.03505	-1.03359	-	-	-	-	0.08834	-	<b>0.172</b>
<b>NE</b>	0.00248	0.01007	0.00827	0.10368	0.08629	1.43555	-1.43355	0.00904	0.00786	<b>0.16701</b>	-	0.05716	-	<b>0.174</b>
<b>PL</b>	0.01075	0.01266	0.02631	0.13657	0.16089	1.43555	-1.43355	0.00893	0.00811	0.16330	<b>0.45181</b>	0.06376	-	<b>0.174</b>
<b>100S</b>	0.03645	0.02869	-	-	0.80206	-1.84729	1.84333	-	0.01448	0.04412	-	<b>0.21641</b>	-	<b>0.191</b>
	0.04699	0.01176	0.23286	0.34136	0.61430	3.35214	-3.34658	0.01107	0.00372	0.01176	0.01702	0.05502	<b>0.38475</b>	<b>0.239</b>
<b>Residual effect: 0.50359</b>														

**Note: Bold diagonal figures indicate the direct effect.**

Here, D50 = days to 50% flowering, PH = plant height, FG = filled grains per panicle, UG = unfilled grains per panicle, NG = number of grains per panicle, SF = spikelet fertility, SS = spikelet sterility, FLL = flag leaf length, FLW = flag leaf width, NT = numbers of tillers per plant, NE = number of effective tillers per plant, PL = panicle length, 100S = 100 seed weight and SY = seed yield per plant.