

Original Research Article

SPECTROPHOTOMETRIC ASSESSMENT OF BENZENE 1,3,5-TRIOLE IN PURE FORMS AND IN BULK SAMPLE BY DIAZOTIZATION COUPLING REACTION

Abstract: The method was published for determining microgram quantities of benzene 1,3,5-triol (phloroglucinol) in aqueous solution and some bulk samples using an easy, quick, and effective spectrophotometric method. The method uses a diazotization as well as coupling process in basic medium to produce a bright yellow water-soluble dye from Phloroglucinol and diazotized 4-Methoxy aniline That is stable and also has a maximum absorption wavelength of 420 nm. With a molar absorptivity of $1.5989 \times 10^4 \text{ l mol}^{-1} \text{ cm}^{-1}$ as well as index of Sand ell sensitivity 0.001 g.cm^{-2} , Beer's law is followed across a concentration range of (2-40) g.ml^{-1} of phloroglucinol. The ideal conditions for all color development are detailed, and even the suggested procedures for determining phloroglucinol in aqueous medium as well as some bulk sample have been successfully employed.

Keywords: Diazotization coupling , phloroglucinol, 4- Methoxy aniline, Spectrophotometric Assessment

Introduction

The bioactive chemical phloroglucinol (1,3,5-trihydroxybenzene or 1,3,5-benzenetriol) has been utilized as a smooth muscle relaxant. Radiation-induced

intracellular as well as destroy to cellular elements like lipid, DNA, as well as protein were dramatically reduced by phloroglucinol. A phenolic chemical is Phloroglucinol.




Figure 1. Phloroglucinol chemical structure

Phloroglucinol is a yellowish white or white crystal with a melting point at 215-219°C. It is slightly water soluble, alcohol soluble, as well as ether. soluble It is employed as an antispasmodic. Its antispasmodic action is greater versus muscle than smooth muscle under spastic conditions. Phloroglucinol is often used to treat colics caused by renal as well as biliary calculi, actually pain in the gastrointestinal tract or urinary tract, pain in the abdomen area of unknown origin, spastic disorders of the female genital order, and dysmenorrhea [1-4]. Phloroglucinol was a non-atropine antispasmodic drug utilized for its spasmolytic actions against the digestive and urinary tract via oral, rectal, intramuscular, or intravenous injection. Phloroglucinol is often utilized to treat aches associated with digestive system is working disorders, renal colics (pain associated with urine diseases), as well as gynecological pains. In the past, the medication was commonly used techniques women get pregnant [5-7].

By using the azo-coupling reaction in alkaline medium, the stable diazotized 4-methoxy aniline agent has been developed to assess. Phloroglucinol in aqueous solution as well as some bulk samples. The yellow outcome was computed utilizing spectrophotometry. The technique for analyzing is straight

forward, quick, as well as precise. It has been successfully used in previously research to determine Phloroglucinol in aqueous solution and various bulk samples in the plasma of human using HPLC/mass spectrometry [8]. additionally with gas chromatography/mass spectrometry [7,9], speed HPLC analyses of naphthodianthrones as well as phloroglucinol from *Hypercom perforatum* extracts are too reported [9-10] obtained by Titrimetric process Another technique that has been mentioned [11-14]. these added flow injection [15] as well as reversed-phase ion-pair chromatography [16].

However, in typical quality control laboratories, especially on an industrial scale, there is no simple and sensitive procedure to follow. As a result, there was still a demand for The technique for analyzing that could help calculate in bulk drugs., bodily fluid as well as commercial products Taking this into account, an attempt has been made to build a suitable, sensitive, as well as verified RP-HPLC method using UV detection for showing stability. The technique's suitability for testing pharmaceutical items has been confirmed. According to recommendations [4,8], the analysis results were validated.

Materials and Methods

Devices

The UV-Visible 160 electronic double-beam calculating spectrometer was used for every spectral as well as absorbance investigations. sensitive

balance (Sartorius BL210S) (Germany).

Bath water

Material and reagents

Every chemicals employed are of the greatest purity possible as well as have not been purified further.

Solution of benzene 1,3,5-triol 500 g.ml⁻¹

0.05g of purified chemical BDH is dissolved in 100 ml 100% ethanol to make this solution., The bulk sample solutions are made using the same process and then moved to a a black bottle containing they will be for at least one week. Fluka Chemical Material Company submitted another bulk sample of benzene 1,3,5-triol with a purity of 98 percent.

Diazotized 4- methoxy aniline 1x10⁻²M reagent solution

Taken and dissolved 0.003 gm of purified 4-methoxy aniline Fluke in deionized water, then introducing 1 M HCl BDH with volume 1 ml and shaking well, accompanied by 0.025 M sodium nitrite BDH with volume 2 ml as well as shaking completely, then diluting to 25 ml and cooling to around 5°C for 30 minutes.

Methodology

A sequence of volumetric flasks 25ml containing to increase volumes of phloroglucinol 500 g.ml^{-1} from the working standard solution were transmitted to support the concentration range $2\text{-}40 \text{ g.ml}^{-1}$ in finished dilution, then 0.1ml of 1M solution sodium hydroxide as well as 2 ml diazotized 4-methoxy aniline agent 0.001 M solution were increased as well as watered down to the mark with deionized water, well integrated. and let aside at the temperature of room for 20 minutes, at 420 nm, the absorbance of the yellow dye produced was compared to a blank sample that included all components excluding Phloroglucinol. It was decided to create a calibration curve.[17,18].

Results and Discussion

Investigation of the optimum reaction conditions: the effects of a variety of variables on the optical properties of an azo dye characteristics have been investigated, as well as the experimental conditions have been adjusted.

Reagent volume impact

The impact of diazonium agent 0.001M with volume between 0.1-5 ml on absorbance ferocity was investigated, as well as 2.5 ml was revealed to be control.**Figure 2.**




Figure 2. influence of reagent volume

Acid impact

It was discovered that the existence of acid enhanced the ferocity for the output composed, so various acids including HNO_3 , H_2SO_4 , CH_3COOH , and HCl were investigated, and it was discovered that every one of the studied acids gave nearly the same intensity, for that HCl was chosen, and it was discovered that 1.5 ml for 1M HCl gave a high sensitivity especially in comparison to those other acids, so it was chosen in following experiments [19].Table 1.

Table 1. influence of HCl acid volume

Volume of acid	Absorbance
0.5	0.442
1	0.674

1.5	0.771
2	0.768
2.5	0.764
3	0.766

Base impact

Because the dye generated in an alkaline medium grew more intense and stable, the influence of various alkaline solutions on the colored result was investigated, including sodium carbonate, potassium hydroxide, ammonium hydroxide, sodium acetate, and sodium hydroxide. Only when the reaction was performed in the sodium hydroxide presence did it achieve maximum sensitivity and stability. The impacts of different NaOH concentrations 0.1-4 M were investigated, with 1 M appearing to be the best. The impact of several quantities of 1 M NaOH was also investigated, ranging from 0.05 to 4 ml, with 0.2 ml being the most effective.

Effect of Order of Addition

The optimal order of additions for the maximum absorption was discovered to be (PHI+B+R) where (PHI= Phloroglucinol material, B=base, as well as R=reagent) which was used in further studies.

Temperature impact

The proposed method's finished product was investigated at various temperatures. The analysis shows that in the temperature range (0-80) °C, the absorbance values remain virtually constant, however at greater temperatures, the amount of absorbance decreases, signaling the breakdown of a product due to protracted heating. At room temperature 20 °C, the colorful product remained steady. As a result, in this procedure, room temperature is chosen. [20]. Figure 3.




Figure 3. influence of temperature on the color product

Reaction Time impact

After 20 minutes, the color ferocity acquired its maximal after the phloroglucinol was interacted instantly with the agent solution. As a result, a 20 minute development time was chosen as the best option in the overall method. For at least 3 hours, the color acquired remained steady.

The best experimental conditions for identifying Phloroglucinol have been identified. The

Diaz onium reaction took place in an acidic medium [8], and 1M hydrochloric acid was chosen [9], as well as the dye generated grew more powerful additionally an alkaline medium stable [10].

Absorption spectra

When ever a dilute solution of Phloroglucinol is coupled with diazotized 4-Methoxy aniline, the yellow colored dye is produced instantaneously. under the above-mentioned conditions, This indicates maximum absorption, in contrast to the colored reagent blank, which shows no absorption at maximum absorption at 420 nm .(Figure.4) The absorption spectra are displayed. The maximum absorption wavelength of 420 nm is still used in later studies.




Figure 4. Absorption spectra

A : Phloroglucinol (20 μg /ml1) with 4- methoxy aniline 1×10^{-2} M color product instead of reagent blank.

B : Reagent blank versus deionized water C: Pure Phloroglucinol versus absolute ethanol

Calibration curve

Under a correlation value of 0.9990 as well as an intercept of 0.0379, a linear relationship between the concentration and absorbance of Phloroglucinol was seen under perfect experimental circumstances spanning the concentration range of 2-40 $\mu\text{g.ml}^{-1}$ (Figure.5). Above 40 $\mu\text{g.ml}^{-1}$ of Phloroglucinol, a negative divergence from Beer's law was detected. The molar absorptivity was $5.2164 \times 10^3 \text{ l.mol}^{-1}.\text{cm}^{-1}$. And table (2) appeared Analytical characteristics of the Phloroglucinol assessment technique established.




Figure 5. Calibration curve of Phloroglucinol

Table 2. Analytical characteristics of the Phloroglucinol assessment technique established

Analytical parameters	Present method
The equation of regression	Y= 0.0414X + 0.1269
Linearity ($\mu\text{g ml}^{-1}$)	(2-40)
Correlation coefficient, r^2	0.9990
Limit of detection LOD ($\mu\text{g ml}^{-1}$)	0.598
LOQ ($\mu\text{g ml}^{-1}$)	1.451
(RSD Average) %	0.956
recovery Average %	100.823
Molar absorptivity ($\text{l mol}^{-1} \text{cm}^{-1}$)	5.2164×10^3
The sensitivity of Sandell ($\mu\text{g cm}^{-2}$)	0.001

Precision and accuracy

Phloroglucinol was measured at three distinct concentrations to establish the calibration graph's accuracy and precision. Table (3) means that the data are exact and accurate enough.


Table 3. Shows the proposed method's precision and accuracy.

No.	Conc. of Phloroglucinol $\mu\text{g ml}^{-1}$		Error %*	Recovery*	R.S.D %*
	present	found			
1	2	1.97	- 1.500	98.500	1.226
2	20	20.33	1.650	101.650	0.932
3	40	40.53	1.320	101.320	0.712

* Average for five assessments

Nature of product and reaction mechanism

Job's technique of the continuous of variations additionally also the ratio mole technique have been unitized to determine the structure (ratio of Phloroglucinol to diazotized 4-methoxy aniline agent) of the yellow azo dye produced. The dye was created by reacting Phloroglucinol with diazotized 4-methoxy aniline agent in a 1:1 ratio, according to the findings., (Figure.6&7), demonstrating a mono azo dye that adopts the following scheme:



Colored Azo dye

Scheme 1. color product formation




Figure 6. Continuous variation plot




Figure 7. Mole ratio plot

The azo dye's supposed stability constant in aqueous solution under the circumstances $2.16 \times 10^6 \text{ l.mole}^{-1}$ has been calculated under the conditions of the experimental procedure

The derived regression equation, as well as the procedure's analytical properties, are given in (Table 3). [21,22].

The technique's applicability

The proposed methodology are already used to detection and quantification of Phloroglucinol in aqueous solutions, and two sample of bulks containing it were

examined, as well as they provided good precision and accuracy, as indicated in the table (Table 3). The proposed method was found to be superior than the standard method . As indicated in table, the study's findings were statistically compared to the conventional strategy utilizing a variation precision ratio (F-test) additionally precision student (test) in the middle of confidence (95 percent) with two degrees of freedom .The F-test and T-test results were lesser than the theoretical amount (F=19.0, t=4.303). The amounts of (F = 2.732) as well as (t =1.023) were also lesser than the theoretical amount (F=19.0, t=4.303) for the technique under investigation. There was no significant contrast identified between the researched approach and the standard strategy (average of five examinations). Furthermore, as shown in Table 3, the approach under consideration is suitably compared to several previously described techniques. displaying the texts, corporate brand names, Phloroglucinol identification in bulk samples using the approved procedure, and recovery of the Phloroglucinol in the suggested approach [4,23-25].

Table 4. The use of the suggested as well as authorized techniques for assessing Phloroglucinol in bulk sampl

Phloroglucinol bulk Samples	Deliberated method		standard method		Official Values (t),(F)
	Recovery %	RSD %	Recovery %	RSD %	
Phloroglucinol pure	100.490	0.956	100.560	1.071	2.732
Phloroglucinol 98%	100.910	1.029	101.330	1.110	(F)Value = 19.0 1.023 (t)Value= 4.303

* Average for five determinations

Conclusions

Depending on its diazotized conjugating reaction with 4-methoxy aniline, a quick, quick, efficient, extremely accurate spectrophotometric technique for determining trace quantities of Phloroglucinol in aqueous solution has been devised, which does not require temperature control or solvent extraction.

References

1. Martindale W. Martindale. *The Extra Pharmacopoeia*. In J.E.F. 31st Edition. London: Royal Pharmaceutical Society, **1996**, pp.44-45.
2. Budavari, S., *The Merck index: an encyclopedia of chemicals, drugs and biologicals*. In M.J.O' Neil, A. Smith and P.E. Heckelman, 13th ed. Whitehouse Station, N.J.: Merck , **2001**, pp.7-8.
3. United States Pharmacopoeia Convention, Inc., *The United States Pharmacopoeia USP 27: The National Formulary: NF 22*, Washington, D.C.USA., April 12-16 , **2000**, pp.233-235.
4. *Stationary Office (Great Britain)*, **2009**. British Pharmacopoeia, 2009. London: Stationary Office, pp.4476-4478.
5. Sasaki, D. Kido, A. and Yoshida, Y., Effect of antispasmodic drugs on the colonic motility. II, 1984. clinical study in man, *International Journal of Clinical Pharmacology Therapy and Toxicology*, **1984**, 22(7), 338–341.
6. Tabassum, S. Afridi, B. and Aman, Z., Phloroglucinol for acceleration of labour: double blind, randomized controlled trial, *Journal of the Pakistan Medical Association*, **2001**, 55,(7), 270–273.
7. Louvel, D. Delvaux, M. Staumont, G., Intracolonic injection of glycerol: a model for abdominal pain in irritable bowel syndrome, *Gastroenterology*, **1996**, 110, (2), 351–361.
8. Cargnin, S. T., de Matos Nunes, J. Haas, J. S., Supercritical fluid extraction and high performance liquid chromatographic determination of benzopyrans and phloroglucinol

derivative in *Hypericum polyanthemum*, *Journal of Chromatography B*, **2009**,878, (1),83–87.

9. Cui Y. and Ang, C. Y. W., Supercritical fluid extraction and high-performance liquid chromatographic determination of phloroglucinol in St. John's wort (*Hypericum perforatum* L, *Journal of Agricultural and Food Chemistry*,**2002**,50 (10), 2755–2759.

10.Tolonen, A. Hohtola, A. and Jalonen, J.,Fast high-performance liquid chromatographic analysis of naphthodianthrones and phloroglucinol from *Hypericum perforatum* extracts, *Phytochemical Analysis*,**2003**,14(5), 306–309

11.Kim, H. Roh H., Lee H. J., Determination of phloroglucinol in human plasma by high-performance liquid chromatography-mass spectrometry, *Journal of Chromatography B*,**2003**, 792(2),307–312.

12.Bilia, A. R. Bergonzi, M. C. Mazzi, G. and Vincieri, F. F, Analysis and stability of the constituents of Artichoke and St. John's wort tinctures by HPLC-DAD and HPLC-MS, *Drug Development and Industrial Pharmacy*, **2002**,28(5), 609–619.

13.Lartigue-Mattei, C. Lauro-Marty, C. Bastide et al. M., Determination of phloroglucinol in human plasma by gas chromatography-mass spectrometry, *Journal of Chromatography*,**1993**, 617(1),140–146.

14.Amin D. and Bashir, W. A., Titrimetric determination of phenol resorcinol and phloroglucinol, *Talanta*,**2004**, 31(4), 283–284.

15. Xie C.-G. and Li,H.F.Determination of phloroglucinol by a new flow injection Chemiluminescence method, *Guang Pu Xue Yu Guang Pu Fen Xi*, **2004**, 24(12),1521–1523.

16. Hasan, N. Afridi, N. S. Khan, S. Siddiqui, M. Z. and Chauhan, M., Simultaneous determination of two antispasmodic drugs in bulk, pharmaceutical products and body fluid by a validated, acetonitrile free, cost effective and stability indicating reverse phase high performance liquid chromatographic method, *Analytical Chemistry*,**2012**, 11(7), 274–281.

17. Al-Rufaie, M.M.; Al-Sharefy, A.N.; Kathem, K.H. New spectrophotometric method for the determination chlorpromazine hydrochloride in pharmaceutical preparations by using oxidative coupling reaction. *Inter. J. Univ. Pharm. Bio. Sci.* **2013**, 2(4), 19-29.

18. Al-Rufaie, M.M., New spectrophotometric method for the determination of Sulfamethoxazole drug, *World Journal of pharmacy and pharmaceutical sciences*.**2016**,5(3), 172-180.

19. Al-Rufaie, M.M.; Jawad, A.A.; Sadiq, H.M. Colorimetric estimation for salbutamol sulphate in pure form and in different types of pharmaceutical. *Int. J. Chem. Tech. Res.*, **2016**, *9(11)*, 432-440.
20. Hanaa, K.A.T.; Al-Rufaie, M.M.; Zahraa, Y.M. Spectrophotometric determination of metoclopramide medicine in bulk form and in pharmaceuticals using orcinol as reagent. *Ovidius Univ. Ann. Chem.*, **2019**, *29(2)*, 85-93.
21. Al-Rufaie, M.M. Modern kinetic spectrophotometric procedure for estimation of furosemide drug as bulk form and in pharmaceuticals preparations. *Curr. Issues Pharm. Med. Sci*, **2016**, *29(4)*, 184-190.
22. Ahmed A.Z. and Al-Rufaie, M.M., Determination of trace amount of chlorpromazine hydrochloride in its pure form and in Pharmaceutical Preparations by using spectrophotometric analysis, *Sapporo Medical Journal*, **2021**, *55(3)*, 398-406.
23. Al-Rufaie, M.M. Al-labban, H.M.Y and Salih , N.S. ,Reduction and assessment of chloramphenicol antibiotic as pure from and in various kinds of pharmaceuticals by utilizing spectrophotometric approach, *Iranian Journal of Organic Chemistry*, **2017**, *9(2)*, 2087-2092.
24. Barrak, M. H., Al-Rufaie, M.M and Motaweq, Z.Y., Thymol quantitative analysis in medicinal formulation types through employing of nano-technology and antimicrobial activity in some pathogenic bacterial isolates, *Nusantara Bioscience* ,**2021**, *13(1)*, 129-137.
25. Gary D. Christian., *Analytical Chemistry.*, 6 th ed. *John Wiley & Sons Ltd.*, India, *3(4)*, **2004**, pp.12-13.