

Original Research Article

Allelopathic Effects of *Conyza Canadensis* Water Extract on Seed Germination of Three Crops

ABSTRACT

The study was designed to explore the allelopathic effects of *Conyza Canadensis* water extract (CCE) on seed germination in pakchoi, spinach and lettuce. Some indicators during the seed germination of the three crops were measured, and the allelopathic effect index was calculated under different concentrations of CCE (20, 40, 60, 80, 100 g/L) treatment. The results show that CCE has a varying degree of inhibitory effect on the germination rate of the seeds of the three crops. Spinach and lettuce have significant allelopathic effects. At higher concentration of CCE, the germination rate of spinach and lettuce significantly decreased ($P < 0.01$). After germination, the root length, seedling height, and fresh weight of the crop were measured, and the allelopathic effects were also calculated. It was found that the CCE had a significant inhibitory effect on root length, but it showed a promoting effect on seedling height and fresh weight of pakchoi, and CCE had a "low promotion-high inhibition" effect on seedling height and fresh weight of lettuce.

Keywords: Seed germination; *Conyza Canadensis* water extract; Allelopathy

1. INTRODUCTION

Allelopathy is mainly a phenomenon in which chemical substances produced and released by plants have a beneficial or adverse effect on their own or other biological life activities. Chemical substances released by plants can enter the environment through leaching and other forms, and are widely present in farmland systems.

Conyza Canadensis is currently considered a typical invasive species and a typical pioneer species in ecological succession. *Conyza Canadensis* secretes agrochemicals into the environment and has an effect on the growth of surrounding plants. Seed germination is an important link in the process of plant growth and development [1], and allelochemicals have a promoting or inhibiting effect on the germination of plant seeds and the growth and development of seedlings [2].

Conyza Canadensis is often found on the edges of farmland, abandoned agricultural land, or abandoned mining land, so its effects on crops, especially the germination of crop seeds, have an undeniable impact on agricultural production and sustainability. Therefore, this study selected three crops that are widely cultivated and in high demand in the market - spinach, pakchoi and lettuce - to study the allelopathy of *Conyza Canadensis* on their seed germination in order to provide more theoretical support for the allelopathy of *Conyza Canadensis* on crop seeds.

2. MATERIAL AND METHODS

2.1 EXPERIMENTAL MATERIALS

Conyza Canadensis was taken from the abandoned agricultural land at the southeast corner of Shandong agricultural engineering University (Zibo, Shandong) in October 2022, and dried naturally. Retrieve and clean before using.

Spinach, pakchoi and lettuce seeds were purchased from Zibo Seed Station.

Comment [i-1]: Please modify your title follows;
Allelopathic Effects of *Conyza canadensis* Water Extract on Seed Germination and Seedling Growth of Three Crops.

Comment [i-2]: Omit dot.

Comment [i-3]: 1. According to the [Asian Journal of Agricultural and Horticultural Research](#), the abstract should be written in Arial font, and the font size is 10. Please check it!
2. An abstract should include the following information: a little information about the background, the study's objectives, the methodology, the results and discussion, and finally, the conclusion.
3. But in this article, the objective, methodology, and conclusion parts are totally lost... Please incorporate this valuable information into your abstract.
4. Justify

Comment [i-4]: Omit.

Comment [i-5]:

Comment [i-6]: 1. Reference
2. Please add space after each paragraph before starting a new one..

Comment [i-7]: Write in small letters; species names should be written in small letters. Please check every point in the body of this paper.

Comment [i-8]: 1. Reference.
2. Your introduction is too short. Please clearly discuss each point and add more references.
3. The introduction should introduce the problem, why the research is going to be conducted, the significance of the study in general, and the aims of the study in particular.
• However, in this article, most of the relevant information, such as research problems and answers to why the research area is interesting, is uncovered.

Comment [i-9]: 1. Should be Arial, Bold, 11 font, left aligned, caps
2. Please add space after each paragraph before starting a new one..
3. It is better if you say plant materials in place of experimental materials.
4.

Comment [i-10]: second level heading should be written in ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined

2.2 EXPERIMENTAL DESIGN

(1) Preparation of extraction solution

- ① Crushing. Cut the dried *Conyza Canadensis* into approximately 5 mm pieces.
- ② Extraction. Immerse the crushed *Conyza Canadensis* sample (100 g) in deionized water (900 ml) and stir well. The samples were immersed at room temperature for 48 h and ultrasounded for 30 min.
- ③ Filter. The original liquor 100 g/L of CCE was obtained after coarse filtration with gauze and vacuum extraction, and stored at 4 °C.
- ④ Preparation of experimental treatment solution. During the experiment, the original liquor was divided into different concentrations of treatment liquid (20 g/L, 40 g/L, 60 g/L, 80 g/L).

(2) Seed pretreatment

Soak and disinfect the seeds with 75% alcohol for 5 min, then rinse thoroughly with deionized water. Seed-soaking. After soaking the seeds in deionized water for 12 hours, seeds with full, uniform seeds were selected for use.

(3) Seed germination test

The soaked cabbage, cabbage and spinach seeds were spread evenly in a pasteurized petri dish covered with filter paper. Five concentrations of CCE (20 g/L, 40 g/L, 60 g/L, 80 g/L, 100 g/L) were set and water was used as control (CK-Z) for seed germination experiment. Three replicates were set for each treatment with 40 seeds per replicate. The culture was carried out in a constant temperature incubator at 25 °C.

During the seed cultivation period, CCE shall be quantitatively (5 ml) applied to each petri dish daily, and added irregularly according to water consumption. A seed is considered to have germinated when its embryonic root breaks through the seed coat. Seed germination rates are regularly observed and recorded on a daily basis, with a germination period of 7 days.

After 7 days, 3-6 typical samples of each crop were selected for each treatment to determine root length, seedling height, and fresh weight.

Table 1. Germination rate test treatment

Treatment	CCE (g/L)
CK-Z	Deionized water
T1	20
T2	40
T3	60
T4	80
T5	100

2.3 PROCESSING OF TEST RESULTS

The germination rate is calculated as follows.

Germination rate (GR) = total number of germinated seeds within 7 days / total number of tested seeds × 100% (1)

$$\text{Germination (\%)} = \frac{\text{Number of seeds germinated}}{\text{Number of seeds placed for germination}} \times 100$$

Comment [i-[11]: make heading
2.2.1. Preparation of extraction solution
2.2.2. Seed pretreatment
2.2.3. Seed germination test

Comment [i-[12]: check it!

Comment [i-[13]: Please specify how many seeds were used per petriplate

Comment [i-[14]: Check it

Comment [i-[15]: Please indicate how root length, seedling height, and fresh weight were determined. Please refer Hailu, M., Muthuswamy, M. and Petros, Y., 2019. Effects of Tannery Effluent on Seed Germination and Growth Performance of Selected Varieties of Maize (*Zea mays* L.). *International Journal of Scientific and Research Publications*, 9(3), pp.242-251.

Comment [i-[16]: Use this formula:
Germination (%) =

$$\frac{\text{Number of seeds germinated}}{\text{Number of seeds placed for germination}} \times 100$$

The response index (RI) is calculated as follows:

$$RI = 1 - C/T \quad (T \geq C) \text{ or } RI = T/C - 1 \quad (T < C) \quad (2)$$

In Eq.2, C is the control value; T is the value of the treatment. RI represents the strength of allelopathy, with a positive value indicating a boosting effect and a negative value indicating an inhibiting effect. Its absolute value reflects the strength of the allelopathy.

The comprehensive allelopathic effects (CAE) represents the arithmetic mean value of each test index RI of the same tested plant treated with CCE [4]:

$$CAE = (RI_1 + RI_2 + RI_3 + RI_4) / 4 \quad (3)$$

In Eq.3, RI₁ to RI₄ are the allelopathic effect indices of germination rate, root length, seedling height and fresh weight, respectively.

Comment [i-[17]: Clearly write down the formula by using equation tools.

2.4 DATA PROCESSING AND STATISTICAL ANALYSIS

Excel 2019 was used to perform mapping, variance analysis and significance testing on the experimental data. Single factor analysis of variance was used to compare the significance of differences between different treatments.

3. RESULTS AND ANALYSIS

3.1 THE EFFECT OF CCE ON THE GERMINATION AND ALLELOPATHIC EFFECT INDEX OF CROP

As the germination of spinach seeds was seriously affected under the treatment of the concentration of water extract of each *Conyza Canadensis*, at the end of the experiment, the germination of spinach seeds only broke through the seed coat and did not grow into seedlings, so the data of root length and seedling height could not be measured. Therefore, there are no other indicators for spinach determination data except GR.

Table 2. The Effect of CCE on the Germination Rate and Allelopathic Effect Index of Crop

Crops	Concentration (g/L)	GR (%)	RI ₁	Root length (mm)	RI ₂	Seedling Height (mm)	RI ₃	Freshweight (mg)	RI ₄
Pakchoi	CK-Z	100±0.00	-	58.55±8.04	-	37.75±4.30	-	30.65±3.65	-
	T1	98.33±0.47	-0.02	48.69±9.05	-0.17	48.51±7.01	0.22	41.37±4.88*	0.26
	T2	98.33±0.47	-0.02	32.88±7.81	-0.44	61.01±11.13**	0.38	55.78±4.74**	0.45
	T3	96.68±0.47*	-0.03	51.90±7.27	-0.11	61.21±9.73**	0.38	49.71±5.64**	0.38
	T4	93.75±1.25*	-0.06	42.13±11.53	-0.28	73.19±13.60**	0.48	68.38±3.21**	0.55
	T5	96.68±0.47*	-0.03	30.33±5.33	-0.48	72.58±3.47**	0.48	54.60±3.65**	0.44
Spinach	CK-Z	60.83±2.49	-	-	-	-	-	-	-
	T1	28.33±1.70**	-0.53	-	-	-	-	-	-
	T2	26.17±0.47**	-0.60	-	-	-	-	-	-
	T3	20.83±1.25**	-0.66	-	-	-	-	-	-
	T4	22.50±2.94**	-0.63	-	-	-	-	-	-
	T5	11.00±1.00*	-1.21	-	-	-	-	-	-

	CK-Z	79.17±2.36	-	36.72±7.80	-	37.60±5.51	-	13.85±1.68	-
Lettuce	T1	67.50±2.16	-0.15	32.18±4.33	-0.12	47.08±5.76*	0.20	17.23±1.48*	0.20
	T2	64.17±2.87	-0.19	32.69±3.61	-0.11	42.53±3.37	0.12	20.15±1.41**	0.31
	T3	45.83±3.40*	-0.42	26.33±3.43**	-0.28	40.39±9.21	0.07	21.47±2.32**	0.35
	T4	10.83±1.86**	-0.86	13.51±4.55**	-0.63	26.39±8.82	-0.30	13.45±0.45	-0.03
	T5	7.50±0.82**	-0.91	12.69±4.87**	-0.65	14.59±16.32*	-0.61	3.28±0.65**	-0.76

Note: The value is the mean ± standard deviation, and "*" represents a significant level $P < 0.05$, "**" indicates a significant level of $P < 0.01$.

From Table 2 it can be seen that the seed GR of the three crops was reduced to varying degrees after treatment, and all the allelopathic effect indices were negative. The GR of spinach and lettuce were significantly reduced compared to the control ($P < 0.05$), except for pakchoi. The GR of spinach was extremely significantly lower than that of the control ($P < 0.01$) under each treatment of different concentrations of CCE, while the lettuce GR was extremely significantly reduced ($P < 0.01$) when the concentration of CCE reached T4. Therefore, each concentration of CCE has a different inhibitory effect on GR in the three crops.

Pakchoi and spinach show different characteristics in terms of root length, seedling height and fresh weight after treatment with CCE. The root length values of pakchoi and lettuce decreased to varying degrees after treatment with CCE, and the allelopathic effect index was negative. Among them, the root length of lettuce decreased significantly when CCE reached T3 compared to the control ($P < 0.01$). But the CCE had a different effect on the height of the seedlings in both crops.

Pakchoi has a gradual increase in seed height with increasing CCE concentration and a positive allelopathy index. When the CCE concentration reached T2, the seedling height of pakchoi was significantly different from that of the control, which indicated that the application of CCE to pakchoi had a promoting effect on the seedling height. After treatment with CCE, the characteristics of seedling height changes in lettuce are different from those in pakchoi, reflecting the characteristics of "low promotion and high inhibition". At CCE concentrations ranging from T1 to T3, CCE promotes the seedling height of lettuce, and the increase in seedling height at T1 concentration was significantly different from the control ($P < 0.05$). However, at high CCE concentrations, there was a significant decrease in seedling height of the lettuce, and there was a significant difference ($P < 0.05$) compared to the control at a concentration of T5.

After the CCE treatment, the fresh weight change properties of pakchoi and lettuce are similar to those of seedling heights. Pakchoi has a gradual increase in fresh weight with increasing CCE concentration and a positive allelopathy index. After the CCE concentration reached T2, the value of fresh weight of pakchoi was significantly different from that of the control ($P < 0.01$). The lettuce still showed the characteristics of "low promotion and high suppression" after the treatment of CCE. At the CCE concentration of T1 to T3, the fresh weight of the lettuce was significantly increased ($P < 0.05$). However, at high concentrations, the fresh weight of the lettuce decreased significantly, and the difference between T5 and the control was extremely significant ($P < 0.01$).

Comment [i-[18]: Please revise it, check space, grammar, etc.

3.2 The comprehensive allelopathic effects of different CCE on crops

Table 3. The comprehensive allelopathic effects of different CCE on crops

Test plants	Concentration (g/L)						
	CK-z	T1	T2	T3	T4	T5	
CAE	Pakchoi	-	0.07	0.09	0.16	0.17	0.10
	Lettuce	-	0.03	0.03	0.07	-0.46	-0.73

Table 3 shows the comprehensive allelopathic effects (CAE) of different concentrations of CCE on various indicators of pakchoi and lettuce. The CAE of CCE on pakchoi was positive. This suggests that CCE has an overall beneficial effect on the germination of pakchoi seeds. The CAE of lettuce was found to be positive at low concentrations and negative at high concentrations, so it was suggested that low concentrations of CCE had a positive effect on seed germination in lettuce, however, high concentrations of CCE had a negative effect on seed germination in lettuce.

4. DISCUSSION

The CCE has different degrees of allelopathic effects on seed germination and seedling growth [5]. The water extract contains phenolic allelopathic substances [6]. Phenolic substances can damage the root tip cell structure of plants, thereby affecting the absorption of nutrients by plants [7]. Studies have shown that excessive levels of phenolic substances can inhibit root growth in plants [8]. In this study, all concentrations of CCE suppressed the GR of the three crops and also affected the root growth of pakchoi and lettuce, in agreement with the above results. However, at the same time, the CCE also showed a promoting effect on the growth of crop height and fresh weight. In particular, for the pakchoi seedling height and fresh weight promotion, significant differences with the control can be achieved. The response of seedling height and fresh weight of lettuce to the CCE was characterized by "low promotion and high inhibition". The different responses of different crops to the CCE may be related to the crop itself [9-11]. For example, *Avena sativa* aqueous extract significantly inhibited root growth and promoted seedling height on seeds of *Elymus nutans* and lettuce [9,12]. The water immersion of the invasive plant (*Ambrosia trifida*.L) had a promoting effect on the plant freshness of pakchoi and lettuce [10]. The effect of CCE on the fresh weight of lettuce is consistent with the research results of Zhang Ruiqi [11] et al., which showed "low promotion, high inhibition" in the effect of herb water extract on *Amygdalus pedunculata*. In this study, the allelopathic substances of *Conyza Canadensis*, had different effects on crops, and the individual biomass of some crops tended to increase under the influence of the allelopathic substances. Therefore, in addition to some ecological restoration functions, it may have a possible ecological value in promoting crop biomass growth.

5. CONCLUSION

CCE has a varying degree of inhibitory effect on crop GR and root growth, but it can significantly promote the growth of pakchoi seedling heights and fresh weight. The seedling height and fresh weight of lettuce are promoted at low CCE concentrations and promoted at high CCE concentrations.

REFERENCES

- [1] Chen, F., Meng, Y.J., Shuai, H.W., Luo, X.F., Zhou, W.G. (2017) Effect of plant allelochemicals on seed germination and its ecological significance. *Chinese Journal of Eco-Agriculture*, 25(01): 36-46.
- [2] Ping, X.Y., Wang, T.M. (2018) Ecological significance of plant allelopathy and progress in allelopathy research in grassland ecosystems. *Acta Prataculturae Sinica*, 27(08): 175-184.
- [3] Xu, L.X., Li, Y.F., Wu, D., Zheng, J.J., Zhang, J.G. (2022) Effects of *Artemisia Argyi* Extracts on Seed Germination and Seeding Growth of Six Grasses. *Seed*, 41(10): 58-64.
- [4] Shen, S.C., Xu, G.F., Zhang, F.D., Jin, G.M., Liu, S.F., Yang, Y.X., Zhang, Y.H. (2017) Allelopathic effects of water Extracts from sweet potato (*Ipomoea batatas*) leaves on five major farming weeds. *Acta Ecologica Sinica*, 37(06): 1931-1938.
- [5] Li, W.Y., Wang, Y., Lin, Y. (2023) Allelopathic Effects of *Camellia oleifera* Abel Extracts on three medicinal plants. *Journal of Chinese Medicinal Materials*, (03): 570-575[2023-4-11].
- [6] Xu, G.F., Liu, J.M., Chao, H.J. (2007) Study on Allelopathy of the Invasive Plant *Conyza Canadensis*. *Acta Agriculturae Boreali-occidentalis Sinica*, (03): 215-218.

[7] Wang, A.K., Bi, Y.F., Wen, X., Wang, Y.K., Li, W.C. (2019) Research Advances of Plant Allelochemicals. *Molecular Plant Breeding.*, 17(17): 5829-5835.

[8] Wang, Y.C., Lü, J.J., Mao, Z.X., Li, Q. (2023) Allelopathic Effect of Native *Artemisia* Species Extracts on Growth of Invasive Weeds in Farmland. *Acta Agriculturae Boreali-occidentalis Sinica.*, 32(01): 117-129.

[9] Li, R., Mu, P., Zhao, G.Q. (2019) Allelopathic Effects of from Different Parts of Oat (*Avena sativa*) Seed on Rapeseed (*Brassica napus*). *Journal of Grassland Science and Technology.*, 27(04): 882-888.

[10] Ai, M.J., Shang, Y. H., Zhao, W.Y. (2022) Allelopathy of Invasive *Ambrosia trifida* L. *Journal of Fujian Agriculture.*, 37(10): 1326-1334.

[11] Zhang, R.Q., Wang, J.X., Wang, X.Q. (2018) Effects of the Water Extract from Four Herbs on Seed Germination and Seeding Growth of *Amygdalus pedunculata*. *Journal of Northwest Forestry University.*, 33(05): 94-99+129.

[12] Zhou, J.J., Wei, W., Xie, W.D. (2021) Allelopathic Effect of Oat (*Avena sativa*) Aqueous Extracts on Seed Germination and Seeding Growth of *Elymus nutans*. *Chinese Journal of Grassland*, 43(08): 18-25.

Comment [1-19]: Edit it.

UNDER PEER REVIEW